

RAIRO

ANALYSE NUMÉRIQUE

ROLF RANNACHER

Some asymptotic error estimates for finite element approximation of minimal surfaces

RAIRO – Analyse numérique, tome 11, n° 2 (1977), p. 181-196.

http://www.numdam.org/item?id=M2AN_1977__11_2_181_0

© AFCET, 1977, tous droits réservés.

L'accès aux archives de la revue « RAIRO – Analyse numérique » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

SOME ASYMPTOTIC ERROR ESTIMATES FOR FINITE ELEMENT APPROXIMATION OF MINIMAL SURFACES (1)

par Rolf RANNACHER (2)

Communiqué par V. THOMÉE

Abstract. — *The solution of a minimal surface problem over a plane domain is approximated by piecewise linear finite elements. Using related results of Johnson/Thomé [5] and a weighted Sobolev norm technique introduced by Nitsche [11], we show the L^∞ -convergence with rate $O(h^2 |\ln h|)$ and the L^2 -convergence with rate $O(h^2)$.*

1. INTRODUCTION

Let Ω be a bounded, strictly convex domain in the plane R^2 with smooth boundary $\partial\Omega \in C^{2,\alpha}$, $0 < \alpha < 1$, and let $g \in C^0(R^2)$ be a real function. We consider the following minimal surface problem for functions $u \in C^{0,1}(\bar{\Omega})$:

$$(V) \quad \int_{\Omega} (1 + |\nabla u|^2)^{1/2} dx = \text{Min.}, \quad u = g \text{ on } \partial\Omega.$$

If g satisfies the bounded slope condition on $\partial\Omega$ and is in the Sobolev space $W^{2,q}(\Omega)$ for some $q \in]2, \infty[$ or in the Hölder space $C^{2,\alpha}(\bar{\Omega})$, then it is known that there is a unique minimizing function

$$u \in W^{2,q}(\Omega) \quad \text{or} \quad u \in C^{2,\alpha}(\bar{\Omega}) \subset W^{2,\infty}(\Omega),$$

respectively, (see [7; 4.2.1]). This solution will be approximated by the simplest finite element method.

Let $\hat{\Omega}_h \supset \bar{\Omega}$, $0 < h \leq h_0 < 1$, be polygonal domains, and let $T_h = \{T_i\}$ be finite triangulations of $\hat{\Omega}_h$, such that the triangles have disjoint interiors and all their edges are the edge of another triangle or of the polygon $\partial\hat{\Omega}_h$. Further all vertices of the inscribed polygonal domain

$$\Omega_h := \cup \{T \in T_h \mid T \subset \bar{\Omega}\} \subset \bar{\Omega} \subset \hat{\Omega}_h$$

(1) Manuscrit reçu le 22 mars 1976.

(2) Institut für Angewandte Mathematik der Universität Bonn. (Fed. Rep. Germany).

are on the boundary $\partial\Omega$. It is assumed that the triangulations T_h are quasi-regular :

(T) Each triangle $T \in T_h$ contains a circle with radius $c_1 h$ and is contained in a circle with radius $c_2 h$.

We define finite dimensional subspaces $S_h \subset W^{1,2}(\hat{\Omega}_h)$ by

$$S_h := \{ v_h \in C(\hat{\Omega}_h) \mid v_h \text{ linear on each } T \in T_h \}$$

and $S_h^0 \subset W_0^{1,2}(\Omega_h) \subset W_0^{1,2}(\Omega)$ by

$$S_h^0 := \{ v_h \in S_h \mid v_h = 0 \text{ on } \hat{\Omega}_h - \Omega_h \}.$$

The usual interpolant $I_h g \in S_h$ of g is determined by

$$I_h g(x) = g(x) \quad \text{for each vertex } x \in T \in T_h. \tag{1}$$

Now the approximating functions $u_h \in S_h$ are defined by the finite problems

$$(V_h) \quad \int_{\Omega_h} (1 + |\nabla u_h|^2)^{1/2} dx = \text{Min.}, \quad u_h - I_h g \in S_h^0.$$

The function $F(\eta) := (1 + |\eta|^2)^{1/2}$ is strictly convex on R^2 .

Thus the existence of unique solutions of (V_h) follows from the continuity and coerciveness of the functional.

Johnson/Thomee [5] have shown for $u \in W^{2,2}(\Omega) \cap W^{1,\infty}(\Omega)$ the rate of convergence

$$\|\nabla(u - u_h)\|_2 = O(h) \tag{2}$$

and the estimate

$$\|\nabla u_h\|_\infty = O(1). \tag{3}$$

Further for $u \in W^{2,q}(\Omega)$, $q > 2$, they derived, using a common duality argument,

$$\|u - u_h\|_p = O(h^2) \quad , \quad 1 \leq p < 2. \tag{4}$$

For uniform triangulations and $u \in C^2(\bar{\Omega}) \cap W^{3,2}(\Omega)$ Mittelmann [6] has shown the L^∞ -estimate

$$\|u - u_h\|_\infty = O(h^{3/2} |\ln h|^{1/2}). \tag{5}$$

In the linear case $F(u) = |\nabla u|^2 - 2fu$ with $u \in W^{2,\infty}(\Omega)$ the analogue of the method (V_h) has the order of convergence (see [8], [13], [11], [12], [4])

$$\|u - u_h\|_\infty = O(h^2 |\ln h|). \tag{6}$$

Recently this was obtained by Frehse [3] for the general variational problem, too :

$$\int_{\Omega} F(\cdot, u, \nabla u) dx = \text{Min.}, \quad u \in W_0^{1,2}(\Omega). \tag{7}$$

He used a Morrey norm estimate

$$\|\nabla(u - u_h)\|_\alpha = O(h) \quad \text{with an } \alpha > 0, \quad (8)$$

which is proven in [2] for zero boundary conditions and uniformly convex functions $F(x, \xi, \cdot)$. These assumptions are not valid in the case of minimal surfaces. In this note we shall prove the L^∞ -estimate (6) for the nonlinear problem (V) without making use of (8). Furthermore the L^p -estimate (4) will be extended to the case $p = 2$.

THEOREM : Assume $\partial\Omega \in C^{2,\alpha}$, $0 < \alpha < 1$, and $g \in W^{2,q}(R^2) \cap W^{2,q}(\partial\Omega)$ and $u \in W^{2,q}(\Omega)$ for some q with $2 < q \leq \infty$. Further let the triangulations T_h be quasi-regular in the sense of (T). Then

$$\|u - u_h\|_\infty = O(h^{2-2/q} |\ln h|), \quad (I)$$

$$\|u - u_h\|_2 = O(h^2). \quad (II)$$

REMARKS : The methods applied in the proof also work with slight modifications for the general variational problem (7) under the assumptions of [3]. This will be of its own interest since the estimate (8) is somewhat hard to prove. For this problem the above L^∞ -estimate can be extended to higher dimensions $n \geq 3$, too. This will be carried out in a forthcoming paper.

Moreover, there is no difficulty to obtain analogously to [11] the order of pointwise convergence $O(h^m)$ for the finite element approximation of a solution $u \in W^{m,\infty}(\Omega)$ of (7) with elements of order $m \geq 3$.

Here and below $L^p(\Omega)$ and $W^{m,p}(\Omega)$, $W_0^{m,p}(\Omega)$ denote the usual real Lebesgue and Sobolev spaces with the corresponding norms

$$\|\cdot\|_p = \|\cdot\|_{p,\Omega} \quad , \quad \|\cdot\|_{m,p} = \|\cdot\|_{m,p,\Omega} \quad , \quad 1 \leq p \leq \infty \quad , \quad m \in N_0.$$

Further we use the abbreviations

$$\partial_i v := \partial v / \partial x^i \quad , \quad i = 1, 2, \quad \nabla^1 v = \nabla v := \text{grad } v \quad , \quad \nabla^2 v := (\partial_i \partial_k v)_{i,k=1,2},$$

for the partial (generalized) derivatives, and c for a positive (generic) constant which is independent of the parameters h and ρ , defined below. Finally, we shall use the usual summation convention.

2. PROOF OF THE THEOREM

First we introduce some notations and technical facts.

The minimizing functions $u \in W^{2,q}(\Omega) \subset C^1(\bar{\Omega})$, $2 < q \leq \infty$, of (V) and $u_h \in S_h$ of (V_h) necessarily satisfy the Euler equations

$$\int_{\Omega} (1 + |\nabla u|^2)^{-1/2} \nabla u \cdot \nabla v \, dx = 0, \quad \forall v \in W_0^{1,2}(\Omega),$$

$$\int_{\Omega_h} (1 + |\nabla u_h|^2)^{-1/2} \nabla u_h \cdot \nabla v_h \, dx = 0, \quad \forall v_h \in S_h^0.$$

Denoting by $F_{ik} := \partial_i \partial_k F$, $i, k = 1, 2$, the second derivatives of the function $F(\eta) := (1 + |\eta|^2)^{1/2}$, $\eta \in R^2$, we obtain

$$F_{ik}(\eta)\xi_i \xi_k \geq (1 + |\eta|^2)^{-3/2} |\xi|^2 \quad , \quad \eta, \xi \in R^2.$$

Combination of the Euler equations yields

$$a^h(v_h, u - u_h) := \int_{\Omega_h} a_{ik}^h(\cdot) \partial_i v_h \partial_k (u - u_h) dx = 0, \quad \forall v_h \in S_h^0, \quad (9)$$

with the $L^\infty(\Omega)$ -functions

$$a_{ik}^h(\cdot) := \int_0^1 F_{ik}(\nabla u_h(\cdot) + t \nabla(u - u_h)(\cdot)) dt, \quad i, k = 1, 2.$$

Further by the result (3),

$$a_{ik}^h \xi_i \xi_k \geq c |\xi|^2 \quad , \quad \xi \in R^2 \quad , \quad \text{on } \bar{\Omega}.$$

Since the a_{ik}^h are discontinuous we introduce the bilinear form

$$a(v, w) := \int_{\Omega} a_{ik}(\cdot) \partial_i v \partial_k w dx, \quad v, w \in W^{1,2}(\Omega),$$

with coefficients

$$a_{ik}(\cdot) := F_{ik}(\nabla u(\cdot)) \in W^{1,q}(\Omega) \quad , \quad i, k = 1, 2.$$

Then the differential operator

$$A := - \partial_k \{ a_{ik}(\cdot) \partial_i \} \quad (11)$$

is uniformly elliptic in Ω and hence satisfies the well known a priori estimate (see [7; 5.2 ff]) :

$$\|v\|_{2,2} \leq c \|Av\|_2 \quad , \quad \forall v \in W_0^{1,2}(\Omega) \cap W^{2,2}(\Omega).$$

From the boundedness of the derivatives of F_{ik} it follows that

$$|a_{ik} - a_{ik}^h| \leq c |\nabla(u - u_h)| \quad \text{on } \Omega. \quad (12)$$

For functions $v \in W^{2,p}(\Omega)$, $2 \leq p \leq \infty$, and $w \in W_0^{1,2}(\Omega) \cap W^{2,2}(\Omega)$ and the corresponding interpolants $I_h v \in S_h$ and $I_h w \in S_h^0$ the following estimates are known (see [1] and [10]) :

$$\|v - I_h v\|_{j,p,\Omega_h} \leq ch^{2-j} \|\nabla^2 v\|_{p,\Omega_h} \quad , \quad 0 \leq j < 2 \quad , \quad 2 \leq p \leq \infty. \quad (13)$$

$$\|v - I_h v\|_{j,\infty,\Omega_h} \leq ch^{2-j-2/p} \|\nabla^2 v\|_{p,\Omega_h}$$

Using in addition Lemma A4 of the Appendix we find with (15)

$$\|w - I_h w\|_{j,2} \leq ch^{2-j} \|w\|_{2,2}, \quad 0 \leq j < 2. \quad (14)$$

Proof of Proposition (I)

In order to make the outline of the proof clear we write its main steps as lemmas.

Set $e_h := u - u_h$ and let $E_h := I_h e_h = I_h u - u_h$ be its interpolant on Ω_h . Observe that $E_h \in S_h^0$.

First we estimate e_h on $\Omega - \Omega_h$:

The assumptions (T) and $\partial\Omega \in C^2$ imply

$$d(\partial\Omega_h, \partial\Omega) := \sup_{x \in \partial\Omega_h} \text{dist}(x, \partial\Omega) \leq ch^2. \quad (15)$$

Thus, we conclude using $u = g$ on $\partial\Omega$, $u_h = I_h g$ on $\Omega - \Omega_h$ and the estimate (13)

$$\begin{aligned} \|e_h\|_{\infty; \Omega - \Omega_h} &\leq \|u - g\|_{\infty; \Omega - \Omega_h} + \|g - I_h g\|_{\infty; \Omega - \Omega_h} \\ &\leq ch^{2-2/q} (\|u\|_{2,q} + \|g\|_{2,q, \mathcal{R}^2}). \end{aligned}$$

Next, let $z_h \in \Omega_h$ be points with the properties

$$|E_h(z_h)| = \|E_h\|_{\infty}.$$

Then, with any disk $B := B_{\tau}(z_h)$, $\tau > 0$,

$$\|E_h\|_{\infty} \leq c\tau^{-2} \int_B |E_h| dx + c\tau \|\nabla E_h\|_{\infty}.$$

Using the well known inverse relation $\|\nabla E_h\|_{\infty} \leq ch^{-1} \|E_h\|_{\infty}$, we find that for $\tau := \delta h$, $\delta > 0$ a constant sufficiently small,

$$\|E_h\|_{\infty} \leq ch^{-2} \int_B |E_h| dx.$$

Thus by the estimate (13)

$$\begin{aligned} \|e_h\|_{\infty; \Omega_h} &\leq \|u - I_h u\|_{\infty; \Omega_h} + \|E_h\|_{\infty} \\ &\leq ch^{2-2/q} \|u\|_{2,q} + ch^{-2} \int_B |e_h| dx. \end{aligned} \quad (17)$$

In the following our main tool will be a modification of the weighted norm technique by Nitsche [11]. With a real parameter $0 < \rho \leq \rho_0$, which will be appropriately coupled with h below, we define the weight function

$$\sigma_h(\cdot) := (|\cdot - z_h|^2 + \rho^2)^{1/2}$$

and the weighted norms (T denoting triangles of T_h)

$$\|\cdot\|_{(v)} := \left(\sum_{T \in \Omega_h} \int_T \sigma_h^v |\cdot|^2 dx \right)^{1/2}, v \in R. \tag{18}$$

Since the points z_h and the corresponding disks $B := B_{\delta h}(z_h)$ will be fixed during the proof, we shall omit the index of σ .

Obviously with constants independent of ρ

$$|\nabla\sigma| \leq c, \quad |\nabla^2\sigma| \leq c\sigma^{-1} \leq c\rho^{-1} \quad \text{on } R^2,$$

and for $\rho \geq c_3 h$, c_3 sufficiently large,

$$\max_{T \in T_h} \{ \max_{x \in T} \sigma^v(x) / \min_{x \in T} \sigma^v(x) \} \leq c, \quad -4 \leq v \leq 4.$$

From this and (13) we conclude the following interpolation estimate for functions $v \in C(\Omega_h) \cap \bigoplus_{T \in \Omega_h} W^{2,2}(T)$ (see [11]) :

$$\|v - I_h v\|_{(v)} + h \|\nabla(v - I_h v)\|_{(v)} \leq ch^2 \|\nabla^2 v\|_{(v)}, \quad -4 \leq v \leq 4. \tag{19}$$

LEMMA 1 : *Let $1 < \beta < 2$. Then there are constants c_4, c_β , independent of h and $\rho \geq c_3 h$, so that*

$$\|e_h\|_{\infty; \Omega_h} \leq c_4 h^{2-2/q} |\ln h| + c_\beta \rho h^{-1} |\ln h| \|\nabla e_h\|_{(-\beta)}^2.$$

Proof. We shall estimate the integral in (17) making use of a common duality argument. Let $G^h \in W_0^{1,2}(\Omega) \cap W^{2,2}(\Omega)$ be the solution of the boundary value problem (smoothed Green function)

$$AG^h - h^{-2} \operatorname{sgn}(e_h) \chi_B \quad \text{in } \Omega, \quad G^h = 0 \quad \text{on } \partial\Omega, \tag{20}$$

with χ_B the characteristic function of the disk B , and let $G_h^h \in S_h^0$ be its Ritz projections defined by

$$a(v_h, G_h^h) = a(v_h, G^h), \quad \forall v_h \in S_h^0. \tag{21}$$

With this setting we obtain by integration by parts denoting

$$\partial_n := -n_k a_{ik} \partial_i, \quad (n_1, n_2) := \text{outward normal to } \partial\Omega,$$

$$h^{-2} \int_B |e_h| dx = \int_{\partial\Omega} e_h \partial_n G^h ds + a(e_h, G^h) \tag{22}$$

and by (9)

$$h^{-2} \int_B |e_h| dx = \int_{\partial\Omega} e_h \partial_n G^h ds + a(e_h, G^h - G_h^h) + (a - a^h)(e_h, G_h^h). \tag{23}$$

Now we shall estimate the three terms on the right.

Since $e_h = g - I_h g$ on $\partial\Omega$, we find using Hölder's inequality and a well known trace theorem (see [9])

$$\left| \int_{\partial\Omega} e_h \partial_n G^h ds \right| \leq c \|g - I_h g\|_{q;\partial\Omega} \|G^h\|_{2,r}, \quad 1/r = 1 - 1/(2q) \quad (24)$$

For $g \in W^{2,q}(R^2) \cap W^{2,q}(\partial\Omega)$ it is proven in [5] that

$$\|g - I_h g\|_{q;\partial\Omega} \leq ch^2.$$

Using Hölder's inequality (notice $r = 2q/(2q - 1) < 2$), Poincaré's inequality for $G^h \in W_0^{1,2}(\Omega)$ and Lemma A2(a) with $\rho \geq c_3 h$, we find (We note that the logarithm only appears for $r = 1$.)

$$\|G^h\|_{2,r} \leq ch^{-1/q} |\ln h|^{1/2} \{ \|\nabla G^h\|_2 + \|\nabla^2 G^h\|_{(2)} \} \leq ch^{-1/q} |\ln h|. \quad (25)$$

Thus

$$\left| \int_{\partial\Omega} e_h \partial_n G^h ds \right| \leq ch^{2-1/q} |\ln h|. \quad (26)$$

The second term in (23) is the same that occurs in the case of linear problems (see [4]). The modified interpolate of u , defined by

$$\hat{I}_h u := \begin{cases} I_h u & \text{on } \Omega_h, \\ I_h g & \text{on } \hat{\Omega}_h - \Omega_h, \end{cases}$$

obviously satisfies $\hat{I}_h u - u_h \in S_h^0$. Hence, we find using (21)

$$\begin{aligned} |a(e_h, G^h - G_h^h)| &= |a(u - \hat{I}_h u, G^h - G_h^h)| \\ &\leq c \|\nabla(u - I_h u)\|_{\infty;\Omega_h} \|\nabla(G^h - G_h^h)\|_1 \\ &\quad + c(\|u\|_{2,q} + \|g\|_{2,q;R^2}) \times \int_{\Omega - \Omega_h} |\nabla G^h| dx. \end{aligned}$$

By Lemma A4 (notice (15)) and Lemma A2(a) (with $\varepsilon = 0, \rho \geq c_3 h$) it follows in the same way as in (25) with $r = 1$

$$\int_{\Omega - \Omega_h} |\nabla G^h| dx \leq ch^2 \|G^h\|_{2,1} \leq ch^2 |\ln h|.$$

Thus, we conclude using the estimate (13) and the result, stated in Lemma A2(b), concerning the convergence of Green functions :

$$|a(e_h, G^h - G_h^h)| \leq ch^{2-2/q} |\ln h|. \quad (27)$$

The third term in (23) comes from the nonlinearity of the problem (V), i.e. from the replacement of the discontinuous coefficients a_{ik}^h by $a_{ik} \in W^{1,q}(\Omega)$. By making use of (12) and Lemma A2(c) (with $\rho \geq c_3 h$) we see (notice $G_h^h = 0$ on $\Omega - \Omega_h$)

$$\begin{aligned} |(a - a^h)(e_h, G_h^h)| &\leq c \int_{\Omega_h} |\nabla e_h|^2 \sigma^{-\beta} |\sigma^\beta \nabla G_h^h| dx \\ &\leq c_\beta \rho h^{-1} |\ln h| \|\nabla e_h\|_{(-\beta)}^2. \end{aligned}$$

Together with (27) and (26) this establishes the desired estimate.

Since $\beta < 2$ our proposition (I) suggests the estimate $\|\nabla e_h\|_{(-\beta)} = O(h^{1-1/q})$ without any logarithmic term. Obviously this would complete the proof of (I). A first step in this direction will be the following :

LEMMA 2 : *Let $1 < \beta < 2$. Then there are constants c_6, c_β , independent of h , so that for $\rho = c_6 h |\ln h|^{3/2}$*

$$\|\nabla e_h\|_{(-\beta)}^2 \leq c_\beta h^{3-\beta-2/q} |\ln h|.$$

Proof. The result (2) gives for $\rho \geq h$

$$\|\nabla e_h\|_{(-\beta)}^2 \leq c \|\nabla e_h\|_{(-2)} \|\nabla e_h\|_2 h^{1-\beta} \leq c h^{2-\beta} \|\nabla e_h\|_{(-2)}. \quad (28)$$

The weighted norm on the right will be estimated in the same way as in [11] and [12]. By (10) we have

$$\begin{aligned} \|\nabla e_h\|_{(-2)}^2 &\leq c |a^h(e_h, \sigma^{-2} e_h)| + c \int_{\Omega_h} |\nabla e_h| |e_h| |\nabla \sigma^{-2}| dx \\ &\leq c |a^h(e_h, \sigma^{-2} e_h)| + c \|\nabla e_h\|_{(-2)} \|e_h\|_{(-4)}. \end{aligned}$$

Since $v_h := I_h(\sigma^{-2} E_h) \in S_h^0$, $E_h := I_h u - u_h$, it follows by (9) that

$$\begin{aligned} |a^h(e_h, \sigma^{-2} e_h)| &= |a^h(e_h, \sigma^{-2}(u - I_h u) + \sigma^{-2} E_h - v_h)| \\ &\leq c \|\nabla e_h\|_{(-2)} \{ \|\nabla(\sigma^{-2}(u - I_h u))\|_{(2)} + \|\nabla(\sigma^{-2} E_h - v_h)\|_{(2)} \}. \end{aligned}$$

Using the estimate (19), we get for $\rho \geq c_3 h$ (notice $\nabla^2 E_h = 0$ on each $T \in T_h$)

$$\begin{aligned} \|\nabla(\sigma^{-2}(u - I_h u))\|_{(2)} &\leq c \{ \|u - I_h u\|_{(-4)} + \|\nabla(u - I_h u)\|_{(-2)} \} \\ &\leq c h \|\nabla^2 u\|_{(-2)} \end{aligned}$$

and

$$\begin{aligned} \|\nabla(\sigma^{-2} E_h - v_h)\|_{(2)} &\leq c h \|\nabla^2(\sigma^{-2} E_h)\|_{(2)} \\ &\leq c h \{ \|e_h\|_{(-6)} + \|\nabla e_h\|_{(-4)} + \|u - I_h u\|_{(-6)} + \|\nabla(u - I_h u)\|_{(-4)} \}. \\ &\leq c \{ \|e_h\|_{(-4)} + h \rho^{-1} \|\nabla e_h\|_{(-2)} + h \|\nabla^2 u\|_{(-2)} \}. \end{aligned}$$

Thus

$$\|\nabla e_h\|_{(-2)} \leq c \|e_h\|_{(-4)} + ch\rho^{-1} \|\nabla e_h\|_{(-2)} + ch \|\nabla^2 u\|_{(-2)}, \quad (29)$$

and with $\rho \geq c_5 h$, c_5 sufficiently large,

$$\|\nabla e_h\|_{(-2)} \leq c \|e_h\|_{(-4)} + ch^{1-2/q} |\ln h|^{1/2} \|u\|_{2,q}. \quad (30)$$

In order to estimate the first term on the right we use a duality argument. With the solution $v \in W_0^{1,2}(\Omega) \cap W^{2,2}(\Omega)$ of the boundary value problem

$$Av = \sigma^{-4} e_h \quad \text{in } \Omega, \quad v = 0 \quad \text{on } \partial\Omega,$$

we obtain by integration by parts

$$\|e_h\|_{(-4)}^2 = \int_{\partial\Omega} e_h \partial_n v \, ds + a(e_h, v)$$

and in addition with $I_h v \in S_h^0$ by (9) (see (23))

$$\|e_h\|_{(-4)}^2 = \int_{\partial\Omega} e_h \partial_n v \, ds + a(e_h, v - I_h v) + (a - a^h)(e_h, I_h v). \quad (31)$$

The three terms on the right will be estimated analogously to those in (23). With $1/r = 1 - 1/(2q)$ we get

$$\begin{aligned} \left| \int_{\partial\Omega} e_h \partial_n v \, ds \right| &\leq c \|g - I_h g\|_{q;\partial\Omega} \|v\|_{2,r} \\ &\leq ch^{2-1/q} |\ln h|^{1/2} \{ \|\nabla v\|_2 + \|\nabla^2 v\|_{(2)} \}. \end{aligned}$$

From the estimate (19) and Lemma A4 it follows that

$$\begin{aligned} |a(e_h, v - I_h v)| &\leq c \|\nabla e_h\|_{(-2)} \|\nabla(v - I_h v)\|_{(2)} + c \|\nabla(u - I_h g)\|_\infty \int_{\Omega - \Omega_h} |\nabla u| \, dx \\ &\leq ch \|\nabla e_h\|_{(-2)} \|\nabla^2 v\|_{(2)} + ch^2 \|v\|_{2,1} \\ &\leq c \{ h \|\nabla e_h\|_{(-2)} + h^2 |\ln h|^{1/2} \} \{ \|\nabla v\|_2 + \|\nabla^2 v\|_{(2)} \}. \end{aligned}$$

Using the estimate (12), Lemma A1 (with $v = 2$) and the result (2), we find

$$\begin{aligned} |(a - a^h)(e_h, I_h v)| &\leq c \|\nabla e_h\|_{(-1)}^2 |\ln h| \{ \|\nabla v\|_2 + \|\nabla^2 v\|_{(2)} \} \\ &\leq ch |\ln h| \|\nabla e_h\|_{(-2)} \{ \|\nabla v\|_2 + \|\nabla^2 v\|_{(2)} \}. \end{aligned}$$

Thus, by Lemma A3, observing that $\|Av\|_{(4)} = \|e_h\|_{(-4)}$,

$$\|e_h\|_{(-4)} \leq ch^{1-1/q} |\ln h| + ch\rho^{-1} |\ln h|^{3/2} \|\nabla e_h\|_{(-2)}.$$

We substitute this in (30), choose $\rho = c_6 h |\ln h|^{3/2}$, c_6 appropriately large, and obtain

$$\|\nabla e_h\|_{(-2)} \leq ch^{1-2/q} |\ln h|. \quad (32)$$

From this the desired estimate follows.

By combination of Lemma 1 and Lemma 2, we find as a first result for $1 < \beta < 2$

$$\|e_h\|_{\infty; \Omega_h} \leq c_\beta h^{3-\beta-2/q} |\ln h|^4. \tag{33}$$

Now, this will be used to improve the estimate of Lemma 2.

LEMMA 3 : Let $1 < \beta_q < 1 + (q - 2)/(3q)$. Then there are constants c_7, c_8 independent of h , so that for $\rho = c_7 h$

$$\|\nabla e_h\|_{(-\beta_q)} \leq c_8 h^{1-1/q}.$$

Proof. Set $\rho \geq c_3 h$ to guarantee the interpolation estimate (19). We start in the same way as in the proof of Lemma 2 concerning the term $\|\nabla e_h\|_{(-2)}$. With $v_h := I_h(\sigma^{-\beta} E_h) \in S_h^0, E_h := I_h u - u_h$, we find analogously to (29)

$$\|\nabla e_h\|_{(-\beta)} \leq c \|e_h\|_{(-\beta-2)} + ch\rho^{-1} \|\nabla e_h\|_{(-\beta)} + ch \|\nabla^2 u\|_{(-\beta)}$$

and for $\rho = c_7 h, c_7$ appropriately large,

$$\|\nabla e_h\|_{(-\beta)} \leq c \|e_h\|_{(-\beta-2)} + ch \|\nabla^2 u\|_{(-\beta)}.$$

Thus, by the result (33)

$$\begin{aligned} \|\nabla e_h\|_{(-\beta)} &\leq ch^{3-\beta-2/q} |\ln h|^4 \|1\|_{(-\beta-2)} + ch^{2-\beta/2-2/q} \|u\|_{2,q} \\ &\leq ch^{1-1/q} \{ h^{2-3\beta/2-1/q} |\ln h|^4 + h^{1-\beta/2-1/q} \}. \end{aligned}$$

Since $q > 2$, we can find some β_q with $1 < \beta_q < 1 + (q - 2)/(3q) < 2$, so that

$$h^{(2-3\beta_q/2-1/q)} |\ln h|^4 + h^{(1-\beta_q/2-1/q)} \leq c.$$

This proves Lemma 3.

Finally, combination of Lemma 3 and Lemma 1 completes the proof of proposition (I).

Proof of Proposition (II)

The proof of Proposition (II) makes use of the estimate (I) for the given $q > 2 (e_h := u - u_h)$:

$$\|e_h\|_{\infty} \leq ch^{2-2/q} |\ln h| \leq c_q h. \tag{36}$$

Let $v \in W_0^{1,2}(\Omega) \cap W^{2,2}(\Omega)$ be the solution of the problem

$$Av = e_h \quad \text{in } \Omega, \quad v = 0 \quad \text{on } \partial\Omega. \tag{37}$$

Using (9), we find analogously to (31) with $I_h v \in S_h^0$

$$\|e_h\|_2^2 = \int_{\partial\Omega} e_h \partial_n v \, ds + a(e_h, v - I_h v) + (a - a^h)(e_h, I_h v). \tag{38}$$

The three terms on the right can be estimated in a similar way as those in (31). We get

$$\left| \int_{\partial\Omega} e_h \partial_n v \, ds \right| \leq c \|g - I_h g\|_{2; \partial\Omega} \|v\|_{2,2} \leq ch^2 \|v\|_{2,2}$$

and, using (14) and the result (2),

$$|a(e_h, v - I_h v)| \leq c \|\nabla e_h\|_2 \|\nabla(v - I_h v)\|_2 \leq ch^2 \|v\|_{2,2}.$$

Further

$$|(a - a^h)(e_h, I_h v)| \leq c \sum_{i,k} \int_{\Omega_h} |a_{ik} - a_{ik}^h| |\nabla e_h| |\nabla I_h v| \, dx.$$

It will be convenient to replace the discontinuous function $|\nabla I_h v|$ by a continuous approximant ${}_h := I_h |\nabla v|$ of $|\nabla v| \in W^1(\Omega)$ on Ω_h . By the boundedness of a_{ik} , a_{ik}^h and the estimate (12), we have

$$|(a - a^h)(e_h, I_h v)| \leq c \int_{\Omega_h} \{ |\nabla e_h| |\nabla(I_h v - v)| + |\nabla e_h| | |\nabla v| - \psi_h | + |\nabla e_h|^2 \psi_h \} \, dx$$

and, using (13) and the result (2),

$$\begin{aligned} |(a - a^h)(e_h, I_h v)| &\leq c \|\nabla e_h\|_2 h \|v\|_{2,2} + c \int_{\Omega_h} |\nabla e_h|^2 \psi_h \, dx \\ &\leq ch^2 \|v\|_{2,2} + c \int_{\Omega_h} |\nabla e_h|^2 \psi_h \, dx. \end{aligned}$$

The integral on the right can be estimated in the same way as the term $\|\nabla e_h\|_{(-2)}$ in the proof of Lemma 2. Defining $v_h := I_h(\psi_h E_h) \in S_h^0$, $E_h := I_h u - u_h$, we have by (9)

$$\begin{aligned} \left| \int_{\Omega_h} |\nabla e_h|^2 \psi_h \, dx \right| &\leq c |a^h(e_h, \psi_h(u - I_h u) + \psi_h E_h - v_h)| \\ &\quad + c \int_{\Omega_h} |\nabla e_h| |e_h| |\nabla \psi_h| \, dx \end{aligned}$$

and hence

$$\begin{aligned} \int_{\Omega_h} |\nabla e_h|^2 \psi_h \, dx &\leq c \|\nabla e_h\|_2 \{ \|\nabla \psi_h(u - I_h u)\|_{2;\Omega_h} + \|\psi_h \nabla(u - I_h u)\|_{2;\Omega_h} \\ &\quad + \|\nabla(\psi_h E_h - v_h)\|_{2;\Omega_h} + \|\nabla \psi_h e_h\|_{2;\Omega_h} \} \\ &\leq c \|\nabla e_h\|_2 \{ \text{(i)} + \text{(ii)} + \text{(iii)} + \text{(iv)} \}. \end{aligned}$$

Using Hölder’s inequality, the Sobolev embedding theorem and (13), it follows that

$$\begin{aligned} \text{(i)} \quad &\leq c \|u - I_h u\|_{\infty, \Omega_h} \|\nabla \psi_h\|_{2, \Omega_h} \leq ch^{2-2/q} \|u\|_{2, q} \|v\|_{2, 2}, \\ \text{(ii)} \quad &\leq c \|\nabla(u - I_h u)\|_{q; \Omega_h} \|\psi_h\|_{1; 2, \Omega_h} \leq ch \|u\|_{2, q} \|v\|_{2, 2}. \end{aligned}$$

Observing $\psi_h \geq 0$, $\nabla^2 \psi_h = \nabla^2 E_h = 0$ on each $T \in T_h$ and the result (3), we find

$$\begin{aligned} \text{(iii)} \quad &\leq ch \left(\sum_{T \subset \Omega_h} \int_T |\nabla^2(\psi_h E_h)|^2 dx \right)^{1/2} \leq ch \|\nabla \psi_h\|_{2, \Omega_h} \|\nabla E_h\|_{\infty} \\ &\leq ch \|u\|_{2, q} \|v\|_{2, 2}. \end{aligned}$$

Finally, our result (36) applies to the crucial term (iv)

$$\text{(iv)} \quad \leq \|\nabla \psi_h\|_{2, \Omega_h} \|e_h\|_{\infty, \Omega_h} \leq ch \|v\|_{2, 2}.$$

This gives

$$|(a - a^h)(e_h, I_h v)| \leq ch \|v\|_{2, 2} \|\nabla e_h\|_2,$$

Thus, by $\|v\|_{2, 2} \leq c \|e_h\|_2$ and the result (2),

$$\|e_h\|_2 = O(h^2).$$

This completes the proof of the theorem.

3. APPENDIX

Here we state some lemmas used in the proof of the theorem. Assume the condition (T) to be satisfied.

LEMMA A1 : Let $v \in W_0^{1,2}(\Omega) \cap W^{2,2}(\Omega)$ and let $I_h v \in S_h^0$ be its interpolant. Then

$$\|\nabla I_h v\|_{\infty} \leq c |\ln h| \|v\|_{2, 2}, \tag{a}$$

and with the weighted norms (18) for $0 \leq \nu \leq 4$ and $\rho \geq c_3 h$

$$\|\sigma^{\nu/2} \nabla I_h v\|_{\infty} \leq c |\ln h| (\|\nabla v\|_{(\nu-2)} + \|\nabla^2 v\|_{(\nu)}). \tag{b}$$

Proof. We shall prove (b). The proof of (a) is similar.

Let $T \in T_h$ be any triangle with $T \subset \Omega_h$ and let $\xi \in T$ be the center of the inscribed circle with radius $c_1 h$ (assumption (T)). The boundary $\partial\Omega$ is of class C^2 and hence satisfies a strong cone condition. The corresponding spherical cone $K := K(\xi, \tau) \subset B(\xi, \tau)$ with vertex ξ , opening $|\Sigma_K|$ and height

$\tau > 0$ (independent of $\xi \in \Omega$ and h) can be cut off to a cone $K_T \subset T$ with volume $|K_T| = ch^2$ and $K_T \subset \subset B(\xi, \tau)$. Then by (19)

$$\begin{aligned} |\sigma^{v/2} \nabla I_h v(\xi)| &\leq ch^{-2} \int_{K_T} \sigma^{v/2} |\nabla I_h v| dx \\ &\leq ch^{-1} \|\nabla(v - I_h v)\|_{(v)} + ch^{-2} \int_{K_T} \sigma^{v/2} |\nabla v| dx \\ &\leq c \|\nabla^2 v\|_{(v)} + ch^{-1} \left(\int_{K_T} \sigma^v |\nabla v|^2 dx \right)^{1/2}. \end{aligned}$$

Now choose a function $\varphi \in C^\infty$ with support in the ball $B(\xi, \tau)$ and the properties (independent of h)

$$0 \leq \varphi \leq 1, \quad |\nabla \varphi| \leq c, \quad \varphi = 1 \quad \text{on } K_T.$$

Using polar coordinates (r, θ) centered in ξ we find with the function

$$p(|x - \xi|) := (|x - \xi|^2 + h^2)^{-1}$$

$$h^{-2} \int_{K_T} \sigma^v |\nabla v|^2 dx \leq c \int_{\Sigma_K} \int_0^\tau r p(r) |\varphi \sigma^{v/2} \nabla v|^2 dr d\theta.$$

According to the special choice of φ it follows by integration by parts

$$\int_0^\tau r p(r) |\varphi \sigma^{v/2} \nabla v|^2 dr = -2 \int_0^\tau \left\{ \int_0^r s p(s) ds \right\} \varphi \sigma^{v/2} |\nabla v| \partial_r (\varphi \sigma^{v/2} |\nabla v|) dr$$

and, using the inequality $ab \leq \varepsilon a^2 + (4\varepsilon)^{-1} b^2$, $\varepsilon > 0$,

$$\int_0^\tau r p(r) |\varphi \sigma^{v/2} \nabla v|^2 dr \leq 4 \int_0^\tau \left\{ \int_0^r s p(s) ds \right\}^2 r^{-2} p(r)^{-1} r |\nabla(\varphi \sigma^{v/2} |\nabla v|)|^2 dr.$$

The function $\left\{ \int_0^r s p(s) ds \right\}^2 r^{-2} p(r)^{-1}$ is continuous and nondecreasing

for $0 \leq r < h$ and hence uniformly bounded by $c |\ln h|^2$ for $r \leq \text{diam}(\Omega)$. This gives

$$h^{-2} \int_{K_T} \sigma^v |\nabla v|^2 dx \leq c |\ln h|^2 (\|\nabla v\|_{(v-2)} + \|\nabla^2 v\|_{(v)})^2$$

and completes the proof.

LEMMA A2 : Let $G^h \in W_0^{1,2}(\Omega) \cap W^{2,2}(\Omega)$ be the smoothed Green functions defined by (20) and let G_h^h be its Ritz projections defined by (21). Then with the weighted norms (18) for $\rho \geq c_3 h$

$$\|\nabla G^h\|_{(e)} + \|\nabla^2 G^h\|_{(2+\varepsilon)} \leq c_\varepsilon \rho h^{-1} \begin{cases} 1 & , \varepsilon > 0 \\ |\ln h|^{1/2} & , \varepsilon = 0, \end{cases} \tag{a}$$

$$\|\nabla(G^h - G_h^h)\|_1 \leq c |\ln h|^{1/2} \|\nabla(G^h - G_h^h)\|_{(2)} \leq c \rho |\ln h|, \tag{b}$$

$$\|\sigma^{1+\varepsilon} \nabla G_h^h\|_\infty \leq c_\varepsilon \rho h^{-1} |\ln h|, \quad \varepsilon > 0. \tag{c}$$

Proof. The estimate (a) and (b) are proven in [4].

Analogously to Lemma A1 we find

$$|\sigma^{1+\varepsilon} \nabla G_h^h(\xi)| \leq c h^{-1} \|\nabla(G_h^h - G^h)\|_{(2+2\varepsilon)} + c |\ln h| \left\{ \|\nabla G^h\|_{(2\varepsilon)} + \|\nabla^2 G^h\|_{(2+2\varepsilon)} \right\}$$

and, using (a) and (b),

$$|\sigma^{1+\varepsilon} \nabla G_h^h(\xi)| \leq c \rho h^{-1} |\ln h|^{1/2} + c_\varepsilon \rho h^{-1} |\ln h|.$$

This proves (c).

LEMMA A3 : Let $v \in W_0^{1,2}(\Omega) \cap W^{2,2}(\tilde{\Omega})$ and let A be the uniformly elliptic differential operator defined by (11). Then with the weighted norms (18) for $0 < \rho < \rho_0$

$$\|\nabla v\|_2 + \|\nabla^2 v\|_{(2)} \leq c \rho^{-1} (1 + |\ln \rho|^{1/2}) \|Av\|_{(4)}.$$

For the sake of completeness we shall sketch a proof of this important a priori estimate. It rests on ideas contained in [12] and [4]. A similar assertion is stated by Nitsche [11] for the Laplace operator.

Proof. Set $\sigma(\cdot) := (|\cdot - z|^2 + \rho^2)^{1/2}$, $0 < \rho \leq \rho_0$, $z \in \Omega$.

Denoting by $y^j := x^j - z^j$, $j = 1, 2$, the components of the vector $x - z$, $x \in \Omega$, we get

$$\|\nabla^2 v\|_{(2)}^2 = \sum_{j=1}^2 \|y^j \nabla^2 v\|_2^2 + \rho^2 \|\nabla^2 v\|_2^2,$$

and by the well known L^2 -estimate,

$$\rho \|\nabla^2 v\|_2 \leq c \rho \|Av\|_2 \leq c \rho^{-1} \|Av\|_{(4)},$$

$$\|y^j \nabla^2 v\|_2 \leq c \{ \|\nabla^2(y^j v)\|_2 + \|\nabla v\|_2 \} \leq c \{ \|A(y^j v)\|_2 + \|\nabla v\|_2 \}.$$

Further, using $M := \max_{i,k} \|a_{ik}\|_{1,q} < \infty$ and $\max_{i,k} \|a_{ik}\|_\infty \leq cM$, we find with $p := 2q/(q - 2)$

$$\|A(y^j v)\|_2 \leq c \{ \|y^j Av\|_2 + M \|v\|_p + M \|\nabla v\|_2 \} \leq c \{ \|y^j Av\|_2 + \|v\|_{1,2} \},$$

and thus by Poincaré's inequality and $\sigma^{-1} \leq c \rho^{-1}$

$$\|\nabla^2 v\|_{(2)} \leq c \{ \|Av\|_{(2)} + \|v\|_{1,2} \} \leq c \{ \rho^{-1} \|Av\|_{(4)} + \|\nabla v\|_2 \}. \tag{39}$$

By the inequality $ab \leq (a^2 + b^2)/2$, it follows that

$$\|\nabla v\|_2^2 \leq a(v, v) \leq c\rho^{-2} |\ln \rho| \|Av\|_{(4)}^2 + c\rho^2 |\ln \rho|^{-1} \|v\|_{(-4)}^2.$$

Denoting by $g(\cdot, \cdot)$ the Green function of A over Ω we obtain

$$\|v\|_{(-4)}^2 = \int \sigma^{-4}(x) \left| \int Av(\xi)g(x, \xi) d\xi \right|^2 dx,$$

and by Hölder's inequality and an interchange of the order of integration

$$\|v\|_{(-4)}^2 \leq \int \sigma^4(\xi) |Av(\xi)|^2 \left(\int \sigma^{-4}(x)g(x, \xi) \left\{ \int \sigma^{-4}(\eta)g(x, \eta) d\eta \right\} dx \right) d\xi.$$

It is well known that the Green function g can be estimated on Ω by (see [4])

$$0 \leq g(x, y) \leq c(1 + |\ln |x - y||).$$

From this we conclude

$$\int \sigma^{-4}(\eta)g(x, \eta) d\eta \leq c\rho^{-2}(1 + |\ln \rho|).$$

It follows that

$$\|v\|_{(-4)}^2 \leq c\rho^{-4}(1 + |\ln \rho|)^2 \|Av\|_{(4)}^2,$$

and thus

$$\|\nabla v\|_2 \leq c\rho^{-1}(1 + |\ln \rho|) \|Av\|_{(4)}.$$

Together with (39) this completes the proof.

Finally, we state a simple boundary estimate, which can be proven by locally reduction to one dimensional integrations.

LEMMA A4. *Let $v \in W^{2,p}(\Omega)$, $1 \leq p < \infty$. Then (see (15))*

$$\int_{\Omega - \Omega_h} |v|^p dx \leq c_p d(\partial\Omega, \partial\Omega_h) \int_{\Omega} \{ |v|^p + |\nabla v|^p \} dx.$$

REFERENCES

1. J. H. BRAMBLE, S. R. HILBERT, *Bounds on a class of linear functionals with applications to Hermite interpolation*, Numer. Math., Vol. 16, 1971, pp. 362-369.
2. J. FREHSE, *Eine a-priori-Abschätzung zur Methode der finiten Elemente in der numerischen Variationsrechnung*. In : Numerische Behandlung von Variations- und Steuerungsproblem, Tagungsband, Bonn. Math. Schr., Vol. 77, 1975, pp. 115-126.
3. J. FREHSE, *Eine gleichmäßige asymptotische Fehlerabschätzung zur Methode der finiten Elemente bei quasilinearen Randwertproblemen*. In : Theory of Nonlinear Operators. Constructive Aspects, Tagungsband der Akademie der Wissenschaften, Berlin (DDR), 1976.

- 4 J FREHSE, R RANNACHER, *Eine L^1 -Fehlerabschätzung für diskrete Grundlösungen in der Methode der finiten Elemente* In Finite Elemente, Tagungsband, Bonn Math Schr, Vol 89, 1976, pp 92-114
- 5 C JOHNSON, V THOMEE, *Error estimates for a finite element approximation of a minimal surface*, Math Comp, Vol 29, 1975, pp 343-349
- 6 H D MITTELMANN, *On pointwise estimates for a finite element solution of nonlinear boundary value problems* To appear
- 7 C B MORREY, *Multiple integrals in the calculus of variations* Springer Berlin-Heidelberg-New York, 1966
- 8 F NATTERER, *Über die punktweise Konvergenz finiter Elemente* Numer Math, Vol 25 1975 pp 67-77
- 9 J NECAS, *Les methodes directes en theorie des equations elliptiques* Masson, Paris, 1967
- 10 J NITSCHKE, *Lineare Spline-Funktionen und die Methode von Ritz für elliptische Randwertaufgaben* Arch Rational Mech Anal, Vol 36, 1970, pp 348-355
- 11 J NITSCHKE, *L^∞ -convergence of finite element approximation 2* Conference on Finite Elements, Rennes (France), 1975
- 12 R RANNACHER, *L^∞ -Konvergenz linearer finiter Elemente* Math Z, Vol 149, 1976, pp 69-77
- 13 R SCOTT, *Optimal L^x -estimates for the finite element method on irregular meshes* Math Comp, Vol 30, 1976