

RAIRO

ANALYSE NUMÉRIQUE

KATSUSHI OHMORI

TERUO USHIJIMA

**A technique of upstream type applied to a linear
nonconforming finite element approximation
of convective diffusion equations**

RAIRO – Analyse numérique, tome 18, n° 3 (1984), p. 309-332.

http://www.numdam.org/item?id=M2AN_1984__18_3_309_0

© AFCET, 1984, tous droits réservés.

L'accès aux archives de la revue « RAIRO – Analyse numérique » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

**A TECHNIQUE OF UPSTREAM TYPE APPLIED
TO A LINEAR NONCONFORMING FINITE ELEMENT
APPROXIMATION OF CONVECTIVE
DIFFUSION EQUATIONS (*)**

by Katsushi OHMORI ⁽¹⁾ and Teruo USHIJIMA ⁽²⁾

Summary — We present a technique of upstream type in a linear nonconforming finite element approximation of convective diffusion equations. It is then shown that this scheme satisfies the discrete maximum principle and leads to an $O(h)$ error estimate in H^1 -norm. Some numerical examples are given for the model problem.

Résumé — On présente une technique de décentrage dans l'approximation par un élément fini de degré un non conforme des équations de diffusion-convection. Ensuite, il est montré que ce schéma satisfait au principe du maximum discret et conduit à l'estimation $O(h)$ d'erreur dans $H^1(\Omega)$. Quelques exemples numériques sont donnés pour le problème modèle.

INTRODUCTION

In this note a technique of upstream type is introduced in a linear nonconforming finite element approximation of convective diffusion equations. The nonconforming element under consideration here is so-called a piecewise linear element using *Loof connections*, which were thoroughly investigated by Crouzeix and Raviart [7] and Temam [16] from the theoretical interest

(*) Received in November 1982

⁽¹⁾ Department of Mathematics, Toyama Mercantile Marine College, 1-2, Ebie-Neriya, Shinminato-shi, Toyama, 933-02, Japan

⁽²⁾ Department of Information Mathematics, The University of Electro-Communications, 1-5-1, Chofugaoka, Chofu-shi, Tokyo, 182, Japan

occured in studying the approximations of incompressible flow problems. For practical computations, see also the recent book by Thomasset [17].

On the other hand, several techniques of upstream type to the usual piecewise linear element were developed in recent years in Japan (Baba and Tabata [1], Ikeda [10], Kanayama [11], Kikuchi and Ushijima [13]). Our present method is an extension of one of such techniques to the considered non-conforming element, which is obtained along the way of the modification of the bilinear form corresponding to the convective term, mentioned in Kikuchi and Ushijima [13]. Then we introduce barycentric domains corresponding to mid-points of sides of all triangles belonging to the triangulation T_h in order to define the lumped regions. Recently, Dervieux and Thomasset [8] also proposed the barycentric domain associated with the linear nonconforming finite element in order to derive an upwind scheme to the convective term. However, their scheme is different from our scheme.

An outline of the paper is as follows. In Section 1, notation and the model problem are presented. Section 2 is devoted to the construction of a lumping method based on the considered nonconforming element. In Section 3, our technique of upstream type is proposed. Then we show the discrete maximum principle for our scheme in Section 4, and an $O(h)$ H^1 error estimate in Section 5. In Section 6, we give some numerical examples.

The authors would like to express their sincere thanks to the referee of this paper for his valuable comments and constructive recommendations which are most helpful to improve an earlier version of the paper. Following his idea Lemma 3 is obtained, which clarifies a feature of our technique of upstream type.

1. NOTATION AND PRELIMINARIES

Let Ω be a polygonal bounded connected domain of \mathbb{R}^2 with the boundary Γ . For a non-negative integer m , let $H^m(\Omega)$ be the usual m th order Sobolev space equipped with the norm and the semi-norm

$$\|v\|_{m,\Omega} = \left(\sum_{|\alpha| \leq m} \|D^\alpha v\|_{0,\Omega}^2 \right)^{1/2}, \quad (1.1)$$

$$|v|_{m,\Omega} = \left(\sum_{|\alpha|=m} \|D^\alpha v\|_{0,\Omega}^2 \right)^{1/2}, \quad (1.2)$$

where $\|\cdot\|_{0,\Omega}$ is the norm of $L^2(\Omega)$. The scalar product in $L^2(\Omega)$ is given by (\cdot, \cdot) . We set as usual

$$H_0^1(\Omega) = \{v \in H^1(\Omega); v|_\Gamma = 0\}. \quad (1.3)$$

We consider the following stationary convective diffusion equation

$$(E) \quad \begin{cases} -v \Delta u + (\mathbf{b} \cdot \nabla) u = f & \text{in } \Omega, \\ u = u_0 & \text{on } \Gamma, \end{cases} \quad (1.4)$$

$$(1.5)$$

where v is a positive constant, $\mathbf{b} = \mathbf{b}(x) \in C^1(\overline{\Omega})^2$, $f \in L^2(\Omega)$ and $u_0 \in H^1(\Omega)$.

Let $a(u, v)$ and $b(u, v)$ be two bilinear forms on $H^1(\Omega) \times H^1(\Omega)$ defined by

$$a(u, v) = \int_{\Omega} \nabla u \cdot \nabla v \, dx, \quad (1.6)$$

$$b(u, v) = \int_{\Omega} (\mathbf{b} \cdot \nabla u) v \, dx. \quad (1.7)$$

Furthermore, we set

$$t(u, v) = va(u, v) + b(u, v). \quad (1.8)$$

We consider the variational formulation (II) for (E) :

$$(II) \quad \begin{cases} \text{Find } u \in H^1(\Omega) \text{ such that} \\ t(u, v) = (f, v) \text{ for all } v \in H_0^1(\Omega), \\ u - u_0 \in H_0^1(\Omega). \end{cases} \quad (1.9)$$

$$(1.10)$$

This problem has a unique solution under the condition that there exists a positive constant α_1 such that

$$v\alpha_0 - 1/2 \cdot \text{div } \mathbf{b} \geq \alpha_1 > 0 \quad \text{in } \Omega, \quad (1.11)$$

where $\alpha_0 > 0$ is less than or equal to the minimum eigenvalue of $-\Delta$ with Dirichlet boundary condition.

It is well known that the *maximum principle* holds for the solution of (II) in the following form (cf. Courant and Hilbert [6]) :

Assume that the solution u of (II) is continuous on $\overline{\Omega}$ and twice continuously differentiable in Ω . Then it holds that

$$\max_{x \in \overline{\Omega}} u(x) \leq \max_{x \in \Gamma} u_0(x) \quad \text{when } f \leq 0 \text{ in } \Omega. \quad (1.12)$$

2. NONCONFORMING FINITE ELEMENT AND LUMPING OPERATOR

In this section we shall consider an approximation of upstream type for the convective term using the linear nonconforming finite element.

Let $\{T_h\}$ be a family of triangulation of $\bar{\Omega}$ made of open triangles K , that is

$$\bar{\Omega} = \bigcup_{K \in T_h} \bar{K}, \quad (2.1)$$

where any two triangles are either disjoint or share at most one side or one vertex. Let h_K be the maximum side length of $K \in T_h$ and ρ_K be the diameter of the inscribed circle in K . Moreover, we set $h = \max_{K \in T_h} h_K$.

In what follows, we assume that $\{T_h\}$ is *regular*. That is, when h tends to 0, there exists a constant $\sigma > 0$, independent of h and K , such that

$$\sigma_K = h_K/\rho_K \leq \sigma \quad \text{for all } K \in T_h. \quad (2.2)$$

Let us recall the linear nonconforming finite element studied by Crouzeix and Raviart [7]. Let B_i , $1 \leq i \leq N$, be the mid-points of sides lying in the interior of Ω and B_i , $N + 1 \leq i \leq N + M$, be the mid-points of sides lying on Γ . Let V_h be the linear nonconforming finite element approximate space of $H^1(\Omega)$ defined by

$$V_h = \{v_h \in L^2(\Omega) : v_h \text{ is linear on } K \in T_h \text{ and is continuous at } B_i, \\ 1 \leq i \leq N + M\}. \quad (2.3)$$

Furthermore, we define

$$V_{0h} = \{v_h \in V_h; v_h = 0 \text{ at } B_i, N + 1 \leq i \leq N + M\}. \quad (2.4)$$

Observe that $V_h \not\subset H^1(\Omega)$ and $V_{0h} \not\subset H_0^1(\Omega)$.

Let w_{ih} , $1 \leq i \leq N + M$, be the elements of V_h such that

$$w_{ih}(B_j) = \delta_{ij} \quad \text{for } 1 \leq i, j \leq N + M, \quad (2.5)$$

where δ_{ij} is the Kronecker delta. Then the sets of functions $\{w_{ih}; 1 \leq i \leq N + M\}$, and $\{w_{ih}; 1 \leq i \leq N\}$, form bases of V_h , and V_{0h} , respectively. This element, however, satisfies the following *compatibility conditions*:

(N1) For any $K_1, K_2 \in T_h$, it holds that

$$\int_{\partial\Gamma_{12}} (v_h|_{K_1} - v_h|_{K_2}) d\gamma = 0 \quad \text{for all } v_h \in V_h, \quad (2.6)$$

where $\partial\Gamma_{12} = \partial K_1 \cap \partial K_2$.

(N2) For any $K \in T_h$, it holds that

$$\int_{\partial K \cap \Gamma} v_h|_K d\gamma = 0 \quad \text{for all } v_h \in V_{0h}. \quad (2.7)$$

We provide the space V_h with the following norm and semi-norm :

$$\| v_h \|_{1,h} = \left(\sum_{K \in T_h} \| v_h \|_{1,K}^2 \right)^{1/2}, \tag{2.8}$$

$$\| v_h \|_h = \left(\sum_{K \in T_h} | v_h |_{1,K}^2 \right)^{1/2}. \tag{2.9}$$

The above conditions (N1) and (N2) imply that $\| \cdot \|_h$ is a norm over the space V_{0h} .

Next, we define the *barycentric domain* associated with the linear non-conforming finite element. For any $K \in T_h$ with vertices $P_{i,K}$, $1 \leq i \leq 3$, let $B_{i,K}$ be the mid-point of the side K'_i opposite to $P_{i,K}$, $1 \leq i \leq 3$, and G_K be the barycenter of K . Consider the triangle S_{ij} , $1 \leq i, j \leq 3, i \neq j$, with vertices $G_K, B_{i,K}$ and $P_{j,K}$, where $k \neq i, j$. We say that S_{ij} is a *barycentric fragment* of K . Then, with each $B_{i,K}$, $1 \leq i \leq 3$, we associate a *barycentric subdomain* S_i^K as follows :

$$S_i^K = \bigcup_{j \neq i} S_{ij}. \tag{2.10}$$

If K_1 and K_2 are adjacent elements having B_i as its common mid-point, we say that $\Omega_i = S_i^{K_1} \cup S_i^{K_2}$ is the *barycentric domain* with respect to B_i . If $B_i \in \Gamma$, we set $\Omega_i = S_i^K$. Furthermore, with each B_i , $1 \leq i \leq N + M$, we associate the index set

$$\Lambda_i = \{ j \neq i;$$

$$B_j \text{ is the mid-point of the side of a triangle having } B_i \text{ as another one } \}. \tag{2.11}$$

For any $j \in \Lambda_i$, $1 \leq i \leq N$, we set as follows :



$$\Gamma_{ij}^S = \partial S_i^K \cap \partial S_j^K. \tag{2.12}$$

If B_i is the mid-point of the side lying in the interior of Ω , we have

$$\partial \Omega_i = \bigcup_{j \in \Lambda_i} \Gamma_{ij}^S. \tag{2.13}$$

In our linear nonconforming finite element approximation, this barycentric domain plays the role of the *lumping region* in the usual conforming finite element approximation (see Kikuchi and Ushijima [13]).

Let $\bar{w}_{i,h}$ be the characteristic function of Ω_i and \bar{V}_h be the linear space spanned by the functions $\bar{w}_{j,h}$, $1 \leq j \leq N + M$. Let L_h be the *lumping operator* from



Barycentric domain Ω_i

Fig. 1. — Lumping region.

V_h onto \bar{V}_h defined by

$$V_h \ni v_h = \sum_{j=1}^{N+M} V_j w_{jh} \mapsto L_h v_h = \bar{v}_h = \sum_{j=1}^{N+M} V_j \bar{w}_{jh} \in \bar{V}_h. \quad (2.14)$$

It is easily seen that the lumping operator L_h satisfies the following properties :

$$\| v_h \|_{0,\Omega} = \| L_h v_h \|_{0,\Omega} \quad \text{for all } v_h \in V_h, \quad (2.15)$$

$$\| v_h - L_h v_h \|_{0,\Omega} \leq h \| v_h \|_h \quad \text{for all } v_h \in V_h. \quad (2.16)$$

3. UPSTREAM-LIKE SCHEME OF THE NONCONFORMING TYPE

We define the following approximate bilinear forms on $V_h \times V_h$:

$$a_h(u_h, v_h) = \sum_{K \in T_h} \int_K \nabla u_h \cdot \nabla v_h \, dx, \tag{3.1}$$

$$b_h(u_h, v_h) = \sum_{K \in T_h} \int_K (\mathbf{b} \cdot \nabla u_h) v_h \, dx. \tag{3.2}$$

Then we set for any $u_h, v_h \in V_h$

$$t_h(u_h, v_h) = \nu a_h(u_h, v_h) + b_h(u_h, v_h). \tag{3.3}$$

In [14] we have considered the following approximate problem of Galerkin type :

$$(\Pi_h) \begin{cases} \text{Find } \bar{u}_h \in V_h \text{ such that} \\ t_h(u_h, v_h) = (f, v_h) \text{ for all } v_h \in V_{0h}, \\ u_h - u_{0h} \in V_{0h}, \end{cases} \tag{3.4}$$

$$\tag{3.5}$$

where $u_{0h} \in V_h$ is chosen so that $u_{0h}(B_i) = u_0(B_i)$, $N + 1 \leq i \leq N + M$.

Now, we shall consider the modification of $b_h(u_h, v_h)$ by using the lumping process with the barycentric domain, following the procedure stated in Kikuchi and Ushijima [13] for the case of conforming piecewise linear approximation. In the first time, we rewrite $b_h(u_h, v_h)$ as follows :

$$b_h(u_h, v_h) = b_h^1(u_h, v_h) + b_h^2(u_h, v_h), \tag{3.6}$$

where

$$b_h^1(u_h, v_h) = \sum_{K \in T_h} \int_K (\text{div } u_h \mathbf{b}) v_h \, dx, \tag{3.7}$$

and

$$b_h^2(u_h, v_h) = - \sum_{K \in T_h} \int_K (\text{div } \mathbf{b}) u_h v_h \, dx. \tag{3.8}$$

Then we modify $b_h^1(u_h, v_h)$ by $b_h^1(u_h, L_h v_h)$. According to the patch-wise application of the Gauss divergence formula, it can be easily verified that

$$b_h^1(u_h, L_h v_h) = \sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^s} (\mathbf{b} \cdot \mathbf{n}_j) u_h \, d\gamma v_h(B_j) + \sum_{j=N+1}^{N+M} \int_{\partial\Omega_j \cap \Gamma} (\mathbf{b} \cdot \mathbf{n}_j) u_h \, d\gamma v_h(B_j), \tag{3.9}$$

where \mathbf{n}_j is the unit outer normal vector along $\partial\Omega_j$. Taking account of (3.9) we define the modified form $\tilde{b}_h^1(u_h, v_h)$ as follows :

$$\begin{aligned} \tilde{b}_h^1(u_h, v_h) = & \sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^S} (\mathbf{b} \cdot \mathbf{n}_j) u_h^{jk} d\gamma v_h(B_j) \\ & + \sum_{j=N+1}^{N+M} \int_{\partial\Omega_j \cap \Gamma} (\mathbf{b} \cdot \mathbf{n}_j) d\gamma u_h(B_j) v_h(B_j), \end{aligned} \quad (3.10)$$

where

$$\left. \begin{aligned} u_h^{jk} &= \lambda_{jk} u_h(B_j) + (1 - \lambda_{jk}) u_h(B_k), \\ \lambda_{jk} &= 1 - \lambda_{kj}, \\ |\lambda_{jk}| &\leq \Lambda \quad (\Lambda \text{ is a constant independent of } j, k \text{ and } h). \end{aligned} \right\} \quad (3.11)$$

Furthermore, $b_h^2(u_h, v_h)$ is modified by $b_h^2(L_h u_h, L_h v_h)$ which is denoted by $\tilde{b}_h^2(u_h, v_h)$. Then we have

$$\begin{aligned} \tilde{b}_h^2(u_h, v_h) = & - \sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^S} (\mathbf{b} \cdot \mathbf{n}_j) d\gamma u_h(B_j) v_h(B_j) \\ & - \sum_{j=N+1}^{N+M} \int_{\partial\Omega_j \cap \Gamma} (\mathbf{b} \cdot \mathbf{n}_j) d\gamma u_h(B_j) v_h(B_j). \end{aligned} \quad (3.12)$$

Thus we can define the *modified* form $\tilde{b}_h(u_h, v_h)$ as follows :

$$\begin{aligned} \tilde{b}_h(u_h, v_h) &= \tilde{b}_h^1(u_h, v_h) + \tilde{b}_h^2(u_h, v_h) \\ &= \sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^S} (\mathbf{b} \cdot \mathbf{n}_j) (u_h^{jk} - u_h(B_j)) d\gamma v_h(B_j). \end{aligned} \quad (3.13)$$

Remark 1 : If we take λ_{jk} as follows, then (3.13) yields the upstream scheme for the convective term :

$$\lambda_{jk} = \begin{cases} 1 & \left(\text{if } \int_{\Gamma_{jk}^S} (\mathbf{b} \cdot \mathbf{n}_j) d\gamma \geq 0 \right), \\ 0 & \text{(otherwise)}. \end{cases} \quad (3.14)$$

Finally, we define our modified form of $t_h(u_h, v_h)$:

$$\tilde{t}_h(u_h, v_h) = \nu a_h(u_h, v_h) + \tilde{b}_h(u_h, v_h). \quad (3.15)$$

Hence, our approximate problem is written as follows :

$$(\tilde{\Pi}_h) \begin{cases} \text{Find } \tilde{u}_h \in V_h \text{ such that} \\ \tilde{t}_h(\tilde{u}_h, v_h) = (f, v_h) \text{ for all } v_h \in V_{0h}, \\ \tilde{u}_h - u_{0h} \in V_{0h}. \end{cases} \tag{3.16}$$

$$\tag{3.17}$$

4. DISCRETE MAXIMUM PRINCIPLE

This section is devoted to the study of the discrete maximum principle for the upstream-like scheme $(\tilde{\Pi}_h)$. From now on, we assume that the triangulation T_h is of acute type, that is, it holds that

$$\tau_K \leq 0 \text{ for all } K \in T_h, \tag{4.1}$$

where

$$\tau_K = \max_{i \neq j} \{ \cos(\nabla\mu_i, \nabla\mu_j)_{\mathbb{R}^2} \} \tag{4.2}$$

and $\mu_i, 1 \leq i \leq 3$, are barycentric coordinates of $x \in K$ with respect to the mid-points $B_{i,K}, 1 \leq i \leq 3$, of the sides of K .

Remark 2 : We note that the above definition of the *acuteness* is equivalent to the usual one (cf. Fujii [9]). Hence (4.1) implies that all the angles of the triangles of T_h are less than or equal to $\pi/2$.

Let

$$\tilde{b}_{jk} = \tilde{b}_h(w_{kh}, w_{jh}) \text{ for } 1 \leq j, k \leq N + M, \tag{4.3}$$

then we have

LEMMA 1 : It holds that

$$\tilde{b}_{jj} = - \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^s} (1 - \lambda_{jk}) (\mathbf{b} \cdot \mathbf{n}_j) d\gamma, \tag{4.4}$$

$$\tilde{b}_{jk} = \begin{cases} \int_{\Gamma_{jk}^s} (1 - \lambda_{jk}) (\mathbf{b} \cdot \mathbf{n}_j) d\gamma & (\text{if } k \in \Lambda_j), \\ 0 & (\text{if } k \notin \Lambda_j), \end{cases} \tag{4.5}$$

and

$$\sum_{k=1}^{N+M} \tilde{b}_{jk} = 0 \text{ for } 1 \leq j \leq N + M. \tag{4.6}$$

Proof : By the definition of $\tilde{b}_h(\cdot, \cdot)$ we have

$$\tilde{b}_{jk} = \sum_{i=1}^{N+M} \delta_{ji} \sum_{m \in \Lambda_i} \int_{\Gamma_{im}^s} (\mathbf{b} \cdot \mathbf{n}_i) (1 - \lambda_{im}) (\delta_{km} - \delta_{ki}) d\gamma,$$

where δ_{ij} is the Kronecker delta. Hence, (4.4) and (4.5) hold. On the other hand, it holds that

$$\sum_{k=1}^{N+M} \tilde{b}_{jk} = \tilde{b}_{jj} + \sum_{k \in \Lambda_j} \tilde{b}_{jk}.$$

Then (4.6) follows from (4.4) and (4.5).

If we take λ_{jk} as in (3.14), then we have from Lemma 1,

$$\tilde{b}_{jk} \leq 0 \quad \text{for } 1 \leq j, k \leq N + M, \quad j \neq k. \tag{4.7}$$

We set

$$a_{jk} = a_h(w_{kh}, w_{jh}) \quad \text{for } 1 \leq j \leq N, \quad 1 \leq k \leq N + M. \tag{4.8}$$

By an analogous discussion to Kikuchi [12] we have

$$\left. \begin{aligned} \sum_{k=1}^{N+M} a_{jk} &= 0 \quad \text{for } 1 \leq j \leq N, \\ a_{jk} &\leq 0 \quad \text{for } 1 \leq j \leq N, \quad 1 \leq k \leq N + M, \quad j \neq k. \end{aligned} \right\} \tag{4.9}$$

The proof of (4.9) can be found in [14].

Now let us return to the scheme $(\tilde{\Pi}_h)$. Observe that $(\tilde{\Pi}_h)$ is equivalent to the following linear system :

$$\left. \begin{aligned} TU + T_1 V &= F, \\ V &= G, \end{aligned} \right\} \tag{4.10}$$

where

$$\left. \begin{aligned} T &= (\tilde{t}_{ij}) = (va_{ij} + \tilde{b}_{ij}) \quad \text{for } 1 \leq i, j \leq N, \\ T_1 &= (\tilde{t}_{ij}) \quad \text{for } 1 \leq i \leq N, \quad N + 1 \leq j \leq N + M, \\ U &= (U_j) = (\tilde{u}_h(B_j)) \quad \text{for } 1 \leq j \leq N, \\ V &= (U_j) \quad \text{for } N + 1 \leq j \leq N + M, \\ F &= (F_j) = (f(B_j) \cdot \text{mes}(\text{supp}(w_{jh}))/3) \quad \text{for } 1 \leq j \leq N, \\ G &= (G_j) = (u_0(B_j)) \quad \text{for } N + 1 \leq j \leq N + M. \end{aligned} \right\} \tag{4.12}$$

We define the matrix

$$T_0 = (\tilde{t}_{ij}) \quad \text{for } 1 \leq i \leq N, \quad 1 \leq j \leq N + M. \tag{4.13}$$

Then we have the following Lemma from (4.6), (4.7) and (4.9).

LEMMA 2 : If we take λ_{jk} as in (3.14), then the matrix T_0 is of non-negative type, that is

$$\left. \begin{aligned} \tilde{t}_{ij} &\leq 0 \quad \text{for } i \neq j, \quad 1 \leq i \leq N, \quad 1 \leq j \leq N + M, \\ \sum_{j=1}^{N+M} \tilde{t}_{ij} &\geq 0 \quad \text{for } 1 \leq i \leq N. \end{aligned} \right\} \quad (4.14)$$

THEOREM 1 : Assume that the triangulation T_h is of acute type and that the matrix T is invertible. If we take λ_{jk} as in (3.14), then we have

$$\max_{1 \leq j \leq N} U_j \leq \max \left(0, \max_{1 \leq j \leq M} G_{N+j} \right) \quad \text{if} \quad \max_{1 \leq j \leq N} F_j \leq 0. \quad (4.15)$$

Proof : This fact comes from general considerations due to Ciarlet [3]. For the sake of completeness we, however, give a direct proof as follows.

First we prove (4.15) in the case of

$$\max_{1 \leq j \leq N} F_j < 0. \quad (4.16)$$

Let $U_i = \max_{1 \leq j \leq N} U_j$. When $U_i \leq 0$, then the assertion is trivial. Then we let $U_i > 0$. Assume that $U_i > \max_{1 \leq j \leq M} U_{N+j}$. Since

$$\sum_{j=1}^N \tilde{t}_{ij} U_j + \sum_{j=1}^M \tilde{t}_{i,N+j} U_{N+j} = F_i \quad \text{for } 1 \leq i \leq N,$$

we have

$$\begin{aligned} \tilde{t}_{ii} U_i &= \sum_{\substack{j=1 \\ j \neq i}}^N (-\tilde{t}_{ij}) U_j + \sum_{j=1}^M (-\tilde{t}_{i,N+j}) U_{N+j} + F_i \\ &\leq -U_i \sum_{\substack{j=1 \\ j \neq i}}^{N+M} \tilde{t}_{ij} + F_i, \end{aligned} \quad (4.17)$$

where we use in the last inequality the fact that the matrix T_0 is non-negative type by Lemma 2. Therefore we have

$$0 > F_i \geq U_i \sum_{j=1}^{N+M} \tilde{t}_{ij} \geq 0, \quad (4.18)$$

which is a contradiction. Thus we obtain (4.15).

Next, we prove (4.15) for the case

$$\max_{1 \leq j \leq N} F_j \leq 0. \quad (4.19)$$

We set a column vector F_ϵ for any $\epsilon > 0$ as follows :

$$F_\epsilon = (F_{\epsilon j}) = (F_j - \epsilon) \quad \text{for } 1 \leq j \leq N. \tag{4.20}$$

Then we have $\max_{1 \leq j \leq N} F_{\epsilon j} < 0$. Let U_ϵ be the solution of the following equation :

$$TU + T_1 V = F_\epsilon, \tag{4.21}$$

$$V = G. \tag{4.22}$$

For the solution U_ϵ of (4.21)-(4.22), (4.15) holds by the first half of this proof. Furthermore, since the matrix T is invertible, we have

$$U_\epsilon = T^{-1}(F_\epsilon - T_1 V) \tag{4.23}$$

and

$$U_\epsilon \rightarrow U \quad \text{as } \epsilon \rightarrow 0. \tag{4.24}$$

Thus our assertion is completely proved.

Remark 3 : In Theorem 1, if the triangulation T_h is of *strictly acute type*, namely if all the angles of triangles of T_h are less than $\pi/2$, then the matrix T is invertible. Another condition to assert the invertibility of T is that $\text{div } \mathbf{b}$ is non-positive in Ω , which will be shown in Theorem 3.

5. ERROR ESTIMATE FOR UPSTREAM-LIKE APPROXIMATION

In the first time, we show that the modified form $\tilde{b}_h(\cdot, \cdot)$ is *admissible* in the sence defined in Kikuchi and Ushijima [13].

THEOREM 2 : *Assume that there exists a constant $D > 0$ independent of j, k and h such that*

$$h \cdot \text{mes} (\Gamma_{jk}^S) \leq D \cdot \text{mes} (S_{jk}), \tag{5.1}$$

then there exists a constant $C > 0$ independent of h ($0 < h \leq \bar{h}$) such that

$$| b_h(u_h, v_h) - \tilde{b}_h(u_h, v_h) | \leq Ch \| u_h \|_{1,h} \| v_h \|_{1,h} \tag{5.2}$$

for all $u_h, v_h \in V_h$.

Proof : We follow the proof of Proposition 2 of [13] with suitable modifications. We may write for all $u_h, v_h \in V_h$

$$\begin{aligned} b_h(u_h, v_h) - \tilde{b}_h(u_h, v_h) &= b_h^1(u_h, v_h) - b_h^1(u_h, L_h v_h) + b_h^1(u_h, L_h v_h) \\ &\quad - \tilde{b}_h^1(u_h, v_h) + b_h^2(u_h, v_h) - \tilde{b}_h^2(u_h, v_h). \end{aligned}$$

Using (2.13) and (2.14), one can easily check that

$$|b_h^1(u_h, v_h) - b_h^1(u_h, L_h v_h)| \leq C_1 h \|u_h\|_{1,h} \|v_h\|_{1,h} \tag{5.3}$$

and

$$|b_h^2(u_h, v_h) - \tilde{b}_h^2(u_h, v_h)| \leq C_2 h \|u_h\|_{1,h} \|v_h\|_{1,h} \tag{5.4}$$

with the appropriate constants $C_i, 1 \leq i \leq 2$.

Thus it suffices to estimate $b_h^1(u_h, L_h v_h) - \tilde{b}_h^1(u_h, v_h)$. From (3.7) and (3.8) we find that

$$\begin{aligned} b_h^1(u_h, L_h v_h) - \tilde{b}_h^1(u_h, v_h) &= \sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^S} (\mathbf{b} \cdot \mathbf{n}_j) (u_h - u_h^{jk}) d\gamma v_h(B_j) \\ &+ \sum_{j=N+1}^{N+M} \int_{\Omega_j \cap \Gamma} (\mathbf{b} \cdot \mathbf{n}_j) (u_h - u_h(B_j)) d\gamma v_h(B_j) = I_1 + I_2. \end{aligned} \tag{5.5}$$

Taking into account that $\Gamma_{jk}^S = \Gamma_{kj}^S$ and $\mathbf{n}_j = -\mathbf{n}_k$, we have

$$\begin{aligned} I_1 &= 1/2 \sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^S} (\mathbf{b} \cdot \mathbf{n}_j) \{ \lambda_{jk} (u_h - u_h(B_j)) \\ &+ (1 - \lambda_{jk}) (u_h - u_h(B_k)) \} d\gamma (v_h(B_j) - v_h(B_k)). \end{aligned} \tag{5.6}$$

On the other hand, since u_h is linear on \bar{S}_{jk} it is easy to check that for any $x \in \Gamma_{jk}^S$

$$|u_h(x) - u_h(B_j)| \leq h \text{mes}(S_{jk})^{-1/2} \|\nabla v_h\|_{L^2(S_{jk})} \tag{5.7}$$

and

$$|v_h(B_j) - v_h(B_k)| \leq h \text{mes}(S_{jk})^{-1/2} \|\nabla v_h\|_{L^2(S_{jk})}. \tag{5.8}$$

Then from the properties of λ_{jk} , (5.7), (5.8) and the assumption (5.1), we find that

$$|I_1| \leq C_3 h \|u_h\|_{1,h} \|v_h\|_{1,h}. \tag{5.9}$$

Next, let us estimate for I_2 . Here we assume that the mid-points of sides lying on $\Gamma, B_i, N + 1 \leq i \leq N + M$, are located consecutively on the boundary in anti-clockwise orientation such that

$$B_j = \text{mid-point of } \overline{P_j P_{j+1}} \text{ for } N + 1 \leq j \leq N + M, \tag{5.10}$$

where $P_j, N + 1 \leq j \leq N + M$, are boundary vertices with $P_{N+M+1} = P_{N+1}$.

Then we may write

$$\begin{aligned}
 I_2 = & \sum_{j=N+1}^{N+M} \int_{P_j}^{P_{j+1}} (\mathbf{b}(B_j) \cdot \mathbf{n}_j) (u_h - u_h(B_j)) \, d\gamma \, v_h(B_j) \\
 & + \sum_{j=N+1}^{N+M} \int_{P_j}^{P_{j+1}} ((\mathbf{b} - \mathbf{b}(B_j)) \cdot \mathbf{n}_j) (u_h - u_h(B_j)) \, d\gamma \, v_h(B_j). \quad (5.11)
 \end{aligned}$$

Since u_h is linear on $\overline{P_j P_{j+1}}$, $N + 1 \leq j \leq N + M$, the first term of I_2 vanishes. Since the second term of I_2 is equal to

$$\begin{aligned}
 \sum_{j=N+1}^{N+M} \left\{ \int_{P_j}^{B_j} ((\mathbf{b} - \mathbf{b}(B_j)) \cdot \mathbf{n}_j) (u_h - u_h(B_j)) \, d\gamma \right. \\
 \left. + \int_{B_j}^{P_{j+1}} ((\mathbf{b} - \mathbf{b}(B_j)) \cdot \mathbf{n}_j) (u_h - u_h(B_j)) \, d\gamma \right\} v_h(B_j),
 \end{aligned}$$

using (5.1), (5.7) and the following fact

$$|(\mathbf{b} - \mathbf{b}(B_j)) \cdot \mathbf{n}_j| \leq C_4 h \quad \text{for } x \in \overline{P_j B_j} \quad (\text{resp. } \overline{B_j P_{j+1}}),$$

we conclude that there exists a constant C_5 such that

$$|I_2| \leq C_5 h \|u_h\|_{1,h} \|v_h\|_{1,h}. \quad (5.12)$$

Combining (5.3), (5.4), (5.9) and (5.12), we obtain (5.2).

Remark 4 : Since from Proposition 4.13 of Temam [16] the following discrete Poincaré inequality holds

$$\|v_h\|_{0,\Omega} \leq C(\Omega) \|v_h\|_h \quad \text{for all } v_h \in V_{0h} \quad (5.13)$$

with a constant $C(\Omega) > 0$ independent of $h \in (0, \bar{h}]$, $\|\cdot\|_{1,h}$ and $\|\cdot\|_h$ are equivalent norms on V_{0h} .

Hence we find a constant $C > 0$ independent of $h \in (0, \bar{h}]$ such that

$$|b_h(u_h, v_h) - \tilde{b}_h(u_h, v_h)| \leq Ch \|u_h\|_h \|v_h\|_h \quad \text{for all } u_h, v_h \in V_{0h}. \quad (5.14)$$

In what follows, we shall restrict our attention to the homogeneous Dirichlet problem for simplicity, which is denoted by (E^0) . Thus we consider the following problem :

$$(\tilde{\Pi}_h^0) \quad \left\{ \begin{array}{l} \text{Find } \tilde{u}_h \in V_{0h} \text{ such that} \\ \tilde{t}_h(\tilde{u}_h, v_h) = (f, v_h) \quad \text{for all } v_h \in V_{0h}. \end{array} \right.$$

LEMMA 3 : It holds that for any $v_h \in V_{0h}$

$$\begin{aligned} \tilde{b}_h^1(v_h, v_h) + \frac{1}{2} \tilde{b}_h^2(v_h, v_h) &= \sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^S} \frac{1}{2} (\mathbf{b} \cdot \mathbf{n}_j) (v_h(B_j) - v_h(B_k))^2 \left(\lambda_{jk} - \frac{1}{2} \right) d\gamma. \end{aligned} \quad (5.15)$$

Proof : Let v_h be an arbitrary element in V_{0h} . From (3.10) and (3.12) we have

$$\tilde{b}_h^1(v_h, v_h) + \frac{1}{2} \tilde{b}_h^2(v_h, v_h) = \sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^S} (\mathbf{b} \cdot \mathbf{n}_j) \left(v_h^{jk} - \frac{1}{2} v_h(B_j) \right) v_h(B_j) d\gamma$$

Since $\Gamma_{jk}^S = \Gamma_{kj}^S$ and $\mathbf{n}_j = -\mathbf{n}_k$, we obtain by using (3.11)

$$\begin{aligned} \tilde{b}_h^1(v_h, v_h) + \frac{1}{2} \tilde{b}_h^2(v_h, v_h) &= \sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^S} \frac{1}{2} (\mathbf{b} \cdot \mathbf{n}_j) \left\{ \left(v_h^{jk} - \frac{1}{2} v_h(B_j) \right) v_h(B_j) - \left(v_h^{kj} - \frac{1}{2} v_h(B_k) \right) v_h(B_k) \right\} d\gamma \\ &= \sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^S} \frac{1}{2} (\mathbf{b} \cdot \mathbf{n}_j) \left(\lambda_{jk} - \frac{1}{2} \right) \{ v_h(B_j)^2 - 2 v_h(B_j) v_h(B_k) + v_h(B_k)^2 \} d\gamma \\ &= \sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^S} \frac{1}{2} (\mathbf{b} \cdot \mathbf{n}_j) (v_h(B_j) - v_h(B_k))^2 \left(\lambda_{jk} - \frac{1}{2} \right) d\gamma. \end{aligned}$$

THEOREM 3 : Assume that there exists a constant $\alpha' > 0$ such that

$$vC(\Omega)^{-2} - \frac{1}{2} \cdot \operatorname{div} \mathbf{b} \geq \alpha' > 0 \quad \text{in } \Omega, \quad (5.17)$$

where $C(\Omega)$ is the constant in the discrete Poincaré inequality (5.13). If we take λ_{jk} as in (3.14), then the problem $(\tilde{\Pi}_h^0)$ has a unique solution $\tilde{u}_h \in V_{0h}$.

Proof : It is sufficient to show the V_{0h} -coercivity of $\tilde{t}_h(\cdot, \cdot)$. From the definition of $\tilde{t}_h(\cdot, \cdot)$ and Lemma 3, we have for all $v_h \in V_{0h}$

$$\begin{aligned} \tilde{t}_h(v_h, v_h) &= \nu a_h(v_h, v_h) + \tilde{b}_h^1(v_h, v_h) + \frac{1}{2} \tilde{b}_h^2(v_h, v_h) + \frac{1}{2} \tilde{b}_h^2(v_h, v_h) \\ &= \nu \|v_h\|_h^2 + \sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^S} \frac{1}{2} (\mathbf{b} \cdot \mathbf{n}_j) (v_h(B_j) - v_h(B_k))^2 \left(\lambda_{jk} - \frac{1}{2} \right) d\gamma \\ &\quad - \frac{1}{2} \sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^S} (\mathbf{b} \cdot \mathbf{n}_j) d\gamma v_h(B_j)^2. \end{aligned} \quad (5.18)$$

By virtue of the choice of λ_{jk} , we find that

$$\sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^S} \frac{1}{2} (\mathbf{b} \cdot \mathbf{n}_j) (v_h(B_j) - v_h(B_k))^2 \left(\lambda_{jk} - \frac{1}{2} \right) d\gamma \geq 0. \quad (5.19)$$

Then we have

$$\begin{aligned} \tilde{t}_h(v_h, v_h) &\geq \nu \|v_h\|_h^2 - \frac{1}{2} \sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^S} (\mathbf{b} \cdot \mathbf{n}_j) d\gamma v_h(B_j)^2 \\ &= \nu \|v_h\|_h^2 - \frac{1}{2} \left\{ \sum_{j=1}^{N+M} \sum_{k \in \Lambda_j} \int_{\Gamma_{jk}^S} (\mathbf{b} \cdot \mathbf{n}_j) d\gamma v_h(B_j)^2 \right. \\ &\quad \left. + \sum_{j=N+1}^{N+M} \int_{\partial\Omega_j \cap \Gamma} (\mathbf{b} \cdot \mathbf{n}_j) d\gamma v_h(B_j)^2 \right\}, \end{aligned}$$

where in the last equality we used the fact that $v_h(B_j)=0$ for $N+1 \leq j \leq N+M$.

According to the patch-wise application of the Gauss divergence formula, it holds that

$$\tilde{t}_h(v_h, v_h) \geq \nu \|v_h\|_h^2 - \frac{1}{2} \sum_{j=1}^{N+M} \int_{\Omega_j} (\operatorname{div} \mathbf{b}) |L_h v_h|^2 dx. \quad (5.20)$$

Then for any constant ε satisfying $0 < \varepsilon < \min \{ 1, \alpha' C(\Omega)^2/\nu \}$ we have

$$\tilde{t}_h(v_h, v_h) \geq \nu\varepsilon \|v_h\|_h^2 \quad \text{for all } v_h \in V_{0h}. \quad (5.21)$$

THEOREM 4 : *Assume the hypotheses of Theorem 3. Then $t_h(\cdot, \cdot)$ is coercive on V_{0h} for any sufficiently small h .*

Proof . For all $v_h \in V_{0h}$ we have

$$t_h(v_h, v_h) = \tilde{t}_h(v_h, v_h) + b_h(v_h, v_h) - \tilde{d}_h(v_h, v_h). \quad (5.22)$$

Therefore, from Theorem 3 and Remark 4 it holds that

$$t_h(v_h, v_h) \geq (\nu\varepsilon - Ch) \|v_h\|_h^2 \quad \text{for all } v_h \in V_{0h}, \quad (5.23)$$

where C is the positive constant independent of h in (5.14).

Thus, we find that there is a constant $h_0 > 0$ such that it holds for some constant $\alpha^* > 0$

$$t_h(v_h, v_h) \geq \alpha^* \|v_h\|_h^2 \quad \text{for all } v_h \in V_{0h}, \quad (5.24)$$

provided that $h \leq h_0$.

Now, let us derive a bound for the error $\| \tilde{u}_h - u_h \|_h$, where \tilde{u}_h is the solution of $(\tilde{\Pi}_h^0)$ with λ_{jk} as in (3.14) and u_h is the solution of (Π_h^0) which is the Galerkin approximation of (E^0) .

THEOREM 5 : *Assume the hypotheses of Theorem 3. Then we have*

$$\| \tilde{u}_h - u_h \|_h \leq Ch \| f \|_{0,\Omega} \tag{5.25}$$

for some constant $C > 0$.

Proof : Since

$$\tilde{t}_h(\tilde{u}_h, v_h) = (f, v_h) = t_h(u_h, v_h) \quad \text{for all } v_h \in V_{0h}, \tag{5.26}$$

we may write

$$t_h(u_h, v_h) - \tilde{t}_h(u_h, v_h) + \tilde{t}_h(u_h, v_h) = \tilde{t}_h(\tilde{u}_h, v_h) \quad \text{for all } v_h \in V_{0h}. \tag{5.27}$$

Hence we have

$$\begin{aligned} \tilde{t}_h(\tilde{u}_h - u_h, v_h) &= t_h(u_h, v_h) - \tilde{t}_h(u_h, v_h) \\ &= b_h(u_h, v_h) - \tilde{b}_h(u_h, v_h). \end{aligned} \tag{5.28}$$

According to Remark 4, it holds that

$$| \tilde{t}_h(\tilde{u}_h - u_h, v_h) | \leq Ch \| u_h \|_h \| v_h \|_h. \tag{5.29}$$

Taking $v_h = \tilde{u}_h - u_h$ in (5.29) we obtain from the coercivity of $\tilde{t}_h(\cdot, \cdot)$

$$\forall \epsilon \| \tilde{u}_h - u_h \|_h^2 \leq Ch \| u_h \|_h \| \tilde{u}_h - u_h \|_h.$$

Hence we have

$$\| \tilde{u}_h - u_h \|_h \leq C' h \| u_h \|_h. \tag{5.30}$$

On the other hand, from the coercivity of $t_h(\cdot, \cdot)$ we can show that

$$\| u_h \|_h \leq 1/\alpha^* \cdot \| f \|_{0,\Omega}. \tag{5.31}$$

Then, from (5.30) and (5.31) the assertion follows.

Remark 5 : In order to obtain the error estimate for $u - u_h$ in the norm $\| \cdot \|_h$, we can apply the *primal hybrid* finite element method introduced by Raviart-Thomas [15], since the linear nonconforming finite element is one of the *hybrid* elements. Then we have the following result :

THEOREM 6 : *In addition to the hypotheses of Theorem 3, suppose that $\{T_h\}$ is regular. If $u \in H^2(\Omega) \cap H_0^1(\Omega)$, then we have*

$$\|u - u_h\|_h \leq Ch \|u\|_{2,\Omega}. \quad (5.32)$$

Using Theorems 5 and 6, we can derive an error estimate for the *upstream-like* approximation with the linear nonconforming finite element.

THEOREM 7 : *Assume the hypotheses of Theorem 6. Then we have*

$$\|u - \tilde{u}_h\|_h \leq Ch(\|u\|_{2,\Omega} + \|f\|_{0,\Omega}). \quad (5.33)$$

6. NUMERICAL EXAMPLES

As an illustration, here we adopt one of the problems treated in Kikuchi and Ushijima [13]. Namely, our model problem is :

$$\left. \begin{aligned} -\nu \Delta u + (\mathbf{b} \cdot \nabla) u &= 1 \quad \text{in } \Omega = (0, 1) \times (0, 1), \\ u &= 0 \quad \text{on } \Gamma, \end{aligned} \right\} \quad (6.1)$$

where $\mathbf{b} = (1, 0)$. In [13], the following initial boundary value problem (6.2) is taken as an approximation of (6.1) for sufficiently small ν in the region far from $x_1 = 1$.

$$\left. \begin{aligned} \frac{\partial u}{\partial x_1} - \nu \frac{\partial^2 u}{\partial x_2^2} &= 1 \quad \text{in } \Omega, \\ u(0, x_2) &= 0 \quad \text{for } 0 < x_2 < 1, \\ u(x_1, 0) = u(x_1, 1) &= 0 \quad \text{for } 0 < x_1 < 1. \end{aligned} \right\} \quad (6.2)$$

Examples of employed meshes are pictured in figure 2, where N denotes a number of elements along the side $x_2 = 0$ (or $x_1 = 0$) of the domain Ω . Figures 3, 4, 5 and 6 show the distributions of the numerical solutions u_h and \tilde{u}_h along the line $x_1 = 1/2$ of the square domain Ω , where u_h is the linear nonconforming finite element approximation and \tilde{u}_h is the linear nonconforming finite element approximation of upstream type. In these figures, continuous curves are the profiles of numerical solutions of the problem (6.2), which are denoted by PEA. Among these results, the Galerkin method gives a strongly oscillating solution for the coarse meshes and the small values of ν , but gives an improved one for sufficiently fine meshes. On the other hand, our method gives a non-oscillating and reasonable solution.

The authors are grateful to Professor Sotoshi Shinbo of Toyama Prefectural College of Technology, who gave us the opportunity to use HITAC M 150 of his college. Besides this computer, we used for the computations in this paper HITAC 8350/M 180 of The University of Electro-Communications and OKITAC 4500 C of Toyama Mercantile Marine College.




Fig. 2. — Finite element meshes for $N = 5$.




Fig. 3. — Distributions of $u_h(0.5, x_2)$ and $\tilde{u}_h(0.5, x_2)$ with $\nu = 0.01$ and $N = 5$.




Fig. 4. — Distributions of $u_h(0.5, x_2)$ and $\tilde{u}_h(0.5, x_2)$ with $v = 0.01$ and $N = 10$.




Fig. 5. — Distributions of $u_h(0.5, x_2)$ and $\tilde{u}_h(0.5, x_2)$ with $v = 0.001$ and $N = 10$.




Fig. 6. — Distributions of $u_h(0.5, x_2)$ and $\tilde{u}_h(0.5, x_2)$ with $\nu = 0.001$ and $N = 20$.

REFERENCES

1. K. BABA and M. TABATA, *On a conservative upwind finite element scheme for convective diffusion equations*, R.A.I.R.O., Anal. Numér., Vol. 15, 1981, pp. 3-25.
2. F. BREZZI, *On the existence, uniqueness and approximation of saddle-point problems arising from Lagrangian multipliers*, R.A.I.R.O., Anal. Numér., Vol. 8, 1974, pp. 129-151.
3. P. G. CIARLET, *Discrete maximum principle for finite-difference operators*, Aeq. Math., Vol. 4, 1970, pp. 338-352.
4. P. G. CIARLET, *The Finite Element Method for Elliptic Problems*, North Holland, Amsterdam, 1978.
5. P. G. CIARLET and P.-A. RAVIART, *Maximum principle and uniform convergence for the finite element method*, Comput. Methods Appl. Mech. Engrg., Vol. 2, 1973, pp. 17-31.
6. R. COURANT and D. HILBERT, *Methods of Mathematical Physics*, Vol. II, Interscience Publishers, New York, 1962.
7. M. CROUZEIX and P.-A. RAVIART, *Conforming and nonconforming finite element methods for solving the stationary Stokes equations I*, R.A.I.R.O., Anal. Numér., Vol. 7, 1973, pp. 33-76.
8. A. DERVIEUX and F. THOMASSET, *Sur l'approximation d'écoulements multilfluides incompressibles visqueux par des éléments finis triangulaires de degré un*, Rapports de Recherche 67 (LABOLIA INRIA), Avril, 1981.
9. H. FUJII, *Some remarks on finite element analysis of time-dependent field problem*, Theory and Practice in Finite Element Structural Analysis, Y. Yamada and R. H. Gallagher, Eds., University of Tokyo Press, 1973, pp. 91-106.
10. T. IKEDA, *Artificial viscosity in finite element approximations to the diffusion equations with drift terms*, H. Fujita and M. Yamaguti, Eds., Lecture Notes in Num. Appl. Anal., Kinokuniya, Tokyo, Vol. 2, 1980, pp. 59-78.
11. H. KANAYAMA, *Discrete models for salinity distribution in a bay — Conservative law and maximum principle*, Proc. Japan Nat. Congr. for Applied Mech., 1980, Theoretical and Applied Mechanics, Vol. 28, 1980, pp. 559-579.
12. F. KIKUCHI, *Discrete maximum principle and artificial viscosity in finite element approximation to convective diffusion equations*, ISAS Report., No. 550, 1977.
13. F. KIKUCHI and T. USHIJIMA, *Theoretical analysis of some finite element schemes for convective diffusion equations*, R. H. Gallagher, D. H. Norrie, J. T. Oden and O. C. Zienkiewicz, Eds., Finite Elements in Fluids, John Wiley & Sons Ltd, Vol. 4, 1982, pp. 67-87.
14. K. OHMORI, *The discrete maximum principle for nonconforming finite element approximations to stationary convective diffusion equations*, Math. Rep. Toyama Univ., Vol. 2, 1979, pp. 33-52. Corrections, *ibid.*, Vol. 4, 1981, pp. 179-182.
15. P.-A. RAVIART and J. M. THOMAS, *Primal hybrid finite element methods for 2nd order elliptic equations*, Math. Comp., Vol. 31, 1977, pp. 391-413.
16. R. TEMAM, *Navier-Stokes Equations*, North Holland, Amsterdam, 1977.
17. F. THOMASSET, *Implementation of Finite Element Methods for Navier-Stokes Equations*, Springer-Verlag, New York, 1982.
18. T. USHIJIMA, *On a certain finite element method of the upstream type applied to convective diffusion problems*, China-France Symposium on the Finite Element Method, 1982.