

RAIRO

MODÉLISATION MATHÉMATIQUE ET ANALYSE NUMÉRIQUE

F.-J. DELVOS

Approximation properties of periodic interpolation by translates of one function

RAIRO – Modélisation mathématique et analyse numérique,
tome 28, n° 2 (1994), p. 177-188.

http://www.numdam.org/item?id=M2AN_1994__28_2_177_0

© AFCET, 1994, tous droits réservés.

L'accès aux archives de la revue « RAIRO – Modélisation mathématique et analyse numérique » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>



**APPROXIMATION PROPERTIES OF PERIODIC INTERPOLATION
BY TRANSLATES OF ONE FUNCTION (*)**

by F.-J. DELVOS ⁽¹⁾

Communicated by P. J. LAURENT

To the memory of Professor Dr. Lothar COLLATZ

Abstract. — In the paper [5] Golomb has derived a Hilbert space approach to periodic splines of odd degree on uniform meshes which were studied systematically for the first time by Quade and Collatz [9]. Golomb's approach has been extended to more general methods of periodic interpolation by translates of a given periodic function g [1, 2, 3, 8]. It is the objective of this paper to investigate approximation properties of these interpolation methods in spaces of periodic functions which are closely related to g and extend the results of [4]. As an application approximation properties of periodic splines of even degree are obtained.

Résumé. — Dans l'article [3] Golomb a présenté une construction hilbertienne des fonctions-spline périodiques de degré impair dans un réseau uniforme, fonctions étudiées systématiquement pour la première fois par Quade et Collatz [7]. La construction de Golomb a été généralisée pour les méthodes d'interpolation par les translates d'une fonction périodique g . Le but de cet article est d'étudier des propriétés d'approximation des méthodes d'interpolation par les translates de g dans des espaces des fonctions périodiques qui sont associées à la fonction g . En application, nous déduisons les propriétés d'approximation des fonctions-spline périodiques de degré pair.

1. THE INTERPOLATION METHOD

Let g be a fixed real valued periodic function from the Wiener algebra $\mathcal{A}_{2\pi}$ of those continuous periodic functions from $\mathcal{C}_{2\pi}$ which possess an absolutely convergent Fourier series. The inner product of $f, g \in \mathcal{C}_{2\pi}$ is defined by

$$(f, g) = \frac{1}{2\pi} \int_0^{2\pi} f(t) g(t)^* dt .$$

The exponential functions are denoted by $e_k(t) = \exp(ikt)$, $k \in \mathbb{Z}$. If

(*) Manuscript received May, 25, 1993.

(¹) Universität Siegen, Lehrstuhl für Mathematik I, Hölderlin-Strasse 3, D-5900 Siegen (Germany).

$g \in \mathcal{A}_{2\pi}$, then

$$g(t) = \sum_{k=-\infty}^{\infty} (g, e_k) e_k(t), \quad \sum_{k=-\infty}^{\infty} |(g, e_k)| < \infty.$$

Let $t_j = 2\pi j/n$, $j \in \mathbb{Z}$, be a uniform mesh depending on $n \in \mathbb{N}$. The discrete inner product is given by

$$\langle f, g \rangle = \frac{1}{n} \sum_{k=0}^{n-1} f(t_k) g(t_k)^*.$$

The discrete Fourier transform and the finite Fourier transform are related by *aliasing* :

$$\langle f, e_j \rangle = \sum_s \langle f, e_{j+sn} \rangle \quad (1)$$

where $j \in \mathbb{Z}$ and $f \in \mathcal{A}_{2\pi}$.

The *basis functions* b_j , $0 < j < n$, are obtained via the discrete Fourier transform from the generating function g :

$$b_j(t) = \langle g(t - \cdot), e_{-j} \rangle = \sum_s \langle g, e_{j+sn} \rangle e_{j+sn}(t). \quad (2)$$

We have the fundamental relations

$$b_j(t_k) = e_j(t_k) b_j(0) \quad (j, k \in \mathbb{Z}). \quad (3)$$

We assume that

$$b_j(0) \neq 0 \quad (0 < j < n) \quad (4)$$

and define

$$Q_n(f)(t) = \langle f, e_0 \rangle + \sum_{k=1}^{n-1} \langle f, e_k \rangle b_k(t)/b_k(0) \quad (5)$$

for any $f \in \mathcal{C}_{2\pi}$.

Discrete Fourier transform yields the interpolation properties

$$Q_n(f)(t_j) = f(t_j) \quad (0 \leq j < n). \quad (6)$$

The space of interpolants is given by (see [2, 3])

$$V_n(g) = \langle 1, g(\cdot - t_1) - g, \dots, g(\cdot - t_{n-1}) - g \rangle. \quad (7)$$

We assume that the finite Fourier transform of the generating function

g possesses certain properties such that the *existence conditions* $b_j(0) \neq 0$ are valid and certain approximation properties can be established.

Let $d_k, k \in \mathbb{Z}$, be a sequence of real numbers satisfying the following relations

$$d_{-k} = d_k, \quad d_k > d_{k+1} > 0 \quad (k \in \mathbb{N}).$$

Moreover we assume that

$$d_{rm} \leq \alpha_r d_m (r, m \in \mathbb{N}), \quad \alpha := \sum_{r=1}^{\infty} \alpha_r < \infty.$$

We consider mainly two cases of generating functions g :

(I)
$$g(t) = \sum_k d_k e_k(t).$$

(II)
$$g(t) = \sum_k -i \cdot \text{sgn}(k) d_k e_k(-\pi/n) e_k(t).$$

The generating function g defining the interpolation process is independent of the number n of interpolation points in case I while it is dependent in case II.

Trigonometric interpolation is characterized by the choice

(III)
$$g(t) = \sum_{k=-m}^m e_k(t) (m = [n/2]).$$

In this case g depends also on n . We refer to [3] for the proof of the validity of the existence condition.

Two examples are given for possible choices of the sequence (d_k) .

Example a :

$$d_k = k^{-q} \quad (k \in \mathbb{N}).$$

Here q is a real number greater than 1. If q is *even*, case I yields periodic odd degree spline interpolation on uniform mesh as studied by Golomb [3]. The generating function is the well known Bernoulli function $P_q(t)$ up to a factor

$$g(t) = \sum_{k=1}^{\infty} k^{-2r} \cos(kt) = \frac{(-1)^r}{2} P_{2r}(t).$$

The function g is a spline of degree $2r$ with deficiency 1, while $g(\cdot - t_k) - g$ is a spline of degree $2r - 1$.

If q is *odd*, then case II yields periodic even degree midpoint spline interpolation on a uniform mesh [3]. The generating function is the *shifted*

Bernoulli function $P_q(t - \pi/n)$ up to a factor

$$g(t) = \sum_{k=1}^{\infty} k^{-2r-1} \sin(k(t - \pi/n)) = \frac{(-1)^r}{2} P_{2r+1}(t - \pi/n).$$

Example b :

$$d_k = e^{-bk} \quad (k \in \mathbb{N}).$$

Here b is a positive real number. Both cases I and II yield rational trigonometric interpolation processes (see [2, 3]).

In case I the generating function is related to the Poisson kernel

$$P_b(t) = \frac{\sinh(b)}{2(\cosh(b) - \cos(t))},$$

$$g(t) = \sum_{k=1}^{\infty} e^{-kb} \cos(kt) = P_b(t) - \frac{1}{2}.$$

In case II the generating function is related to the *shifted conjugate* Poisson kernel

$$Q_b(t) = \frac{\sin(t)}{2(\cosh(b) - \cos(t))},$$

$$g(t) = \sum_{k=1}^{\infty} e^{-kb} \sin(k(t - \pi/n)) = Q_b(t - \pi/n).$$

2. UNIFORM BOUNDEDNESS OF THE INTERPOLATION PROJECTORS

The Wiener algebra $\mathcal{A}_{2\pi}$ is a dense subalgebra of $\mathcal{C}_{2\pi}$. $\mathcal{A}_{2\pi}$ is a Banach space of continuous periodic functions with respect to the norm (see [6])

$$\|f\|_a = \sum_{k=-\infty}^{\infty} |(g, e_k)|.$$

Moreover, we have

$$\|f\|_{\infty} \leq \|f\|_a \quad (f \in \mathcal{A}_{2\pi})$$

where

$$\|f\|_{\infty} = \sup \{|f(t)| : t \in \mathbb{R}\}.$$

We will first investigate approximation properties of the Fourier partial sum projector which is in some sense a universal approximation operator. The

Fourier partial sum projector F_n is defined by

$$F_n(f) = \sum_{k=-m}^m (f, e_k) e_k \quad (m = [n/2]).$$

Clearly, F_n is a bounded linear operator on $\mathcal{C}_{2\pi}$. On the other hand, it is well known that the norms $\|F_n\|$, $n \in \mathbb{N}$, are not uniformly bounded. According to the Banach Steinhaus principle [7], $F_n(f)$, $n \in \mathbb{N}$, is not convergent for every continuous periodic function. Therefore it is natural to consider F_n as a bounded linear operator from $\mathcal{A}_{2\pi}$ into $\mathcal{C}_{2\pi}$. It follows from the definition of the norms that

$$\|F_n(f)\|_\infty \leq \|f\|_a \quad (f \in \mathcal{A}_{2\pi}).$$

Thus, the norms $\|F_n\|$, $n \in \mathbb{N}$, are uniformly bounded. Since

$$F_n(e_k) = e_k(|k| < n/2),$$

an application of the Banach Steinhaus principle yields uniform convergence

$$\lim_{n \rightarrow \infty} \|f - F_n(f)\|_\infty = 0 \quad (f \in \mathcal{A}_{2\pi}).$$

Replacing $\mathcal{C}_{2\pi}$ by $\mathcal{A}_{2\pi}$ is not a severe restriction from the practical point of view since $\mathcal{A}_{2\pi}$ contains all Lipschitz-continuous periodic functions [6] and in particular those functions from $\mathcal{C}_{2\pi}$ which are piecewise \mathcal{C}^1 [10].

Next we will establish an analogous result for the interpolation projectors Q_n , $n \in \mathbb{N}$. The interpolation projector Q_n is a bounded linear projector from $\mathcal{A}_{2\pi}$ into $\mathcal{C}_{2\pi}$ with $Q_n(f) \in \mathcal{A}_{2\pi}$ for $f \in \mathcal{A}_{2\pi}$.

THEOREM 1: *There is a positive constant c independent of n such that*

$$\|Q_n(f)\|_\infty \leq c \|f\|_a \quad (f \in \mathcal{A}_{2\pi}),$$

i.e., the interpolation projectors Q_n are uniformly bounded as projectors from $\mathcal{A}_{2\pi}$ into $\mathcal{C}_{2\pi}$.

Proof: We start with case I:

$$g(t) = \sum_k d_k e_k(t).$$

Recall that

$$b_j(t) = \sum_s (g, e_{j+sn}) e_{j+sn}(t) = \sum_s d_{j+sn} e_{j+sn}(t)$$

where $0 < j < n$. Since $d_{j+sn} > 0$ we obtain

$$|b_j(t)| \leq b_j(0), \quad 0 < j < n.$$

Next remember that

$$Q_n(f)(t) = \langle f, e_0 \rangle + \sum_{k=1}^{n-1} \langle f, e_k \rangle b_k(t)/b_k(0).$$

This implies

$$|Q_n(f)(t)| \leq \sum_{k=0}^{n-1} |\langle f, e_k \rangle|.$$

Since $\langle f, e_j \rangle = \sum_s (f, e_{j+sn})$, we can conclude

$$|Q_n(f)(t)| \leq \sum_{k=0}^{n-1} \left| \sum_s (f, e_{k+sn}) \right| \leq \sum_{k=0}^{n-1} \sum_s |(f, e_{k+sn})| = \|f\|_a.$$

Thus the assertion holds for case I.

Next we consider case II. Recall that

$$b_j(t) = \sum_s (g, e_{j+sn}) e_{j+sn}(t).$$

Since

$$g(t) = \sum_k -i \cdot \text{sgn}(k) d_k e_k(\pi/n) e_k(t),$$

we obtain

$$|b_j(t)| = \left| \sum_{s=0}^{\infty} d_{j+sn} (-1)^s e_{sn}(t) - \sum_{s=1}^{\infty} d_{-j+sn} (-1)^s e_{-sn}(t) \right|,$$

$$|b_j(0)| = \sum_{s=0}^{\infty} (d_{j+sn} + d_{n-j+sn}) (-1)^s$$

where $0 < j < n$.

Due to the properties of the sequence $d = (d_k)$ we obtain

$$|b_j(0)| \geq d_j - d_n \geq d_j(1 - \alpha_2),$$

$$|b_j(t)| \leq d_j + \sum_{s=1}^{\infty} d_{j+sn} + \sum_{s=1}^{\infty} d_{-j+sn} \leq d_j(1 + \alpha)$$

where $0 < j < n$. This implies

$$|b_j(t)/b_j(0)| \leq (1 + \alpha)/(1 - \alpha_2)$$

where $t \in \mathbb{R}$ and $0 < t < n$. Proceeding as in the proof for case I we can

conclude

$$|Q_n(f)(t)| \leq (1 + \alpha)/(1 - \alpha_2) \|f\|_U.$$

Thus the assertion holds also for case II.

For completeness we consider also case III (trigonometric interpolation). Recall that

$$g(t) = \sum_{k=-m}^m e_k(t) \quad (m = [n/2]),$$

$$b_j(t) = \sum_s (g, e_{j+sn}) e_{j+sn}(t).$$

This implies

$$b_j(t) = e_j(t), \quad 0 < j < n/2, \quad b_j(t) = e_{-n+j}(t), \quad n/2 < j < n$$

for odd n and

$$b_j(t) = e_j(t), \quad 0 < j < n/2, \quad b_j(t) = e_{-n+j}(t), \quad n/2 < j < n,$$

$$b_{n/2}(t) = e_{n/2}(t) + e_{-n/2}(t)$$

for even n . Again we have

$$|b_j(t)/b_j(0)| \leq 1, \quad 0 < j < n.$$

Thus the assertion holds also for case III.

To establish uniform convergence of $Q_n(f)$ for $f \in \mathcal{A}_{2\pi}$ via the Banach-Steinhaus principle [7] we have to verify uniform convergence of $Q_n(f)$ on a dense subset of $\mathcal{A}_{2\pi}$. This will be done in the following section.

3. APPROXIMATION OF THE EXPONENTIALS

The approximation properties of Q_n will first be investigated for the exponential functions e_j .

THEOREM 2 : *For any $k \in \mathbb{Z}$ the asymptotic error relation*

$$\|e_k - Q_n(e_k)\|_\infty = \mathcal{O}(d_{[n/2]}) \quad (n \rightarrow \infty)$$

holds.

Proof : We consider case I. Then we have

$$g(t) = \sum_k d_k e_k(t) = d_0 + \sum_{k=1}^\infty 2 d_k \cos(kt),$$

i.e., $g(t)$ is real valued. This implies $Q_n(\bar{f}) = \overline{Q_n(f)}$.

Since $\bar{e}_k = e_{-k}$ and $Q_n(e_0) = e_0$ we may assume $0 < k \leq n/2$. Recall that

$$b_k(t) = \sum_s d_{k+sn} e_{k+sn}(t) = e_k(t) \sum_s d_{k+sn} e_{sn}(t).$$

Then we obtain

$$\begin{aligned} |e_k(t) - Q_n(e_k)(t)| &= |e_k(t) - b_k(t)/b_k(0)| = \\ &= |b_k(0) - b_k(t) e_{-k}(t)| / |b_k(0)| \\ &\leq \sum_{s \neq 0} 2 d_{k+sn} / d_k \leq \sum_{s=1}^{\infty} 2 d_{s[n/2]} / d_k \leq (2 \alpha / d_k) d_{[n/2]}. \end{aligned}$$

Thus, the assertion holds for case I.

We consider now case II. Then we have

$$g(t) = \sum_k -i \cdot \text{sgn}(k) d_k e_k(-\pi/n) e_k(t) = \sum_{k=1}^{\infty} 2 d_k \sin(k(t - \pi/n)),$$

i.e., $g(t)$ is real valued which implies $Q_n(\bar{f}) = \overline{Q_n(f)}$. Since $\bar{e}_k = e_{-k}$ and $Q_n(e_0) = e_0$, again we may assume $0 < k \leq n/2$. Recall that

$$b_k(t) = -i \cdot e_k(t) \left(\sum_{s=0}^{\infty} d_{k+sn} (-1)^s e_{sn}(t) - \sum_{s=1}^{\infty} d_{-k+sn} (-1)^s e_{-sn}(t) \right).$$

Then we obtain

$$\begin{aligned} |e_k(t) - Q_n(e_k)(t)| &= |e_k(t) - b_k(t)/b_k(0)| = \\ &= |b_k(0) - b_k(t) e_{-k}(t)| / |b_k(0)| \\ &\leq \left| \sum_{s=1}^{\infty} d_{k+sn} (-1)^s (1 - e_{sn}(t)) - \sum_{s=1}^{\infty} d_{-k+sn} (-1)^s (1 - e_{-sn}(t)) \right| / |b_k(0)| \\ &\leq (2 \alpha / ((1 - \alpha_2) d_k)) d_{[n/2]}. \end{aligned}$$

Thus, the assertion holds also for case II.

For the sake of completeness we note that Theorem 2 is trivially true for case III (*trigonometric interpolation*) since in this case we have

$$e_k = Q_n(e_k) \quad (|k| < [n/2]).$$

THEOREM 3 : For any $f \in \mathcal{A}_{2\pi}$ $Q_n(f)$ converges uniformly to f as n tends to infinity :

$$\|f - Q_n(f)\|_{\infty} \rightarrow 0 \quad (n \rightarrow \infty).$$

Proof: Note that the exponentials form a dense subalgebra of $\mathcal{A}_{2\pi}$. In view of Theorem 1 and Theorem 2 the Banach Steinhaus principle is applicable and yields a proof of Theorem 3.

4. QUANTITATIVE ERROR BOUNDS

We use the sequence $d = (d_k)$ to introduce a *smooth* subspace $\mathcal{A}_{2\pi}^d$ of the Wiener algebra $\mathcal{A}_{2\pi}$:

$$\mathcal{A}_{2\pi}^d := \left\{ f \in \mathcal{A}_{2\pi} : \sum_{k \neq 0} |(f, e_k)|/d_k < \infty \right\} .$$

Then we can introduce a linear operator $D : \mathcal{A}_{2\pi}^d \rightarrow \mathcal{A}_{2\pi}$ being defined by

$$D(f) = \sum_{k \neq 0} d_k^{-1} (f, e_k) e_k .$$

The space $\mathcal{A}_{2\pi}^d$ and the operator D are related to the generating function g and the interpolation projector Q_n . It is useful to investigate first the approximation properties in $\mathcal{A}_{2\pi}^d$ of the Fourier partial sum projector F_n .

THEOREM 4 : *If $f \in \mathcal{A}_{2\pi}^d$ then*

$$\|f - F_n(f)\|_{\infty} = o(d_{[n/2]}) \quad (n \rightarrow \infty) .$$

Proof: Note first that

$$F_n(D(f)) = D(F_n(f)) \quad (f \in \mathcal{A}_{2\pi}^d) .$$

Then we have

$$\begin{aligned} |f(t) - F_n(f)(t)| &\leq \sum_{|k| > [n/2]} |(f, e_k)| = \|f - F_n(f)\|_a = \\ &= \sum_{|k| > [n/2]} |(D(f)f, e_k)| d_k \leq \left(\sum_{|k| > [n/2]} |(D(f)f, e_k)| \right) d_{[n/2]} \\ &= \|D(f) - F_n(D(f))\|_a \cdot d_{[n/2]} , \end{aligned}$$

i.e., we have

$$\|f - F_n(f)\|_{\infty} \leq \|f - F_n(f)\|_a \leq \|D(f) - F_n(D(f))\|_a \cdot d_{[n/2]} .$$

Since $D(f) \in \mathcal{A}_{2\pi}$, $\|D(f) - F_n(D(f))\|_a = o(1)(n \rightarrow \infty)$, and the proof of Theorem 4 is complete.

THEOREM 5 : If $f \in \mathcal{A}_{2\pi}^d$ then

$$\|f - Q_n(f)\|_\infty = \mathcal{O}(d_{[n/2]}) \quad (n \rightarrow \infty).$$

Proof : We have

$$\begin{aligned} |f(t) - Q_n(f)(t)| &\leq |f(t) - F_n(f)(t)| + |F_n(f)(t) - Q_n(F_n(f))(t)| \\ &\quad + |Q_n(F_n(f))(t) - Q_n(f)(t)|. \end{aligned}$$

In view of Theorem 1 we have

$$|Q_n(F_n(f))(t) - Q_n(f)(t)| \leq c \|f - F_n(f)\|_a.$$

This implies

$$\begin{aligned} |f(t) - Q_n(f)(t)| &\leq C \|f - F_n(f)\|_a + |F_n(f)(t) - Q_n(F_n(f))(t)| \\ &\leq C \|f - F_n(f)\|_a + \sum_{|k| \leq [n/2]} |(f, e_k)| \cdot \|e_k - Q_n(e_k)\|_\infty \end{aligned}$$

where $C > 0$ is a constant.

In view of Theorem 2 we can conclude

$$\begin{aligned} |f(t) - Q_n(f)(t)| &\leq \\ &\leq C \|f - F_n(f)\|_a + \left(\sum_{0 < |k| \leq [n/2]} |(f, e_k)| \right) \cdot (2\alpha / ((1 - \alpha_2) d_k)) d_{[n/2]} \\ &\leq C \|f - F_n(f)\|_a + (2\alpha / (1 - \alpha_2)) d_{[n/2]} \|D(f)\|_a. \end{aligned}$$

Now an application of Theorem 4 completes the proof of Theorem 5.

The approximation properties of the trigonometric interpolation projector are more closely related to those of the Fourier partial sum projector.

THEOREM 6 : Let Q_n be the projector of trigonometric interpolation. Then for any $f \in \mathcal{A}_{2\pi}^d$ the asymptotic error relation

$$\|f - Q_n(f)\|_\infty = o(d_{[n/2]}) \quad (n \rightarrow \infty)$$

holds.

Proof : As in the proof of Theorem 5 we have

$$\begin{aligned} |f(t) - Q_n(f)(t)| &\leq |f(t) - F_n(f)(t)| + |F_n(f)(t) - Q_n(F_n(f))(t)| \\ &\quad + |Q_n(F_n(f))(t) - Q_n(f)(t)|. \end{aligned}$$

Since Q_n is the trigonometric interpolation projector, we have

$$F_n(f) = Q_n(F_n(f)).$$

Taking into account Theorem 1 we obtain

$$|f(t) - Q_n(f)(t)| \leq C \|f - F_n(f)\|_a$$

where $C > 0$ is a constant. Now an application of Theorem 4 completes the proof.

As an application we consider

$$d_k = k^{-q} \quad (k \in \mathbb{N}, q \in \mathbb{N}, q > 1).$$

Then we have

$$\mathcal{A}_{2\pi}^d := \left\{ f \in \mathcal{A}_{2\pi} : \sum_{k \neq 0} |k|^q |(f, e_k)| < \infty \right\},$$

i.e.,

$$\mathcal{A}_{2\pi}^d = \{f \in \mathcal{C}_{2\pi}^q : D^q f \in \mathcal{A}_{2\pi}\}$$

where $D^q f$ denotes the q -th derivative of f .

THEOREM 7: For any $f \in \mathcal{C}_{2\pi}^q$ with $D^q f \in \mathcal{A}_{2\pi}$ the asymptotic error relation

$$\|f - Q_n(f)\|_\infty = \mathcal{O}(n^{-q}) \quad (n \rightarrow \infty)$$

holds.

If $q = 2r$, $r \in \mathbb{N}$, case I yields the error estimate of Golomb [5] for odd degree periodic splines.

If $q = 2r + 1$, $r \in \mathbb{N}$, case II extends the error estimate of Golomb to even degree periodic midpoint splines.

REFERENCES

- [1] E. W. CHENEY, 1992, Approximation by functions of nonclassical form. In *Approximation Theory, Spline Functions and Applications*, edited by S. P. Singh, *NATO ASI Series C*, **356**, 1-18.
- [2] F.-J. DELVOS, 1985, Interpolation of odd periodic functions on uniform meshes. In *Delay Equations, Approximation, and Applications* (Eds. G. Meinardus, G. Nürnberger), *ISNM 74*, Birkhäuser Verlag, Basel.
- [3] F.-J. DELVOS, 1987, Periodic interpolation on uniform meshes, *Journal of Approximation Theory*, **39**, 71-80.

- [4] F.-J. DELVOS, 1985, Convergence of interpolation on by translation, *In Alfred Haar Memorial Volume, Colloquia Mathematica Societatis Janos Bolyai*, **49** 273-287.
- [5] M. GOLOMB, 1968, Approximation by periodic spline interpolation on uniform meshes, *Journal of Approximation Theory*, **1**, 26-65.
- [6] Y. KATZNELSON, 1976, *An introduction to harmonic analysis*, Dover, New York.
- [7] P.-J. LAURENT, 1972, *Approximation et optimisation*. Herrmann, Paris.
- [8] F. LOCHER, 1981, Interpolation on uniform meshes by translates of one function and related attenuation factors, *Mathematics of Computation*, **37**, 403-416.
- [9] W. QUADE and L. COLLATZ, 1938, Zur Interpolationstheorie der reellen periodischen Funktionen, *Preußische Akademie der Wissenschaften (Math. Phys. Klasse)*, **30**, 383-429.
- [10] G. TOLSTOW, 1976, *Fourier Series*, Dover, New York.