

AN ANALYSIS TECHNIQUE FOR STABILIZED FINITE ELEMENT SOLUTION OF INCOMPRESSIBLE FLOWS*

TOMÁS CHACÓN REBOLLO¹

Abstract. This paper presents an extension to stabilized methods of the standard technique for the numerical analysis of mixed methods. We prove that the stability of stabilized methods follows from an underlying discrete inf-sup condition, plus a uniform separation property between bubble and velocity finite element spaces. We apply the technique introduced to prove the stability of stabilized spectral element methods so as stabilized solution of the primitive equations of the ocean.

Mathematics Subject Classification. 65N30, 76M10.

Received: November 24, 1999. Revised: October 19, 2000.

1. INTRODUCTION AND MOTIVATION

This paper deals with the numerical analysis of the solution of incompressible flow problems by stabilized finite elements. We shall be interested in the Oseen equations (Stokes equations plus a linear transport term), also called in some works “linearized Navier-Stokes equations”.

Stabilized methods provide efficient and computationally cheap techniques to solve incompressible fluids. Historically, these methods have been the object of a specific analysis, different from that of mixed methods. Indeed, the proof of stability is not based upon the existence of a discrete velocity – pressure inf-sup condition, but rather upon specific arguments that strongly rely on the elementwise regularity of finite element functions. Based upon such kind of arguments, the papers of Hugues, Franca and Balestra [21] and Hughes and Franca [20] contained an error analysis that was improved in Brezzi and Douglas [8] and in Pierre [25]. In Franca and Stenberg [15] a general stability and error analysis technique was introduced, which was summarized in Franca, Hugues and Stenberg [16]. Also, the paper of Tobiska and Verfürth [27] develops an analysis of stability and convergence for the solution of Navier-Stokes equations by stabilized methods.

Another way of analysis is suggested by the relationship between stabilized and mixed methods. In Franca and Frey [14] it is proved that the Streamline Upwind/Petrov-Galerkin (SUPG) method is equivalent to the standard mixed method constructed with the mini-element. This equivalence is understood in the sense that both methods yield the same formulation if the degrees of freedom associated to the bubbles are eliminated by static condensation. This equivalence yields the stability of SUPG method from that of the mixed method

Keywords and phrases. Oseen equations, finite elements, mixed methods, stabilized methods, discrete inf-sup condition, spectral methods, primitive equations.

* This research was partially supported by Spanish M. E. C. Project MAR97-1055-C02-02 and by EU HCM ERBCHB ICT 94.1823 Grant.

¹ Departamento de Ecuaciones Diferenciales y Análisis Numérico, Universidad de Sevilla. C/ Tarfia, s/n. 41080 Sevilla, Spain.
e-mail: chacon@numer.us.es

constructed with the mini-element. It is a direct consequence of the fact that this element satisfies the discrete inf-sup condition. Such analysis is essentially performed in Chacón Rebollo [10].

We address in this paper the question of whether this way of analysis may be applied to stabilized methods other than SUPG. We develop a technique for the numerical analysis of stabilized methods that gives a positive answer to that question. Concretely, we prove the existence of an underlying discrete inf-sup condition from which we deduce the stability of stabilized methods. Once this point has been set up, our technique allows to analyze stabilized methods as if they were mixed methods (Th. 1). They appear as internal approximations of a weak formulation, whose stability relies on an inf-sup condition. Then, our analysis may be applied to more complex situations, where we use the tools provided by functional analysis to obtain gains with respect to the standard analysis. We include in this paper two of such applications:

- To prove the stability of a spectral element approximation of the generalized Stokes equations, introduced in Gervasio and Saleri [19]. Here, we obtain L^2 estimates for the pressure, while the standard analysis, used in that paper, allows only to estimate a seminorm of the pressure gradient.
- To solve a linear model of primitive equations of the ocean by stabilized finite elements. For such equations, there is some lack of regularity for the convection term, so that the pressure has only L^p regularity, for some $p \in (1, 2)$. In this case, we obtain L^p estimates for the discrete pressure, and prove convergence in $H^1 \times L^p$ norm to the continuous solution. The standard analysis in this context would be quite difficult to be carried on.

Our analysis may also be applied to nonlinear flows. For instance, in Chacón Rebollo and Domínguez Delgado [11], it is applied to the analysis of the approximation of Navier-Stokes equations by stabilized methods, in parallel to the analysis of their approximations by mixed methods. Stability and error estimates are derived. This analysis also applies to nonlinear stabilized methods, such as the optimal one introduced in Russo [26]. Up to our knowledge, the standard analysis is unable to handle nonlinear stabilization, which turns out to be rather simple to manage with our technique.

Another possible application is the analysis of the solution of Oseen equations by the reduced Q_1/Q_1 stabilized methods introduced in Knobloch and Tobiska [22]. This is a new family of computationally cheap methods that may be directly analyzed with our analysis. In fact, all hypothesis of Theorem 1 are readily proved to be satisfied, using the analysis developed in that paper.

We would like to point out that the analysis technique that we introduce is rather complex from a technical point of view. However, we think that it is worth to be used, as it essentially reduces the difficulties of the analysis of stabilized methods to that of mixed method. Moreover, we have tried to present the technique in a systematic way, so that it may be applied to situations other than the considered here, with relative ease.

The paper is organized as follows. In Section 2 we introduce an abstract discretization of Oseen equations, whose stability is analyzed in Section 3. In Section 4, we apply the abstract theory to stabilized methods. Section 5 is devoted to the analysis of spectral element stabilized methods. Finally, in Section 6 we solve a linear model of primitive equations of the ocean by stabilized finite elements.

2. ABSTRACT DISCRETIZATION

In this Section we introduce an abstract discretization of Oseen equations which is the base of our analysis.

Let us consider a connected bounded domain $\Omega \subset \mathbf{R}^d$ ($d = 2$ or 3), with Lipschitz-continuous boundary Γ . We are given a “driving” velocity field $\mathbf{u} : \Omega \rightarrow \mathbf{R}^d$, that we assume to be divergence-free. Our purpose is to solve numerically the following boundary value problem:

$$\left\{ \begin{array}{ll} \text{Find } \mathbf{y} : \Omega \rightarrow \mathbf{R}^d, \quad p : \Omega \rightarrow \mathbf{R} & \text{such that} \\ \mathbf{u} \cdot \nabla \mathbf{y} - \nu \Delta \mathbf{y} + \nabla p = \mathbf{f}, \quad \nabla \cdot \mathbf{y} = 0 & \text{in } \Omega, \\ \mathbf{y} = 0 & \text{on } \Gamma. \end{array} \right. \quad (1)$$

Here, $\nu > 0$ is the viscosity coefficient, and $\mathbf{f} \in [H^{-1}(\Omega)]^d$ is a given source term. Only homogeneous Dirichlet boundary conditions are considered, in order to not introduce nonessential difficulties in our derivation.

Let us define the bilinear form on $[H_0^1(\Omega)]^d \times [H_0^1(\Omega)]^d$,

$$a(\mathbf{w}, \mathbf{v}) = (\mathbf{u} \cdot \nabla \mathbf{w}, \mathbf{v}) + \nu(\nabla \mathbf{w}, \nabla \mathbf{v}), \quad \forall \mathbf{w}, \mathbf{v} \in [H_0^1(\Omega)]^d, \quad (2)$$

where we denote by (\cdot, \cdot) the L^2 scalar product, either for scalar, vector or tensor functions. If we assume that $\mathbf{u} \in [L^p(\Omega)]^d$ for some $p > d$, and $\nabla \cdot \mathbf{u} = 0$, then $a(\cdot, \cdot)$ is well defined and is continuous and $[H_0^1(\Omega)]^d$ -elliptic; *i.e.*, it verifies

$$a(\mathbf{w}, \mathbf{v}) \leq \mathcal{M}(\mathbf{u}) |\mathbf{w}|_1 |\mathbf{v}|_1, \quad a(\mathbf{v}, \mathbf{v}) \geq \nu |\mathbf{v}|_1^2 \quad \forall \mathbf{v}, \mathbf{w} \in [H_0^1(\Omega)]^d. \quad (3)$$

Here, we have denoted by $|\cdot|_1$ the $[H^1(\Omega)]^d$ seminorm. Also, $\mathcal{M}(\mathbf{u}) = C(\|\mathbf{u}\|_{0,p} + \nu)$ for some constant C appearing from Sobolev injections, where $\|\cdot\|_{0,p}$ denotes the L^p norm.

The form $a(\cdot, \cdot)$ defines a linear bounded operator \mathcal{A} from $[H_0^1(\Omega)]^d$ into $[H^{-1}(\Omega)]^d$, given by

$$\langle \mathcal{A}\mathbf{w}, \mathbf{v} \rangle = a(\mathbf{w}, \mathbf{v}), \quad \forall \mathbf{w}, \mathbf{v} \in [H_0^1(\Omega)]^d.$$

Thus, $\mathcal{A}\mathbf{w} = \mathbf{u} \cdot \nabla \mathbf{w} - \nu \Delta \mathbf{w}$.

The standard mixed formulation of problem (1) reads as follows:

$$\begin{cases} \text{Obtain } (\mathbf{y}, p) \in [H_0^1(\Omega)]^d \times L_0^2(\Omega) \text{ such that} \\ B(\mathbf{y}, p; \mathbf{v}, q) = \langle \mathbf{f}, \mathbf{v} \rangle, \quad \forall (\mathbf{v}, q) \in [H_0^1(\Omega)]^d \times L_0^2(\Omega); \end{cases} \quad (4)$$

where

$$B(\mathbf{y}, p; \mathbf{v}, q) = a(\mathbf{y}, \mathbf{v}) - (p, \nabla \cdot \mathbf{v}) - (\nabla \cdot \mathbf{y}, q).$$

Also, $\langle \cdot, \cdot \rangle$ stands for the $[H^{-1}(\Omega)]^d - [H_0^1(\Omega)]^d$ duality, and $L_0^2(\Omega)$ is the subspace of $L^2(\Omega)$ given by

$$L_0^2(\Omega) = \left\{ q \in L^2(\Omega) \text{ such that } \int_{\Omega} q \, dx = 0 \right\}.$$

The pair of spaces $([H_0^1(\Omega)]^d, L_0^2(\Omega))$ verifies the continuous inf-sup condition (*cf.* Girault and Raviart [17]). Then, due to properties (3), problem (4) has a unique solution that depends continuously on the data \mathbf{f} .

In order to describe our abstract discretization of problem (4) we shall consider two families of subspaces $\{Y_h\}_{h>0}$ and $\{Z_h\}_{h>0}$ of $[H_0^1(\Omega)]^d$ and another family of subspaces $\{M_h\}_{h>0}$ of $L_0^2(\Omega)$, all of them of finite dimension. These spaces may be, for instance, standard finite element spaces. We shall also consider a family of bilinear continuous forms on $[H_0^1(\Omega)]^d \times [H_0^1(\Omega)]^d$, $\{\mathbf{S}_h(\cdot, \cdot)\}_{h>0}$. These forms are assumed to be coercive in H^1 norm on Z_h .

We shall denote by \mathcal{R}_h the “static condensation” operator

$$\mathcal{R}_h : [H^{-1}(\Omega)]^d \rightarrow Z_h,$$

defined as follows. Given $\varphi \in [H^{-1}(\Omega)]^d$, $\mathcal{R}_h(\varphi)$ is the only element of Z_h that satisfies

$$\mathbf{S}_h(\mathcal{R}_h(\varphi), \mathbf{z}_h) = \langle \varphi, \mathbf{z}_h \rangle, \quad \forall \mathbf{z}_h \in Z_h. \quad (5)$$

We discretize problem (4) by

$$\begin{cases} \text{Obtain } (\mathbf{y}_h, p_h) \in Y_h \times M_h \text{ such that} \\ B_h(\mathbf{y}_h, p_h; \mathbf{v}_h, q_h) = F_h(\mathbf{v}_h, q_h), \quad \forall (\mathbf{v}_h, q_h) \in Y_h \times M_h; \end{cases} \quad (6)$$

where

$$\begin{aligned} B_h(\mathbf{w}, r; \mathbf{v}, q) &= B(\mathbf{w}, r; \mathbf{v}, q) - \mathbf{S}_h(\mathcal{R}_h(\mathcal{B}\mathbf{v} + \nabla q), \mathcal{R}_h(\mathcal{A}\mathbf{w} + \nabla r)); \\ F_h(\mathbf{v}, q) &= \langle \mathbf{f}, \mathbf{v} \rangle - \mathbf{S}_h(\mathcal{R}_h(\mathcal{B}\mathbf{v} + \nabla q), \mathcal{R}_h(\mathbf{f})); \end{aligned}$$

where \mathcal{B} denotes the operator

$$\mathcal{B}\mathbf{w} = -\mathbf{u} \cdot \nabla \mathbf{w} + \varepsilon \nu \Delta \mathbf{w}, \quad \forall \mathbf{w} \in [H_0^1(\Omega)]^d,$$

for a given $\varepsilon \in \mathbf{R}$.

We shall use method (6) as an abstract framework to analyze various standard stabilized methods. To describe these methods, we shall consider affine-equivalent finite element spaces, as described in Hughes, Franca and Balestra [21]. Assume that the domain Ω is polyhedral. Let us consider a triangulation \mathcal{T}_h of Ω formed by either simplicial or parallelepipedic elements. We assume that the elements of \mathcal{T}_h are affine-transformed of a reference element K^* (either the unit simplex or parallelepiped), in the sense of Ciarlet [13]. Given an integer number $k \geq 0$, and an element $K \in \mathcal{T}_h$, denote by $P_k(K)$ the space of polynomials of degree smaller than, or equal to, k , defined on K . Also, denote by $Q_k(K)$ the space of polynomials of degree smaller than, or equal to, k , in each variable, defined on K . Denote by $R_k(K)$ either $P_k(K)$, if K is a triangle or tetrahedron, or $Q_k(K)$ if K is a quadrilateral or hexaedron. Given two integer numbers $m \geq 1$, $l \geq 0$, consider the following finite element spaces.

$$Y_h^{(m)} = \left\{ \mathbf{v} \in [H_0^1(\Omega)]^d \mid \mathbf{v}|_K \in [R_m(K)]^d, \forall K \in \mathcal{T}_h \right\}; \quad (7)$$

$$M_h^{(l)} = \left\{ q \in L_0^2(\Omega) \mid q|_K \in R_l(K), \forall K \in \mathcal{T}_h \right\}, \quad (8)$$

or

$$M_h^{(l)} = \left\{ q \in L_0^2(\Omega) \cap C^0(\Omega) \mid q|_K \in R_l(K), \forall K \in \mathcal{T}_h \right\}. \quad (9)$$

We consider the following stabilized methods.

$$\begin{cases} \text{Find } (\mathbf{y}_h, p_h) \in Y_h^{(m)} \times M_h^{(l)} \text{ such that} \\ B_S(\mathbf{y}_h, p_h; \mathbf{v}_h, q_h) = F_S(\mathbf{v}_h, q_h), \quad \forall (\mathbf{v}_h, q_h) \in Y_h^{(m)} \times M_h^{(l)}; \end{cases} \quad (10)$$

where

$$\begin{aligned} B_S(\mathbf{w}, r; \mathbf{v}, q) &= B(\mathbf{w}, r; \mathbf{v}, q) - \sum_{K \in \mathcal{T}_h} \tau_K (\mathcal{B}\mathbf{v} + \nabla q; \mathcal{A}\mathbf{w} + \nabla r)_K; \\ F_S(\mathbf{v}, q) &= \langle \mathbf{f}, \mathbf{v} \rangle - \sum_{K \in \mathcal{T}_h} \tau_K (\mathcal{B}\mathbf{v} + \nabla q, \mathbf{f})_K, \end{aligned}$$

where the τ_K are given stabilizing coefficients, and $(\cdot, \cdot)_K$ denotes the inner product in $[L^2(K)]^d$. When $l = m = 1$, method (10) is independent of the actual value of the coefficient ε , and it is known as Streamline

Upwind/Petrov-Galerkin (SUPG) method. For other values of $m \geq 1$ and $l \geq 0$, when $\varepsilon = -1, 0$ and 1 , method (10) is respectively known as Adjoint stabilized (AdS), generalized SUPG and Galerkin-Least Squares (GaLS) method.

Typically, the coefficients τ_K are continuous functions of the local Péclet number on element K ,

$$Pe_K = \frac{\mathbf{U}_K h_K}{\nu} \quad \text{with} \quad \mathbf{U}_K = \left[\int_K |\mathbf{u}|^p \right]^{1/p};$$

$$\tau_K(Pe_K) = A \frac{h_K}{\mathbf{U}_K} \min(Pe_K, P) = \begin{cases} A \frac{h_K^2}{\nu} & \text{if } Pe_K \leq P, \\ A P \frac{h_K}{\mathbf{U}_K} & \text{if } Pe_K > P; \end{cases} \quad (11)$$

where A is a numerical constant and P is a preset threshold for the Péclet number. This allows on one hand to introduce some suitable stabilization of high frequency components of the transport operator (of order h_K), due to convection dominance (Large Pe_K). Also, this introduces low levels of numerical diffusion (of order h_K^2) in regions where diffusion is dominant (Low Pe_K). On the other hand, this stabilizes the spurious modes of the pressure gradient.

Also, for reasons of computability, in practice the convection velocity \mathbf{u} is replaced in the stabilizing terms by some stable interpolate $\mathbf{u}_h \in Y_h^{(m)}$. We shall assume it so in our analysis.

The standard analysis of stabilized methods, summarized in Franca, Hughes and Stenberg [16], states that SUPG and GaLS methods are stable for any positive coefficients τ_K , and that AdS and generalized SUPG methods are stable if the τ_K are small enough. The obtention of optimal bounds for these coefficients to ensure stability requires the computation of the best constant C_I in the inverse inequality

$$C_I \sum_{K \in \mathcal{T}_h} h_K^2 \|\Delta \mathbf{v}_h\|_K^2 \leq \|\nabla \mathbf{v}_h\|_0^2, \quad \forall \mathbf{v}_h \in Y_h^{(m)}. \quad (12)$$

That analysis applies to either continuous pressures combined with velocities of arbitrary interpolation degree, or to discontinuous pressures combined with high-degree interpolation velocities. Concretely, it holds under the following condition:

$$\text{Either } M_h^{(l)} \subset C^0(\bar{\Omega}), \text{ or } m \geq n, \quad (13)$$

where

$$n = \begin{cases} d & \text{if } \mathcal{T}_h \text{ is formed by triangles or tetrahedra, and} \\ 2 & \text{if } \mathcal{T}_h \text{ is formed by quadrilaterals or hexaedra.} \end{cases}$$

In Tobiska and Verfürth [27] this restriction is removed by introducing in the structure of the method some additional terms that take into account interelement pressure jump terms. However, it seems that method (10), without these jump terms, is not able to stabilize the discretization of discontinuous pressures combined with low-degree velocities.

In this paper we shall analyze methods satisfying condition (13). Our analysis also applies to general discretizations that do not necessarily satisfy this condition. However, its proof requires a rather lengthy derivation that shall appear in a forthcoming paper.

Notice that method (6) applies to general internal approximations of $[H_0^1(\Omega)]^d$ and $L_0^2(\Omega)$, while stabilized methods only apply to approximations by piecewise smooth functions. We are, thus, considering a genuine generalization of stabilized methods.

In the next two Sections we first develop a stability and convergence analysis for the abstract method (6) which extends the standard analysis of mixed methods, and next apply it to analyze the stabilized methods (10).

3. ANALYSIS OF ABSTRACT METHOD

In this section we prove that the stability of the abstract method (6) follows from a discrete inf-sup Brezzi-Babuška condition, similarly to mixed methods.

The stability of abstract method (6), in addition to the inf-sup condition, requires the following hypotheses on the new elements appearing in method (6):

Hypothesis 1. There exists a constant $C_0 > 0$ independent of h such that

$$|\mathbf{y}_h|_1 + |\mathbf{z}_h|_1 \leq C_0 |\mathbf{y}_h + \mathbf{z}_h|_1, \quad \forall \mathbf{y}_h \in Y_h, \mathbf{z}_h \in Z_h, \quad \forall h > 0. \quad (14)$$

Hypothesis 2. There exist two constants $\nu_s > 0, M_s > 0$ such that

$$|\mathbf{S}_h(\mathbf{w}_h, \mathbf{v}_h)| \leq M_s |\mathbf{w}_h|_1 |\mathbf{v}_h|_1, \quad \mathbf{S}_h(\mathbf{v}_h, \mathbf{v}_h) \geq \nu_s |\mathbf{v}_h|_1^2, \quad \forall \mathbf{w}_h, \mathbf{v}_h \in Z_h.$$

Both hypotheses play a crucial role in the obtention of estimates for both velocity and pressure, and thus in the proof of stability of method (6). Hypothesis 1 is a generalization of the well known H_0^1 -orthogonality between piecewise affine and bubble finite elements. Hypothesis 2 is a generalization of the fact that the stabilizing coefficients in (11) are of order h_K^2 .

Let us recall the definition of stability for method (6) (*cf.* Babuška [3], Brezzi [7]):

Definition 1. Method (6) is said to be stable on $Y_h \times M_h$ if there is a constant $\gamma > 0$ independent of h such that for any $(\mathbf{w}, r) \in Y_h \times M_h$,

$$\sup_{\substack{(\mathbf{v}, q) \in Y_h \times M_h \\ (\mathbf{v}, q) \neq (0, 0)}} \frac{B_h(\mathbf{w}, r; \mathbf{v}, q)}{|\mathbf{v}|_1 + \|q\|_0} \geq \gamma (|\mathbf{w}|_1 + \|r\|_0);$$

$$\sup_{\substack{(\mathbf{v}, q) \in Y_h \times M_h \\ (\mathbf{v}, q) \neq (0, 0)}} \frac{B_h(\mathbf{v}, q; \mathbf{w}, r)}{|\mathbf{v}|_1 + \|q\|_0} \geq \gamma (|\mathbf{w}|_1 + \|r\|_0).$$

□

We now state our basic stability result.

Theorem 1. Assume that the pairs of spaces $\{(Y_h + Z_h, M_h)\}_{h>0}$ satisfy a uniform discrete Brezzi-Babuška condition, and that Hypotheses 1 and 2 hold. Assume that at least one of the two following sentences hold:

- i) Z_h and Y_h are orthogonal with respect to the $[H_0^1(\Omega)]^d$ inner product and $\nu_s > 0$, or
- ii) $\nu_s \geq \left(\frac{1-\varepsilon}{2}\right)^2 \nu$, when $\varepsilon \neq 1$, or $\nu_s > 0$ when $\varepsilon = 1$.

Then, the abstract method (6) is stable.

From this theorem we deduce the main result of this paper:

Theorem 2. Assume that the family of triangulations $\{\mathcal{T}_h\}_{h>0}$ is regular. Assume that condition (13) holds. Then, the stabilized method (10) coincides with an abstract method (6) constructed with a finite element space Z_h of bubble functions and a bilinear form S_h , verifying

1. The pairs of spaces $\{Y_h + Z_h, M_h\}_{h>0}$ satisfy a uniform discrete inf-sup condition.
2. The pairs of spaces $\{Y_h, Z_h\}_{h>0}$ satisfy Hypothesis 1.
3. The forms $\{S_h\}_{h>0}$ satisfy Hypothesis 2.

As a consequence,

- GaLS and SUPG methods are stable for any $A > 0$ in (11)
- The general stabilized method (10) is stable if $A \leq A_0 \left(\frac{2}{\varepsilon - 1} \right)^2$, where A_0 is a computable positive constant. In particular, AdS method is stable if $A \leq A_0$, and generalized SUPG method is stable if $A \leq 4A_0$.

Thus, under our analysis, the stability of stabilized methods follows from a discrete inf-sup condition, similarly to mixed methods. We shall prove this result in Section 4. In addition, we shall prove that the constant A_0 depends on the aspect ratio of the grid and on the reference elements of spaces Y_h and M_h , and shall give computable fine estimates for this constant.

Proof of Theorem 1.

Velocity estimate. We shall treat separately cases **i)** and **ii)**.

i) Assume that spaces Z_h and Y_h are orthogonal with respect to the $[H_0^1(\Omega)]^d$ inner product. In this case, all methods (10) coincide, independently of the actual value of ε , as such orthogonality implies $\mathcal{R}_h(\Delta \mathbf{w}) = 0$, $\forall \mathbf{w} \in Y_h$.

Consider a pair $(\mathbf{w}_h, r_h) \in Y_h \times M_h$. Define $\mathbf{c}_h = \mathcal{R}_h(\mathcal{A}\mathbf{w}_h + \nabla r_h)$. As $\mathcal{R}_h(\Delta \mathbf{v}_h) = 0$, then $\mathbf{c}_h = \mathcal{R}_h(-\mathcal{B}\mathbf{w}_h + \nabla r_h)$. Consequently,

$$B_h(\mathbf{w}_h, r_h; \mathbf{w}_h, -r_h) = a(\mathbf{w}_h, \mathbf{w}_h) + \mathbf{S}_h(\mathbf{c}_h, \mathbf{c}_h) \geq \nu |\mathbf{w}_h|_1^2 + \nu_s |\mathbf{c}_h|_1^2.$$

ii) Consider a pair $(\mathbf{w}_h, r_h) \in Y_h \times M_h$. Define $\mathbf{c}_h = \mathcal{R}_h(\mathcal{A}\mathbf{w}_h + \nabla r_h)$. Then,

$$\begin{aligned} B(\mathbf{w}_h, r_h; \mathbf{w}_h, -r_h) &= a(\mathbf{w}_h, \mathbf{w}_h) + \mathbf{S}_h(\mathbf{c}_h, \mathbf{c}_h) + (1 - \varepsilon) \nu \mathbf{S}_h(\mathcal{R}_h(\Delta \mathbf{w}_h), \mathbf{c}_h) \\ &= a(\mathbf{w}_h, \mathbf{w}_h) + \mathbf{S}_h(\mathbf{c}_h, \mathbf{c}_h) - (1 - \varepsilon) \nu (\nabla \mathbf{w}_h, \nabla \mathbf{c}_h) \end{aligned} \quad (15)$$

Due to Hypothesis 1,

$$|(\nabla \mathbf{y}_h, \nabla \mathbf{z}_h)| \leq (1 - \delta_0) |\mathbf{y}_h|_1 |\mathbf{z}_h|_1, \quad \forall \mathbf{y}_h \in Y_h, \mathbf{z}_h \in Z_h, \text{ where } \delta_0 = \frac{2}{C_0^2}. \quad (16)$$

Then, using Young's inequality, (15) implies

$$B(\mathbf{w}_h, r_h; \mathbf{w}_h, -r_h) \geq \tilde{\nu} |\mathbf{w}_h|_1^2 + \tilde{\nu}_s |\mathbf{c}_h|_1^2, \quad (17)$$

where

$$\tilde{\nu} = \nu \left[1 - (1 - \delta_0) \frac{|1 - \varepsilon|}{2} \mu \right], \quad \tilde{\nu}_s = \nu_s - \nu (1 - \delta_0) \frac{|1 - \varepsilon|}{2} \mu^{-1}$$

for any $\mu > 0$. When $\varepsilon = 1$, $\tilde{\nu} = \nu$ and $\tilde{\nu}_s = \nu_s > 0$. When $\varepsilon \neq 1$, we may choose

$$\frac{\nu}{\nu_s} \frac{|1 - \varepsilon|}{2} (1 - \delta_0) < \mu < \frac{2}{|1 - \varepsilon|} (1 - \delta_0)^{-1},$$

and then $\tilde{\nu} > 0$, $\tilde{\nu}_s > 0$.

Denote

$$S = \sup_{\substack{(\mathbf{v}, q) \in Y_h \times M_h \\ (\mathbf{v}, q) \neq (0, 0)}} \frac{B_h(\mathbf{w}_h, r_h; \mathbf{v}, q)}{|\mathbf{v}|_1 + \|q\|_0}.$$

Then, in all cases

$$\tilde{\nu} |\mathbf{w}_h|_1^2 + \tilde{\nu}_s |\mathbf{c}_h|_1^2 \leq (|\mathbf{w}_h|_1 + \|r_h\|_0) S, \quad (18)$$

where for case **i**) we define $\tilde{\nu} = \nu$ and $\tilde{\nu}_s = \nu$.

Pressure estimate. Consider a nonzero element $\mathbf{v}_h \in Y_h$. We have

$$(r_h, \nabla \cdot \mathbf{v}_h) = -B_h(\mathbf{w}_h, r_h; \mathbf{v}_h, 0) + a(\mathbf{w}_h, \mathbf{v}_h) - \mathbf{S}_h(\mathcal{R}_h(\mathcal{B}\mathbf{v}_h), \mathbf{c}_h). \quad (19)$$

Remark that $B_h(\mathbf{w}_h, r_h; -\mathbf{v}_h, 0) \leq S |\mathbf{v}_h|_1$. Observe also that

$$\begin{aligned} \mathbf{S}_h(\mathcal{R}_h(\mathcal{B}\mathbf{v}_h), \mathbf{c}_h) &= \langle \mathcal{B}\mathbf{v}_h, \mathbf{c}_h \rangle = -(\mathbf{u} \cdot \nabla \mathbf{v}_h, \mathbf{c}_h) - \varepsilon \nu (\nabla \mathbf{v}_h, \nabla \mathbf{c}_h) \\ &\leq [\mathcal{M}(\mathbf{u}) + |\varepsilon - 1| \nu] |\mathbf{v}_h|_1 |\mathbf{c}_h|_1. \end{aligned}$$

Consequently,

$$\begin{aligned} (r_h, \nabla \cdot \mathbf{v}_h) &\leq \{S + \mathcal{M}(\mathbf{u}) |\mathbf{w}_h|_1 + [\mathcal{M}(\mathbf{u}) + |\varepsilon - 1| \nu] |\mathbf{c}_h|_1\} |\mathbf{v}_h|_1 \leq \\ &\leq C_1 (S + |\mathbf{w}_h|_1 + |\mathbf{c}_h|_1) |\mathbf{v}_h|_1, \end{aligned}$$

where $C_1 = \max\{1, \mathcal{M}(\mathbf{u}) + |\varepsilon - 1| \nu\}$.

Also, given a nonzero element $\mathbf{z}_h \in Z_h$,

$$\begin{aligned} (r_h, \nabla \cdot \mathbf{z}_h) &= -\langle \nabla r_h, \mathbf{z}_h \rangle = -\mathbf{S}_h(\mathcal{R}_h(\nabla r_h), \mathbf{z}_h) \\ &= \mathbf{S}_h(\mathcal{R}_h(\mathcal{A}\mathbf{w}_h), \mathbf{z}_h) - \mathbf{S}_h(\mathbf{c}_h, \mathbf{z}_h) \\ &\leq \mathcal{M}_s [|\mathcal{R}_h(\mathcal{A}\mathbf{w}_h)|_1 + |\mathbf{c}_h|_1] |\mathbf{z}_h|_1 \\ &\leq \mathcal{M}_s [\nu_s^{-1} |\mathcal{A}\mathbf{w}_h|_{-1} + |\mathbf{c}_h|_1] |\mathbf{z}_h|_1 \\ &\leq C_2 (|\mathbf{w}_h|_1 + |\mathbf{c}_h|_1) |\mathbf{z}_h|_1, \end{aligned} \quad (20)$$

where $C_2 = \mathcal{M}_s \max\{\nu_s^{-1} \mathcal{M}(\mathbf{u}), 1\}$. Then, using Hypothesis 1,

$$\begin{aligned} (r_h, \nabla \cdot (\mathbf{z}_h + \mathbf{v}_h)) &\leq C_3 (S + |\mathbf{w}_h|_1 + |\mathbf{c}_h|_1) (|\mathbf{v}_h|_1 + |\mathbf{z}_h|_1) \\ &\leq C_0 C_3 (S + |\mathbf{w}_h|_1 + |\mathbf{c}_h|_1) |\mathbf{v}_h + \mathbf{z}_h|_1, \end{aligned} \quad (21)$$

where $C_3 = \max\{C_1, C_2\}$. Now, we use the discrete inf-sup condition: There exists a constant $\alpha > 0$ such that

$$\alpha \|q_h\|_0 \leq \sup_{\mathbf{x}_h \in Y_h + Z_h} \frac{(q_h, \nabla \cdot \mathbf{x}_h)}{|\mathbf{x}_h|_1}, \quad \forall q_h \in M_h.$$

Therefore,

$$\|r_h\|_0 \leq C_4 (S + |\mathbf{w}_h|_1 + |\mathbf{c}_h|_1), \quad (22)$$

where $C_4 = \alpha^{-1} C_0 C_3$.

Conclusion. Combining (18) and (22) and applying Young's inequality yields

$$\begin{aligned} \tilde{\nu} |\mathbf{w}_h|_1^2 + \tilde{\nu}_s |\mathbf{c}_h|_1^2 &\leq C_4 S^2 + [(1 + C_4) |\mathbf{w}_h|_1 + C_4 |\mathbf{c}_h|_1] S \\ &\leq \frac{1}{2} [(1 + C_4) \varepsilon_1 |\mathbf{w}_h|_1^2 + C_4 \varepsilon_2 |\mathbf{c}_h|_1^2] \\ &\quad + [C_4 + \frac{1}{2} (1 + C_4) \varepsilon_1^{-1} + \frac{1}{2} C_4 \varepsilon_2^{-1}] S^2, \end{aligned}$$

for any $\varepsilon_1 > 0$, $\varepsilon_2 > 0$. Let us take $\varepsilon_1 = \frac{\tilde{\nu}}{1 + C_4}$, $\varepsilon_2 = \frac{\tilde{\nu}_s}{C_4}$. Then,

$$\tilde{\nu} |\mathbf{w}_h|_1^2 + \tilde{\nu}_s |\mathbf{c}_h|_1^2 \leq C_5^2 S^2, \quad (23)$$

where $C_5 = \left(2C_4 + \frac{(1+C_4)^2}{\tilde{\nu}} + \frac{C_4^2}{\tilde{\nu}_s}\right)^{1/2}$. Thus,

$$|\mathbf{w}_h|_1 \leq \frac{C_5}{\sqrt{\tilde{\nu}}} S, \quad |\mathbf{c}_h|_1 \leq \frac{C_5}{\sqrt{\tilde{\nu}_s}} S. \quad (24)$$

Combining now (23) with (22), we obtain

$$\|r_h\|_0 \leq C_6 S, \quad \text{where } C_6 = C_4 + C_4 C_5 \left(\frac{1}{\sqrt{\tilde{\nu}}} + \frac{1}{\sqrt{\tilde{\nu}_s}}\right). \quad (25)$$

From (24) and (25) we finally deduce

$$S \geq \gamma (|\mathbf{w}_h|_1 + \|r_h\|_0 + |\mathbf{c}_h|_1), \quad \text{where } \gamma = \left\{C_6 + \frac{C_5}{\sqrt{\tilde{\nu}}} + \frac{C_5}{\sqrt{\tilde{\nu}_s}}\right\}^{-1}. \quad (26)$$

The proof of the second inequality in Definition 1 follows from similar arguments. \square

The following result closes the equivalence between discrete inf-sup condition and stability of method (6). Thus, the stability analysis of mixed method and method (6) are fully parallel.

Theorem 3. *Assume that abstract method (6) is stable for some $\nu_s > 0$. Assume that Hypothesis 1 and 2 hold. Then, the pairs of spaces $\{Y_h + Z_h, M_h\}_{h>0}$ satisfy the discrete Brezzi-Babuška condition.*

We omit the proof of this result as it again follows from arguments similar to those used in the proof of Theorem 1.

The stability of form B_h yields the well-posedness of our method, and allows to derive error estimates, similarly to the standard analysis of mixed methods:

Corollary 1. *Under the hypotheses of Theorem 1, problem (6) admits a unique solution $(\mathbf{y}_h, p_h) \in Y_h \times M_h$, that verifies, for some constant $C > 0$,*

$$|\mathbf{y}_h|_1 + \|p_h\|_0 + |\mathbf{z}_h|_1 \leq C \|\mathbf{f}\|_{-1}, \quad (27)$$

and

$$|\mathbf{y} - \mathbf{y}_h|_1 + \|p - p_h\|_0 + |\mathbf{z}_h|_1 \leq C \left[\inf_{\mathbf{v}_h \in Y_h} |\mathbf{y} - \mathbf{v}_h|_1 + \inf_{q_h \in M_h} \|p - q_h\|_0 \right], \quad (28)$$

where $\mathbf{z}_h = \mathcal{R}_h(\mathcal{A}\mathbf{y}_h + \nabla p_h - \mathbf{f})$. \square

Remark 1. From this result, the ‘‘bubble’’ space Z_h appears as a control space for high-frequency components of the residual $\mathcal{A}\mathbf{y}_h + \nabla p_h - \mathbf{f}$. In fact, (28) shows that the high frequency components of the residual which are representable on Z_h , via the condensation operator \mathcal{R}_h , are bounded.

4. APPLICATION TO STABILIZED METHODS

In this section we prove that stabilized methods (10) may be formulated as particular cases of abstract method (6), and then apply the general stability analysis of Section 3.

Our derivation starts from the construction of virtual bubbles developed in Baiocchi *et al.* [4]. Let us recall the main result of that paper, that we adapt to our context. Consider a Hilbert space $(H, (\cdot, \cdot)_H)$. Given a subset B of H of finite dimension, we define the abstract static condensation operator $\mathcal{R} : H' \rightarrow B$ by:

Given $\varphi \in H'$, $\mathcal{R}(\varphi)$ is the only element of B that satisfies

$$(\mathcal{R}(\varphi), \zeta)_H = \langle \varphi, \zeta \rangle, \quad \forall \zeta \in B.$$

Consider also a subspace W of H' of finite dimension endowed with an inner product $(\cdot, \cdot)_W$. With these ingredients, we may re-write the concept of space of virtual bubbles reproducing an operator on W . This is done as follows.

Definition 2. Consider a self-adjoint operator T on W . We say that B is a space of virtual bubbles reproducing T on $(W, (\cdot, \cdot)_W)$ with respect to the inner product $(\cdot, \cdot)_H$ if

$$(\mathcal{R}(\mathbf{w}_1), \mathcal{R}(\mathbf{w}_2))_H = (T\mathbf{w}_2, \mathbf{w}_1)_W, \quad \forall \mathbf{w}_1, \mathbf{w}_2 \in W.$$

A slight modification of the analysis made in Baiocchi et al. [4], proves the following:

Theorem 4. Let H be an infinite-dimensional Hilbert space. Consider a subset W of H' of finite dimension endowed with an inner product $(\cdot, \cdot)_W$. Let B_0 be a finite-dimensional subspace of H satisfying the following property:

$$\text{If } \langle w, b \rangle = 0, \quad \forall b \in B_0, \quad \text{for some } w \in W, \quad \text{then } w = 0. \quad (29)$$

Then, there exists a constant $\mu_0 > 0$ depending only on B_0 such that if $0 < \tau \leq \mu_0$, there exists a finite-dimensional space of virtual bubbles $B \subset H$, that reproduces the operator τI on W with respect to the inner product $(\cdot, \cdot)_H$.

Space B may be constructed as a subspace of $B_0 \oplus N$, N being any subspace of H of dimension $\dim(W)$ such that any function $n \in N$ satisfies

$$\langle w, n \rangle = 0, \quad \forall w \in W, \quad \langle n, b \rangle_H = 0, \quad \forall b \in B_0. \quad (30)$$

On this base, we may perform the analysis of stabilized methods as particular cases of the abstract method (6):

Proof of Theorem 2.

Step 1: Embedding of stabilized method in abstract method.

Let us denote $W_j^* = [R_j(K^*)]^d$ for some integer $j \geq 0$ (recall that K^* denotes the reference element). By Theorem 4, there exists a constant $\mu^* > 0$ such that if $0 < \tau^* \leq \mu^*$, there exists a finite-dimensional space of virtual bubbles $B_j^* \subset [H_0^1(K^*)]^d$, that reproduces the operator $\tau^* I^*$ on W_j^* with respect to the inner product on $[H_0^1(K^*)]^d$.

Indeed, the elements of W_j^* are elements of $[H^{-1}(K^*)]^d$ if we identify the $H_0^1 - H^{-1}$ duality with the L^2 inner product. We consider W_j^* to be endowed with the L^2 inner product. Consider a polynomial function $\Phi : K^* \mapsto \mathbf{R}$ such that $\Phi = 0$ on ∂K^* , and $\Phi > 0$ in $\text{int}(K^*)$. Define the set $B_0 = \{\Phi w^*, | w^* \in W_j^*\}$. Then, B_0 is a subspace of $[H_0^1(K^*)]^d$ of dimension $\dim(W_j^*)$ satisfying property (29): Denote by $(\cdot, \cdot)_*$ the standard inner product on $[L^2(K^*)]^d$. If for some $w^* \in W_j^*$ we have $\langle w^*, b \rangle = (w^*, b)_* = 0$ for any $b \in B_0$, by taking $b = \Phi w^*$ we deduce $w^* = 0$.

Let us now introduce the following elements:

- The bubble finite element space $B_h^{(j)} \subset [H_0^1(\Omega)]^d$ generated by the reference space B_j^* on triangulation \mathcal{T}_h .
- The finite element space $W_h^{(j)} \subset [L^2(\Omega)]^d$ generated by the reference space W_j^* on triangulation \mathcal{T}_h ; i.e.,

$$W_h^{(j)} = \left\{ \mathbf{w} \in [L^2(\Omega)]^d \mid \mathbf{w}|_K \in [R_j(K)]^d, \forall K \in \mathcal{T}_h \right\}. \quad (31)$$

- The inner forms on $[H_0^1(\Omega)]^d$,

$$\mathbf{S}_h(\mathbf{w}, \mathbf{v}) = \sum_{K \in \mathcal{T}_h} \mathbf{S}_K(\mathbf{w}|_K, \mathbf{v}|_K), \quad \text{where} \quad (32)$$

$$\begin{aligned} \mathbf{S}_K(\mathbf{w}, \mathbf{v}) &= \beta_K \int_K (C_K \nabla \mathbf{w}) : \nabla \mathbf{v} \, dx \\ &= \beta_K \sum_{j,k,l=1}^d \int_K (C_K)_{jk} \partial_l w_k \partial_l v_j \, dx, \quad \text{with } \beta_K = \tau^* \tau_K^{-1} h_K^2, \end{aligned} \quad (33)$$

for any $\mathbf{w} = (w_1, \dots, w_d)$, $\mathbf{v} = (v_1, \dots, v_d) \in [H^1(K)]^d$,

where C_K is the matrix defined as follows: There exists a one-to-one affine mapping F_K from the reference element K^* into K . Its equations are of the form $x = A_K x^* + b_K$, where $A_K = \nabla F_K$ is a nonsingular $d \times d$ matrix and $b_K = F_K(0)$ is a vector of \mathbf{R}^d . We define the matrix C_K by $C_K = \frac{1}{h_K^2} A_K A_K^t$.

- The static condensation operator acting on $B_h^{(j)}$ associated to \mathbf{S}_h , that we denote $\mathcal{R}_h^{(j)}$.

Then, we have the following representation lemma for the stabilizing terms:

Lemma 1. *Assume $\mathbf{f} \in [L^2(\Omega)]^d$. Assume $\tau^* \leq \mu^*$. Then, $\forall \mathbf{w}_1, \mathbf{w}_2 \in W_h^{(j)}$,*

$$\sum_{K \in \mathcal{T}_h} \tau_K (\mathbf{w}_2 - \mathbf{f}, \mathbf{w}_1)_K = \mathbf{S}_h \left(\mathcal{R}_h^{(j)}(\mathbf{w}_1), \mathcal{R}_h^{(j)}(\mathbf{w}_2 - \mathbf{f}_h) \right), \quad (34)$$

where \mathbf{f}_h is the L^2 orthogonal projection of \mathbf{f} onto $W_h^{(j)}$.

This lemma is proved in the Appendix.

As a consequence, the stabilized method (10) coincides with the abstract method (6) constructed with spaces $Y_h = Y_h^{(m)}$, $M_h = M_h^{(l)}$, $Z_h = B_h^{(j)}$ (for a fixed positive parameter $\tau^* \leq \mu^*$), with $j = \max\{2m - 1, l - 1\}$, the form \mathbf{S}_h given by (32), and with $\mathcal{R}_h^{(j)}(\mathbf{f})$ replaced by $\mathcal{R}_h^{(j)}(\mathbf{f}_h)$, \mathbf{f}_h being the L^2 orthogonal projection of \mathbf{f} onto $W_h^{(j)}$.

Indeed, let us recall that we are assuming that in the stabilizing terms of method (10) the velocity \mathbf{u} is being replaced by some stable interpolate $\mathbf{u}_h \in Y_h^{(m)}$. Then, for each element $K \in \mathcal{T}_h$, $W_h^{(j)}(K)$ contains the set

$$\left\{ \sum_{K \in \mathcal{T}_h} (\mathbf{u}_h \cdot \nabla \mathbf{v}_h)|_K 1_K, \sum_{K \in \mathcal{T}_h} (\Delta \mathbf{v}_h)|_K 1_K, \sum_{K \in \mathcal{T}_h} (\nabla q_h)|_K 1_K, \text{ for } \mathbf{v}_h \in Y_h^{(m)}, q_h \in M_h^{(l)} \right\},$$

where 1_K denotes the characteristic function of K . Then, it is enough to apply Lemma 1 to obtain the formal embedding.

Step 2: Proof of Hypothesis 1.

This is based on the general result that follows.

Lemma 2. *Assume that the family of triangulations $\{\mathcal{T}_h\}_{h>0}$ is regular. Assume Y_h, Z_h are finite element subspaces of $[H_0^1(\Omega)]^d$ affine equivalent to reference spaces Y^* and Z^* , respectively, satisfying $Y^* \cap Z^* = \{0\}$ and such that Y^* contains the constant functions. Then, the pairs of spaces $\{Y_h, Z_h\}_{h>0}$ satisfy Hypothesis 1.*

This Lemma is proved in the Appendix.

We may now prove that there exists a bubble finite element space $B_h^{(j)}$ such that the pairs $\{Y_h^{(m)}, B_h^{(j)}\}_{h>0}$ satisfy Hypothesis 1 and the representation formula (34) holds. Indeed, replace the function Φ that defines space B_0 by a power Φ^i , for some integer i large enough, to ensure $B_0 \cap [R_m(K^*)]^d = \emptyset$. Let N be any subspace

of dimension $\dim(W_j^*)$ of $[H_0^1(K^*)]^d$ formed by non-polynomial functions that satisfy property (30). Then, the bubble space B_j^* given by Theorem 4 satisfies $B_j^* \cap [R_m(K^*)]^d = \emptyset$. Moreover, as $m \geq 1$, then $[R_m(K^*)]^d$ contains the constant functions and thus Lemma 2 holds.

Step 3: Proof of Hypothesis 2.

Following the derivation of Lemma 2, we obtain

$$\mathbf{S}_h(\mathbf{w}, \mathbf{w}) \geq \nu_{sh} |\mathbf{w}|_1^2, \quad |\mathbf{S}_h(\mathbf{w}, \mathbf{v})| \leq M_{sh} |\mathbf{w}|_1 |\mathbf{v}|_1, \quad \forall \mathbf{w}, \mathbf{v} \in [H_0^1(\Omega)]^d,$$

where $\nu_{sh} = \Lambda \min_{K \in \mathcal{T}_h} \{\beta_K\}$, $M_{sh} = M \max_{K \in \mathcal{T}_h} \{\beta_K\}$, the constants Λ and M being given by (81). Observe that

$$\alpha_2^{-1} \tau^* \leq \beta_K \leq \alpha_1^{-1} \tau^*, \quad \forall K \in \mathcal{T}_h,$$

where α_1 (assuming $h_K \leq 1$) and α_2 are given by

$$\alpha_1 = A \min \left\{ \frac{1}{\nu}, \frac{P}{\|\mathbf{u}\|_{0,p}} \right\}, \quad \alpha_2 = \frac{A}{\nu}. \quad (35)$$

Then, $\nu_{sh} \geq \nu_s$, $M_{sh} \geq M_s$ uniformly in h , with

$$\nu_s = \alpha_2^{-1} \tau^* \Lambda, \quad M_s = \alpha_1^{-1} \tau^* M.$$

Step 4: Discrete inf-sup condition.

This will be based upon the following:

Lemma 3. *Assume that for each $h > 0$ there exists a subspace $Z_h \subset [H_0^1(\Omega)]^d$ and a coercive bilinear form $\hat{\mathbf{S}}_h$ on $[H_0^1(\Omega)]^d$ such that $Z_h \cap Y_h^{(m)} = \{0\}$ and*

$$\sum_{K \in \mathcal{T}_h} h_K^2 \|\nabla q_h\|_{0,K}^2 \leq \hat{C} \hat{\mathbf{S}}_h \left(\hat{\mathcal{R}}(\nabla q_h), \hat{\mathcal{R}}(\nabla q_h) \right) \quad \forall q_h \in M_h^{(l)}, \quad (36)$$

for some constant $\hat{C} > 0$, where $\hat{\mathcal{R}}$ denotes the static condensation operator on Z_h with respect to the form $\hat{\mathbf{S}}_h$.

Assume that the family of forms $\{\hat{\mathbf{S}}_h\}_{h>0}$ is uniformly coercive on $[H_0^1(\Omega)]^d$. Then, under condition (13), the pairs of spaces $\{Y_h^{(m)} + Z_h, M_h^{(l)}\}_{h>0}$ satisfy the discrete inf-sup condition.

Proof of Lemma 3. Under condition (13), it is proved in Franca *et al.* [16] – using the trick of Verfürth [28] – that there exist two constants $C_1 > 0$, $C_2 > 0$ such that $\forall q_h \in M_h^{(l)}$,

$$\sup_{\mathbf{v} \in Y_h^{(m)} - \{0\}} \frac{(\nabla \cdot \mathbf{v}_h, q_h)}{|\mathbf{v}_h|_1} \geq C_1 \|q_h\|_0 - C_2 \left(\sum_{K \in \mathcal{T}_h} h_K^2 \|\nabla q_h\|_{0,K}^2 \right)^{1/2}. \quad (37)$$

Then,

$$\sup_{\mathbf{v} \in Y_h^{(m)} - \{0\}} \frac{(\nabla \cdot \mathbf{v}_h, q_h)}{|\mathbf{v}_h|_1} \geq C_1 \|q_h\|_0 - C_3 \left[\hat{\mathbf{S}}_h \left(\hat{\mathcal{R}}(\nabla q_h), \hat{\mathcal{R}}(\nabla q_h) \right) \right]^{1/2}, \quad (38)$$

where $C_3 = \hat{C} C_2$. Consider a nonzero element $\tilde{\mathbf{v}}_h \in Y_h^{(m)}$, such that

$$|\tilde{\mathbf{v}}_h|_1 = 1, \quad \frac{(\nabla \cdot \tilde{\mathbf{v}}_h, q_h)}{|\tilde{\mathbf{v}}_h|_1} = \sup_{\mathbf{v} \in Y_h^{(m)} - \{0\}} \frac{(\nabla \cdot \mathbf{v}_h, q_h)}{|\mathbf{v}_h|_1}.$$

Denote $\mathbf{z}_h = \hat{\mathcal{R}}(\nabla q_h)$. If $\mathbf{z}_h = 0$, then

$$(\nabla \cdot \tilde{\mathbf{v}}_h, q_h) \geq C_1 \|q_h\|_0.$$

If $\mathbf{z}_h \neq 0$, denote $\tilde{\mathbf{z}}_h = \mathbf{z}_h / |\mathbf{z}_h|_1$. Given a constant $C_4 \geq 0$, from (38) and (36) we obtain

$$\begin{aligned} (\nabla \cdot (\tilde{\mathbf{v}}_h - C_4 \tilde{\mathbf{z}}_h), q_h) &\geq C_1 \|q_h\|_0 - C_3 \left[\hat{\mathbf{S}}_h(\mathbf{z}_h, \mathbf{z}_h) \right]^{1/2} + C_4 \hat{\mathbf{S}}_h(\mathbf{z}_h, \tilde{\mathbf{z}}_h) \\ &\geq C_1 \|q_h\|_0 + \left\{ C_4 \left[\hat{\mathbf{S}}_h(\tilde{\mathbf{z}}_h, \tilde{\mathbf{z}}_h) \right]^{1/2} - C_3 \right\} |\mathbf{z}_h|_1 \left[\hat{\mathbf{S}}_h(\tilde{\mathbf{z}}_h, \tilde{\mathbf{z}}_h) \right]^{1/2} \\ &\geq C_1 \|q_h\|_0 + (C_4 \hat{\nu}^{1/2} - C_3) |\mathbf{z}_h|_1 \left[\hat{\mathbf{S}}_h(\tilde{\mathbf{z}}_h, \tilde{\mathbf{z}}_h) \right]^{1/2}, \end{aligned}$$

where $\hat{\nu}$ is the uniform coerciveness constant of the forms $\hat{\mathbf{S}}_h$. Let us take $C_4 = \hat{\nu}^{-1/2} C_3$. Observe that $\tilde{\mathbf{v}}_h - C_4 \tilde{\mathbf{z}}_h \neq 0$ as $Y_h^{(m)} \cap Z_h = \{0\}$. Define $\tilde{\mathbf{x}}_h = \frac{\tilde{\mathbf{v}}_h - C_4 \tilde{\mathbf{z}}_h}{|\tilde{\mathbf{v}}_h - C_4 \tilde{\mathbf{z}}_h|_1} \in Y_h^{(m)} + Z_h$. Then,

$$(\nabla \cdot \tilde{\mathbf{x}}_h, q_h) \geq \frac{C_1}{1 + C_4} \|q_h\|_0. \quad (39)$$

This completes the proof of Lemma 3. \square

In our case, this result holds with $Z_h = B_h^{(j)}$. Indeed, on one hand $B_h^{(j)} \cap Y_h^{(m)} = \emptyset$. On the other hand, let us take $\mathbf{w}_1 = \mathbf{w}_2 = \nabla q_h$, $\mathbf{f} = 0$ in the representation formula (34). Then,

$$\sum_{K \in \mathcal{T}_h} \tau_K \|\nabla q_h\|_{0,K}^2 = \mathbf{S}_h \left(\mathcal{R}_h^{(j)}(\nabla q_h), \mathcal{R}_h^{(j)}(\nabla q_h) \right). \quad (40)$$

and (36) follows because the coefficients τ_K are of order h_K^2 : $\alpha_1 h_K^2 \leq \tau_K \leq \alpha_2 h_K^2$. Finally, by Step 3 the forms \mathbf{S}_h are uniformly coercive.

Step 5: Conclusion.

We now apply Theorem 1:

- SUPG method corresponds to $m = l = 1$, for any ε . In this case, $Y_h^{(m)}$ and $B_h^{(j)}$ are H_0^1 -orthogonal. Then, from Theorem 1, it is stable for any $A > 0$.
- GaLS method corresponds to $\varepsilon = 1$. Then, from Theorem 1, it is also stable for any $A > 0$.
- In the remaining cases, $Y_h^{(m)}$ and $B_h^{(j)}$ are not necessarily H_0^1 -orthogonal, and $\varepsilon \neq 1$. Let us assume

$$A \leq A_0 \left(\frac{2}{\varepsilon - 1} \right)^2, \quad \text{with } A_0 = \Lambda \mu^*.$$

Take τ^* in the closed interval $\left[\Lambda \Lambda^{-1} \left(\frac{\varepsilon - 1}{2} \right)^2, \mu^* \right]$. As $\tau^* \leq \mu^*$, then all the preceding analysis applies.

Also, $\nu_s = A^{-1} \tau^* \Lambda \nu \geq \left(\frac{\varepsilon - 1}{2} \right)^2 \nu$. Then, from Theorem 1, the general stabilized method (10) is stable. AdS and generalized SUPG methods respectively correspond to $\varepsilon = -1$ and $\varepsilon = 0$ and therefore they are respectively stable if $A \leq \Lambda \mu^*$ and $A \leq 4 \Lambda \mu^*$.

This completes the proof of Theorem 2. \square

Remark 2. Stability of AdS method.

The stability of AdS method may be proved without using the uniform separation property. Indeed, in Baiocchi *et al.* [4] it is proved that there exists a bubble subspace B_h of $[H_0^1(\Omega)]^d$ (not necessarily a finite

element space) such that for any $(\mathbf{v}_h, q_h), (\mathbf{w}_h, r_h) \in Y_h^{(m)} \times M_h^{(l)}$ we have

$$\sum_{K \in \mathcal{T}_h} \tau_K (\mathcal{B}\mathbf{v}_h + \nabla q_h, \mathcal{A}\mathbf{w}_h + \nabla r_h - \mathbf{f})_K = a(\mathcal{R}_h(\mathcal{B}\mathbf{v}_h + \nabla q_h), \mathcal{R}_h(\mathcal{A}\mathbf{w}_h + \nabla r_h - \mathbf{f})), \quad (41)$$

where \mathcal{R}_h is the static condensation operator on B_h with respect to the bilinear form $a(\cdot, \cdot)$. This occurs whenever $\tau_K \leq \mu_K$ for some positive μ_K .

As it is proved in Baiocchi *et al.* [4], this implies that a pair $(\mathbf{y}_h, r_h) \in Y_h^{(m)} \times M_h^{(l)}$ is a solution of AdS method if and only if the pair $(\mathbf{y}_h + \mathbf{b}_h, p_h) \in (Y_h^{(m)} + B_h) \times M_h^{(l)}$, where $\mathbf{b}_h = \mathcal{R}_h(\mathbf{f} - (\mathcal{A}\mathbf{y}_h + \nabla p_h))$, is a solution of the mixed method constructed with spaces $Y_h = Y_h^{(m)} + B_h$ and $M_h = M_h^{(l)}$:

$$B(\mathbf{y}_h + \mathbf{b}_h, p_h; \mathbf{v}_h, q_h) = \langle \mathbf{f}, \mathbf{v}_h \rangle, \quad \forall (\mathbf{v}_h, q_h) \in (Y_h^{(m)} + B_h) \times M_h^{(l)}. \quad (42)$$

Thus, to apply the standard analysis of mixed methods to AdS method it is enough to prove that the family of pairs of spaces $\left\{ Y_h^{(m)} + B_h, M_h^{(l)} \right\}_{h>0}$ satisfy the discrete inf-sup condition. This may be proved by Lemma 3 starting from (41), once we prove that the upper bounds μ_K for the stabilizing coefficients are of order h_K^2 .

Notice that equation (42) provides two control equations for the large and small scale components of ∇p_h . Indeed, (42) is equivalent to

$$(p_h, \nabla \cdot \mathbf{v}_h) = a(\mathbf{y}_h + \mathbf{b}_h, \mathbf{v}_h) - \langle \mathbf{f}, \mathbf{v}_h \rangle, \quad \forall \mathbf{v}_h \in Y_h;$$

$$(p_h, \nabla \cdot \mathbf{z}_h) = a(\mathbf{y}_h + \mathbf{b}_h, \mathbf{z}_h) - \langle \mathbf{f}, \mathbf{z}_h \rangle, \quad \forall \mathbf{z}_h \in Z_h.$$

In the case of stabilized methods other than AdS, we no longer have $\mathcal{B} = \mathcal{A}^*$ in (41). Then, we cannot write the method under the structure (42). In this case, the control equations for ∇p_h are (19) and (20).

Remark 3. Computability of stability bounds.

In Baiocchi *et al.* [4], Section 3.1, a general technique for estimating μ^* is derived. The parameter μ^* depends only on the reference element B_j^* . It must be computed once for each actual space W_j^* associated to a pair $(Y_h^{(m)}, M_h^{(l)})$. In the case of two-dimensional triangular elements, for instance, for $m = l = 1$, this technique yields the estimate $\mu^* = 1/320$. If $m = l = 2$, $\mu^* = 3/5120$.

Also, the constant Λ may be computed from the aspect ratio of the family of triangulations $\{\mathcal{T}_h\}_{h>0}$. Recall that $\Lambda = C_1^2$, C_1 being the constant appearing in (80). From Ciarlet [13], this constant is

$$C_1 = \frac{1}{2h^*\sigma},$$

where h^* is the diameter of the reference element K^* and σ is the aspect ratio of the family,

$$\sigma = \sup_{h>0} \max_{K \in \mathcal{T}_h} \frac{h_K}{\rho_K},$$

ρ_K denoting the internal diameter of element K . This technique to estimate the stability bounds simplifies the standard one, that requires computing the best constant in the inverse inequality (12). This simplification is particularly clear if we observe that σ may be preset “*a priori*” if the triangulations are constructed in order to have

$$\frac{h_K}{\rho_K} \leq \sigma, \quad \forall K \in \mathcal{T}_h, \forall h > 0.$$

Remark 4. Error estimates.

Let us finally make some comments about the obtention of error estimates. One may experiment some concern by the fact that to represent the stabilized method as an abstract method, in the stabilizing terms

the second member \mathbf{f} is replaced by its L^2 interpolate on $W_h^{(j)}$, \mathbf{f}_h . However, we still obtain error estimates of optimal order. This is proved for Navier-Stokes equations in [11]. This proof may readily be adapted to Oseen equations.

5. APPLICATION TO STABILIZED SPECTRAL ELEMENT METHOD

In Gervasio and Saleri [19], a stabilized spectral element (SSE) method for solving the unsteady Navier-Stokes equations is derived. We shall apply here our analysis to the approximation of the Oseen equations by such method. Our main contribution is to prove that the stability of the discretization is due to an underlying discrete inf-sup condition. This allows to obtain L^2 estimates for the pressure. Oseen equations are here considered as a model problem for the linear problems that appear after time discretization of Navier-Stokes equations.

Let us start by describing the discretization of Oseen equations by the SSE method. Assume Ω to be polygonal. Consider a partition \mathcal{T}_h of Ω in parallelograms ($d = 2$) or parallelepipeds ($d = 3$), where h still denotes the largest diameter of the elements of \mathcal{T}_h .

Consider an integer number $N \geq 1$. Denote by $\{\xi_i\}_{i=1}^{N+1}$ and by $\{\omega_i\}_{i=1}^{N+1}$ the nodes and weights of the Gauss-Lobatto Legendre quadrature formulas defined on $(-1, 1)$. Assume, for instance, $d = 3$. For $\hat{u}_N, \hat{v}_N \in Q_N(K^*)$ ($K^* = (-1, 1)^d$ being the reference element), we define the discrete inner product,

$$(\hat{u}_N, \hat{v}_N)_{N, K^*} = \sum_{i,j,k=1}^{N+1} \omega_i \omega_j \omega_k \hat{u}_N(\xi_i, \xi_j, \xi_k) \hat{v}_N(\xi_i, \xi_j, \xi_k);$$

while for $u_N, v_N \in Q_N(K)$ we set

$$(u_h, v_h)_{N, K} = \sum_{i,j,k=1}^{N+1} \omega_i \omega_j \omega_k |\det A_K| u_N(P_{ijk}^{(K)}) v_N(P_{ijk}^{(K)});$$

where $P_{ijk}^{(K)} = F_K(\xi_i, \xi_j, \xi_k)$; $i, j, k = 1, \dots, N+1, \forall K \in \mathcal{T}_h$.

Let us define the space

$$W_{\mathcal{H}} = \{v \in L^2(\Omega) \mid v|_K \in Q_N(K), \forall K \in \mathcal{T}_h\},$$

where $\mathcal{H} = (h, N^{-1})$ is our actual discretization parameter.

Given $u_{\mathcal{H}}, v_{\mathcal{H}} \in W_{\mathcal{H}}$ we set

$$(u_{\mathcal{H}}, v_{\mathcal{H}})_{\mathcal{H}} = \sum_{K \in \mathcal{T}_h} (u_{\mathcal{H}|_K}, v_{\mathcal{H}|_K})_{N, K}; \quad \|u_{\mathcal{H}}\|_{\mathcal{H}} = (u_{\mathcal{H}}, u_{\mathcal{H}})_{\mathcal{H}}^{1/2}.$$

We respectively define the discrete inner products $(\cdot, \cdot)_{N, K^*}$, $(\cdot, \cdot)_{N, K}$ and $(\cdot, \cdot)_{\mathcal{H}}$ on a similar manner for vector functions of $[Q_N(K^*)]^d$, $[Q_N(K)]^d$, and $[W_{\mathcal{H}}]^d$.

We also define the spectral element spaces

$$Y_{\mathcal{H}} = V_{\mathcal{H}}^d \cap [H_0^1(\Omega)]^d, \quad M_{\mathcal{H}} = V_{\mathcal{H}} \cap L_0^2(\Omega),$$

where $V_{\mathcal{H}}$ is defined by

$$V_{\mathcal{H}} = \{v \in C^0(\bar{\Omega}) \mid v|_K \in Q_N(K), \forall K \in \mathcal{T}_h\}.$$

We shall consider the following SSE approximation of Oseen equations (1):

$$\begin{cases} \text{Obtain } (\mathbf{y}_{\mathcal{H}}, p_{\mathcal{H}}) \in Y_{\mathcal{H}} \times M_{\mathcal{H}} \text{ such that} \\ B_{\mathcal{H}}(\mathbf{y}_{\mathcal{H}}, p_{\mathcal{H}}; \mathbf{v}_{\mathcal{H}}, q_{\mathcal{H}}) = F_{\mathcal{H}}(\mathbf{v}_{\mathcal{H}}, q_{\mathcal{H}}), \quad \forall (\mathbf{v}_{\mathcal{H}}, q_{\mathcal{H}}) \in Y_{\mathcal{H}} \times M_{\mathcal{H}}; \end{cases} \quad (43)$$

where

$$\begin{aligned} B_{\mathcal{H}}(\mathbf{w}_{\mathcal{H}}, r_{\mathcal{H}}; \mathbf{v}_{\mathcal{H}}, q_{\mathcal{H}}) &= B(\mathbf{w}_{\mathcal{H}}, r_{\mathcal{H}}; \mathbf{v}_{\mathcal{H}}, q_{\mathcal{H}}) - \sum_{K \in \mathcal{T}_h} \tau_K (\mathcal{B}\mathbf{v}_{\mathcal{H}} + \nabla q_{\mathcal{H}}; \mathcal{A}\mathbf{w}_{\mathcal{H}} + \nabla r_{\mathcal{H}})_{N,K}; \\ F_{\mathcal{H}}(\mathbf{v}_{\mathcal{H}}, q_{\mathcal{H}}) &= \langle \mathbf{f}, \mathbf{v}_{\mathcal{H}} \rangle - \sum_{K \in \mathcal{T}_h} \tau_K (\mathcal{B}\mathbf{v}_{\mathcal{H}} + \nabla q_{\mathcal{H}}, \mathbf{f})_{N,K}. \end{aligned}$$

The essential difference between SSE method and stabilized method (10) is that the L^2 inner products $(\cdot, \cdot)_K$ that appear in (10) in the stabilizing terms are here replaced by the discrete inner products $(\cdot, \cdot)_{N,K}$. In Gervasio and Saleri [19], the discrete inner product $(\cdot, \cdot)_{\mathcal{H}}$ is also used to approximate the integral terms appearing in form B . Here, for simplicity we prefer to consider the above discretization. However, we may extend our analysis to the actual discretization considered in that paper if the pressures are approximated by piecewise polynomials of degree at most $N - 1$ (see Rem. 5).

In Gervasio and Saleri [19], the stabilizing coefficients τ_K are still given by (11), with

$$P = 2 \frac{N^2}{m}, \quad A = \frac{m}{4N^4}, \quad \text{for some } m > 0.$$

The parameter m is determined in that paper in order to obtain uniform-in-time stability of the linear problems that arise after time discretization. We shall simply assume that the stabilizing coefficients τ_K are given by (11).

Our analysis allows to state the following result:

Theorem 5. *Assume the triangulations $\{\mathcal{T}_h\}_{h>0}$ are regular. Then, the SSE method (43) is stable for any $A > 0$ if $\varepsilon = 1$, and for $0 < A < \left(\frac{2}{1-\varepsilon}\right)^2 \hat{A}_0$ if $\varepsilon \neq 1$, where \hat{A}_0 is a computable positive constant.*

As a consequence, if $\mathbf{f} \in [C^0(\Omega)]^d$, problem (43) admits a unique solution that satisfies

$$\|\mathbf{y}_{\mathcal{H}}\|_1 + \|p_{\mathcal{H}}\|_0 \leq C \|\mathbf{f}\|_{C^0}, \quad (44)$$

for some constant $C > 0$ independent of \mathcal{H} .

Proof. We proceed as in the proof of Theorem 2.

Step 1: Embedding of SSE method in abstract method.

Let us define the local interpolation operator $I_N^K : C^0(K) \rightarrow Q_N(K)$ by

$$(I_N^K w)(P_{ijk}^{(K)}) = w(P_{ijk}^{(K)}), \quad i, j, k = 1, \dots, N + 1.$$

Consider the space of piecewise continuous functions on \mathcal{T}_h ,

$$C_{p,h}(\Omega) = \{v \in L^2(\Omega) \mid v|_K \in C^0(K), \forall K \in \mathcal{T}_h\};$$

and define the global interpolation operator $I_{\mathcal{H}} : C_{p,h}(\Omega) \rightarrow W_{\mathcal{H}}$ by

$$(I_{\mathcal{H}} w)|_K = I_N^K(w|_K), \quad \forall K \in \mathcal{T}_h.$$

Observe that $C^0(\bar{\Omega}) \subset C_{p,h}(\Omega)$ and that $I_{\mathcal{H}} w \in V_{\mathcal{H}}$ if $w \in C^0(\bar{\Omega})$.

The following representation formula holds:

Lemma 4. *There exists a finite-dimensional bubble finite element space $Z_{\mathcal{H}} \subset [H_0^1(\Omega)]^d$ such that ,*

$$\mathbf{S}_h(\mathcal{R}_{\mathcal{H}}(I_{\mathcal{H}}\mathbf{v}_1), \mathcal{R}_{\mathcal{H}}(I_{\mathcal{H}}\mathbf{v}_2)) = \sum_{K \in \mathcal{T}_h} \tau_K(\mathbf{v}_1, \mathbf{v}_2)_{N,K}, \quad \forall \mathbf{v}_1, \mathbf{v}_2 \in [C_{p,h}(\Omega)]^d; \quad (45)$$

where \mathbf{S}_h is the bilinear form defined by (32).

This lemma is proved in the Appendix.

As a consequence, for all $\mathbf{w}_{\mathcal{H}}, \mathbf{v}_{\mathcal{H}} \in Y_{\mathcal{H}}; r_{\mathcal{H}}, q_{\mathcal{H}} \in M_{\mathcal{H}}$,

$$\begin{aligned} B_{\mathcal{H}}(\mathbf{w}_{\mathcal{H}}, r_{\mathcal{H}}; \mathbf{v}_{\mathcal{H}}, q_{\mathcal{H}}) &= B(\mathbf{w}_{\mathcal{H}}, r_{\mathcal{H}}; \mathbf{v}_{\mathcal{H}}, q_{\mathcal{H}}) \\ &\quad - \mathbf{S}_h(\mathcal{R}_{\mathcal{H}}(I_{\mathcal{H}}(\mathcal{B}\mathbf{v}_{\mathcal{H}} + \nabla q_{\mathcal{H}})), \mathcal{R}_{\mathcal{H}}(I_{\mathcal{H}}(\mathcal{A}\mathbf{w}_{\mathcal{H}} + \nabla r_{\mathcal{H}}))); \\ F_{\mathcal{H}}(\mathbf{v}_{\mathcal{H}}, q_{\mathcal{H}}) &= \langle \mathbf{f}, \mathbf{v}_{\mathcal{H}} \rangle - \mathbf{S}_h(\mathcal{R}_{\mathcal{H}}(I_{\mathcal{H}}(\mathcal{B}\mathbf{v}_{\mathcal{H}} + \nabla q_{\mathcal{H}})), \mathcal{R}_{\mathcal{H}}(I_{\mathcal{H}}\mathbf{f})). \end{aligned} \quad (46)$$

This occurs because $\mathbf{f} \in [C^0(\Omega)]^d$ and $\mathcal{B}\mathbf{v}_{\mathcal{H}} + \nabla q_{\mathcal{H}}, \mathcal{A}\mathbf{w}_{\mathcal{H}} + \nabla r_{\mathcal{H}} \in [C_{p,h}(\Omega)]^d$.

Steps 2 and 3: Proof of Hypotheses 1 and 2.

Hypotheses 1 and 2 have respectively been proved in the Steps 3 and 4 of the proof of Theorem 2.

Step 4: Discrete inf-sup condition.

Observe that if $r_{\mathcal{H}} \in M_{\mathcal{H}}$, then $I_{\mathcal{H}}(\nabla r_{\mathcal{H}}) = \nabla r_{\mathcal{H}}$, because $I_N^K(q_N) = q_N, \forall q_N \in Q_N(K)$. Then, by (45),

$$\mathbf{S}_h(\mathcal{R}_{\mathcal{H}}(\nabla r_{\mathcal{H}}), \mathcal{R}_{\mathcal{H}}(\nabla r_{\mathcal{H}})) = \sum_{K \in \mathcal{T}_h} \tau_K(\nabla r_{\mathcal{H}}, \nabla r_{\mathcal{H}})_{N,K}, \quad \forall r_{\mathcal{H}} \in M_{\mathcal{H}}.$$

By Bernardi and Maday [5],

$$\|q_N\|_{0,K} \leq \|q_N\|_{N,K} \leq 3 \|q_N\|_{0,K}, \quad \forall q_N \in Q_N(K). \quad (47)$$

These estimates are obtained by affine transportation of similar estimates obtained in the reference element. As the coefficients τ_K are of order h_K^2 , then there exists a constant $C > 0$ such that

$$\sum_{K \in \mathcal{T}_h} h_K^2 \|\nabla r_{\mathcal{H}}\|_{0,K}^2 \leq C \mathbf{S}_h(\mathcal{R}_{\mathcal{H}}(\nabla r_{\mathcal{H}}), \mathcal{R}_{\mathcal{H}}(\nabla r_{\mathcal{H}})).$$

Then, by Lemma 3, the pairs of spaces $\{Y_{\mathcal{H}} + Z_{\mathcal{H}}, M_{\mathcal{H}}\}_{\mathcal{H} > 0}$ satisfy the discrete inf-sup condition.

Step 5: Conclusion.

Following the proof of Theorem 1, we prove that if $\nu_s \geq \frac{(1-\varepsilon)^2}{4} \nu$ when $\varepsilon \neq 1$, or $\nu_s > 0$ when $\varepsilon = 1$, then the form $B_{\mathcal{H}}$ is stable. Then, problem (43) admits a unique solution that satisfies, for some constant $C > 0$,

$$|\mathbf{y}_{\mathcal{H}}|_1 + \|p_{\mathcal{H}}\|_0 + |\mathbf{z}_{\mathcal{H}}|_1 \leq C (\|\mathbf{f}\|_{-1} + |\mathcal{R}_{\mathcal{H}}(I_{\mathcal{H}}\mathbf{f})|_1),$$

where $\mathbf{z}_{\mathcal{H}} = \mathcal{R}_{\mathcal{H}}(I_{\mathcal{H}}(\mathcal{A}\mathbf{y}_{\mathcal{H}}) + \nabla p_{\mathcal{H}})$. As $\mathbf{f} \in [C^0(\Omega)]^d$, using (47),

$$|\mathcal{R}_{\mathcal{H}}(I_{\mathcal{H}}\mathbf{f})|_1 \leq \nu_s^{-1} \|I_{\mathcal{H}}\mathbf{f}\|_0 \leq \nu_s^{-1} \|I_{\mathcal{H}}\mathbf{f}\|_{\mathcal{H}} = \nu_s^{-1} \|\mathbf{f}\|_{\mathcal{H}} \leq C \|\mathbf{f}\|_{C^0}.$$

Thus, estimate (44) follows.

The remaining of the proof is similar to the conclusion of the proof of Theorem 2. \square

Remark 5. A slight modification of the above argument allows to prove an underlying inf-sup condition and thus the stability for a stabilized full spectral element discretization of Oseen equations.

Indeed, let us replace the pressure space $M_{\mathcal{H}}$ by $M_{\mathcal{H}'}$, with $\mathcal{H}' = (h, (N-1)^{-1})$, for $N \geq 2$; *i.e.*, we consider pressures of degree at most $N-1$ elementwise. We consider the following discrete problem:

$$\begin{cases} \text{Obtain } (\mathbf{y}_{\mathcal{H}}, p_{\mathcal{H}'}) \in Y_{\mathcal{H}} \times M_{\mathcal{H}'} \text{ such that} \\ B'_{\mathcal{H}}(\mathbf{y}_{\mathcal{H}}, p_{\mathcal{H}'}; \mathbf{v}_{\mathcal{H}}, q_{\mathcal{H}'}) = F'_{\mathcal{H}}(\mathbf{v}_{\mathcal{H}}, q_{\mathcal{H}'}), \quad \forall (\mathbf{v}_{\mathcal{H}}, q_{\mathcal{H}'}) \in Y_{\mathcal{H}} \times M_{\mathcal{H}'}; \end{cases} \quad (48)$$

where

$$\begin{aligned} B'_{\mathcal{H}}(\mathbf{w}_{\mathcal{H}}, r_{\mathcal{H}'}; \mathbf{v}_{\mathcal{H}}, q_{\mathcal{H}'}) &= \frac{1}{2}[(\mathbf{u} \cdot \nabla \mathbf{w}_{\mathcal{H}}, \mathbf{v}_{\mathcal{H}})_{\mathcal{H}} - (\mathbf{u} \cdot \nabla \mathbf{v}_{\mathcal{H}}, \mathbf{w}_{\mathcal{H}})_{\mathcal{H}}] + \nu (\nabla \mathbf{w}_{\mathcal{H}}, \nabla \mathbf{v}_{\mathcal{H}})_{\mathcal{H}} \\ &\quad - (\nabla \cdot \mathbf{w}_{\mathcal{H}}, q_{\mathcal{H}'})_{\mathcal{H}} - (r_{\mathcal{H}'}, \nabla \cdot \mathbf{v}_{\mathcal{H}})_{\mathcal{H}} \\ &\quad - \sum_{K \in \mathcal{T}_h} \tau_K (\mathcal{B} \mathbf{v}_{\mathcal{H}} + \nabla q_{\mathcal{H}'}; \mathcal{A} \mathbf{w}_{\mathcal{H}} + \nabla r_{\mathcal{H}'})_{N,K}; \\ F'_{\mathcal{H}}(\mathbf{v}_{\mathcal{H}}, q_{\mathcal{H}'}) &= (\mathbf{f}, \mathbf{v}_{\mathcal{H}})_{\mathcal{H}} - \sum_{K \in \mathcal{T}_h} \tau_K (\mathcal{B} \mathbf{v}_{\mathcal{H}} + \nabla q_{\mathcal{H}'}, \mathbf{f})_{N,K}. \end{aligned}$$

Then, our analysis allows to prove that the form $B'_{\mathcal{H}}$ is stable. This holds because the quadrature formula

$$\int_{K^*} g \, dx^* \simeq \sum_{i,j,k=0}^N \omega_i \omega_j \omega_k g(\xi_i, \xi_j, \xi_k)$$

is exact for $g \in \mathbf{Q}_{2N-1}(K^*)$.

6. SOLUTION OF LINEAR PRIMITIVE EQUATIONS

In this section we apply our analysis to the solution of a linear model for the primitive equations of the ocean by a penalty stabilized technique. This model includes the main difficulty of these equations: The vertical convection is degenerated. This makes the pressure to be only in some space L^p for $1 < p < 2$. We prove a discrete inf-sup condition in this norm, and prove the convergence of the approximated solutions to a weak solution of the continuous problem.

To describe our model equations, let us consider a connected 2D bounded domain $\omega \subset \mathbf{R}^2$, and a piecewise continuous function $D : \bar{\omega} \rightarrow \mathbf{R}$ such that $D(\mathbf{x}) > 0$, $\forall \mathbf{x} = (x_1, x_2) \in \omega$. This function represents the sea depth. We consider the domain

$$\Omega = \{(\mathbf{x}, z) \in \mathbf{R}^3 \mid \mathbf{x} \in \omega, -D(\mathbf{x}) < z < 0\},$$

which is intended to represent a piece of the ocean with flat surface. To avoid some technical complexities, we shall assume that ω is polygonal and D is piecewise affine on some triangulation of $\bar{\omega}$, so that Ω is polyhedral. Our analysis can be extended to piecewise C^1 depth functions, similarly to the analysis of the approximation of primitive equations by mixed methods (*cf.* Chacón Rebollo and Guillén González [12]).

We assume the domain Ω to be Lipschitz continuous. This occurs, for instance, if the normal derivative of D satisfies $\frac{\partial D}{\partial n} \leq \alpha$ for some $\alpha < 0$ a. e. on the part of $\partial\omega$ where $D = 0$. Notice that D may be zero partially or totally on $\partial\omega$. Also, that we allow the sea bottom to have vertical walls when D has a jump, and sidewalls if $D > 0$ on a part of $\partial\omega$.

We also consider the following subsets of $\partial\Omega$:

$$\Gamma_s = \{(\mathbf{x}, 0) \in \mathbf{R}^3 \mid \mathbf{x} \in \bar{\omega}\}, \quad (\text{sea surface}),$$

$$\Gamma_b = \partial\Omega - \Gamma_s \quad (\text{sea bottom and, eventually, sidewalls}).$$

We assume known a convection velocity $\mathbf{W} = (w_1, w_2, w_3)$ on Ω , such that

$$\begin{cases} \mathbf{w} = (w_1, w_2) \in [H^1(\Omega)]^2, w_3 \in L^2(\Omega), \\ \nabla \cdot \mathbf{W} = 0 \text{ in } \Omega, w_3|_{\Gamma_s} = 0, w_3 \cdot n_3|_{\Gamma_b} = 0, \mathbf{w}|_{\Gamma_b} = 0, \end{cases} \quad (49)$$

where n_3 denotes the third component of the outward normal to $\partial\Omega$, $\mathbf{n} = (n_1, n_2, n_3)$. We are thus forcing the incompressibility of the sea water (Boussinesq's hypothesis). The first boundary condition means that we assume the sea surface to not move in the vertical direction (rigid lid hypothesis), while the second and third ones are rather technical boundary conditions, meaning that we treat the whole Γ_b as a solid wall.

We also assume known a distributed source term \mathbf{f} , representing the effects of temperature, salinity and Coriolis force (assumed to be constant on the whole domain for simplicity), and a "surface wind tension" \mathbf{g} . We set the following problem:

$$\begin{cases} \text{Obtain } \mathbf{y} : \bar{\Omega} \rightarrow \mathbf{R}^2, \mathbf{y} = (y_1, y_2), & \text{(horizontal velocity)} \\ \text{and } p : \omega \rightarrow \mathbf{R} \text{ such that} & \text{(surface pressure)} \\ \mathbf{W} \cdot \nabla \mathbf{y} - \nu \Delta \mathbf{y} + \nabla_H p = \mathbf{f} & \text{in } \Omega, \\ \nabla_H \cdot \langle \mathbf{y} \rangle = 0 & \text{in } \omega, \\ \mathbf{y}|_{\Gamma_b} = 0, \quad \nu \frac{\partial \mathbf{y}}{\partial n}|_{\Gamma_s} = \mathbf{g}. \end{cases} \quad (50)$$

Here, $\nabla_H = (\partial_1, \partial_2)$ stands for the horizontal gradient, and the symbols $\langle \cdot \rangle$ denote vertical mean,

$$\langle \mathbf{y} \rangle(\mathbf{x}) = \int_{-D(\mathbf{x})}^0 \mathbf{y}(\mathbf{x}, z) dz, \quad \text{for } \mathbf{x} \in \omega.$$

In this problem the surface pressure p acts as a Lagrange multiplier associated to the condition $\nabla_H \cdot \langle \mathbf{y} \rangle = 0$.

Problem (50) is a reduced version of a linear model of the primitive equations of the ocean (introduced in Lions, Temam and Wang [24]), that reads as follows:

$$\begin{cases} \text{Obtain a velocity field } (\mathbf{y}, y_3) : \bar{\Omega} \rightarrow \mathbf{R}^3, \\ \text{and a pressure } P : \Omega \rightarrow \mathbf{R} \text{ such that} \\ \mathbf{W} \cdot \nabla \mathbf{y} - \nu \Delta \mathbf{y} + \nabla_H P = \mathbf{f} & \text{in } \Omega, \\ \nabla \cdot (\mathbf{y}, y_3) = 0 & \text{in } \Omega, \\ \partial_3 P = -\rho g & \text{in } \Omega, \\ \mathbf{y}|_{\Gamma_b} = 0, \quad \nu \frac{\partial \mathbf{y}}{\partial n}|_{\Gamma_s} = \mathbf{g}, \\ y_3 \cdot n_3|_{\Gamma_b} = 0, \quad y_3|_{\Gamma_s} = 0. \end{cases} \quad (51)$$

Here, ρ represents the sea water density, assumed to be constant, and g the acceleration of the gravity.

This model is formally obtained from the Navier-Stokes equations by neglecting in the vertical momentum equation all forces (convection, diffusion and Coriolis) but the gravity. This leads to the hydrostatic pressure approximation. A rigorous derivation of this approximation is found in Besson and Laydi [2], as an asymptotic limit as the ratio between vertical and horizontal dimensions tends to zero. The physically meaningful – nonlinear – problem would be to find a "fixed point" of equations (50), in the sense that $\mathbf{y} = \mathbf{w}$. This justifies the choice of regularity and boundary conditions satisfied by \mathbf{w} (see (49)).

Equations (50) may be viewed as a model problem for the nonlinear primitive equations, much as the Oseen equations are a linear model for the Navier-Stokes equations.

Remark 6. Problems (50) and (51) are equivalent. The key point for this equivalence is the following: If a horizontal velocity $\mathbf{y} = (y_1, y_2) \in [H^1(\Omega)]^2$ satisfies $\mathbf{y}|_{\Gamma_b} = 0$, then

$$\langle \nabla_H \cdot \mathbf{y} \rangle = \nabla_H \cdot \langle \mathbf{y} \rangle.$$

As a consequence, there exists a vertical velocity $y_3 \in L^2(\Omega)$ such that

$$\nabla \cdot (\mathbf{y}, y_3) = 0 \text{ in } \Omega, \quad y_3|_{\Gamma_s} = 0 \text{ and } y_3 \cdot \mathbf{n}_3|_{\Gamma_b} = 0$$

if and only if

$$y_3(\mathbf{x}, x_3) = \int_{x_3}^0 \nabla_H \cdot \mathbf{y}(\mathbf{x}, s) \, ds \quad \text{in } \Omega, \quad (52)$$

and

$$\nabla_H \cdot \langle \mathbf{y} \rangle = 0 \text{ in } \omega.$$

This allows to eliminate the vertical velocity y_3 from problem (51). Also, the condition $\partial_3 P = -\rho g$ allows to recover the pressure P from the surface pressure p , by

$$P(\mathbf{x}, z) = \rho g z + p(\mathbf{x}). \quad (53)$$

A rigorous proof of this equivalence may be found in Lewandowski [23].

To give a variational formulation to problem (51), let us define the spaces

$$V_k = \{\mathbf{v} = (v_1, v_2) \in [W^{1,k}(\Omega)]^2 \mid \mathbf{v}|_{\Gamma_b} = 0\} \quad \text{for } k \geq 1, \text{ integer};$$

$$L_D^\alpha(\omega) = \{q : \omega \rightarrow \mathbf{R} \text{ measurable such that } \int_\omega D(\mathbf{x}) |q(\mathbf{x})|^\alpha \, dx < +\infty\} \text{ for } \alpha \geq 1;$$

$$L_{D,0}^\alpha(\omega) = L_D^\alpha(\omega)/\mathbf{R} \text{ (quotient space).}$$

Spaces $L_D^\alpha(\omega)$ and $L_{D,0}^\alpha(\omega)$ are Banach spaces – reflexive if $1 < \alpha < +\infty$ –, respectively endowed with the norms

$$\|q\|_{L_D^\alpha(\omega)} = \left[\int_\omega D(\mathbf{x}) |q(\mathbf{x})|^\alpha \, dx \right]^{1/\alpha},$$

$$\|q\|_{L_{D,0}^\alpha(\omega)} = \inf_{c \in \mathbf{R}} \|q + c\|_{L_D^\alpha(\omega)}.$$

Space $L_D^\alpha(\omega)$ is isomorphic, and, more specifically, isometric, to the space

$$L^\alpha(\partial_3, \Omega) = \{q \in L^\alpha(\Omega) \text{ such that } \partial_3 q = 0\}.$$

Indeed, we identify each $q \in L_D^\alpha(\omega)$ with its extension to Ω as a constant function with respect to the x_3 variable. Then, we have $\|q\|_{L_D^\alpha(\omega)} = \|q\|_{L^\alpha(\Omega)}$.

Moreover, if we consider the space $L_0^\alpha(\partial_3, \Omega) = L^\alpha(\partial_3, \Omega)/\mathbf{R}$, then $L_{D,0}^\alpha(\omega)$ and $L_0^\alpha(\partial_3, \Omega)$ also are isomorphic, and $\|q\|_{L_{D,0}^\alpha(\omega)} = \|q\|_{L_0^\alpha(\partial_3, \Omega)}$, $\forall q \in L_{D,0}^\alpha(\omega)$.

We further assume $\mathbf{f} \in V_2'$ and $\mathbf{g} \in [H^{-1/2}(\Gamma_s)]^d$, the dual space of $[H^{1/2}(\Gamma_s)]^d$. This space is well defined as Γ_s is C^∞ .

We consider the following weak formulation of problem (51):

$$\begin{cases} \text{Obtain } (\mathbf{y}, p) \in V_2 \times L_{D,0}^{3/2}(\omega) \text{ such that} \\ B^{(PE)}(\mathbf{y}, p; \mathbf{v}, q) = F(\mathbf{v}); \quad \forall (\mathbf{v}, q) \in V_4 \times L_{D,0}^2(\omega); \end{cases} \quad (54)$$

where

$$\begin{aligned} B^{(PE)}(\mathbf{y}, p; \mathbf{v}, q) &= \langle \mathbf{W} \cdot \nabla \mathbf{y}, \mathbf{v} \rangle_{V_4' - V_4} + \nu (\nabla \mathbf{y}, \nabla \mathbf{v})_\Omega - (p, \nabla_H \cdot \langle \mathbf{v} \rangle)_\omega \\ &\quad - (\nabla_H \cdot \langle \mathbf{y} \rangle, q)_\omega, \end{aligned}$$

$$F(\mathbf{v}) = \langle \mathbf{f}, \mathbf{v} \rangle_{V_2' - V_2} + \langle \mathbf{g}, \mathbf{v} \rangle_{H^{-1/2}(\Gamma_s) - H^{1/2}(\Gamma_s)}.$$

This form is well defined, due to the following:

Lemma 5. *The following statements hold.*

- i) Consider a function $\mathbf{W} = (\mathbf{w}, w_3) \in V_2 \times L^2(\Omega)$ such that $\nabla \cdot \mathbf{W} = 0$, $w_3|_{\Gamma_s} = 0$. Then, $\forall \mathbf{u} \in V_2$, $\mathbf{W} \cdot \nabla \mathbf{u} \in V_k'$ for $k \geq 3$, and

$$\|\mathbf{W} \cdot \nabla \mathbf{u}\|_{V_k'} \leq \hat{C}_k |\mathbf{w}|_{1,\Omega} |\mathbf{u}|_{1,\Omega}, \quad (55)$$

for some constant $\hat{C}_k > 0$.

- ii) If $\mathbf{w} \in V_k$ for some $k \geq 1$, then $\langle \mathbf{w} \rangle \in [W^{1,k}(\omega)]^2$ and $\partial_i \langle \mathbf{w} \rangle = \langle \partial_i \mathbf{w} \rangle$, $i = 1, 2$.

Proof. i) Observe that, given $\mathbf{w} \in V_2$, and $w_3 \in L^2(\Omega)$ such that $\partial_3 w_3 = -\nabla_H \cdot \mathbf{w}$, and $w_3|_{\Gamma_s} = 0$ we have

$$w_3(\mathbf{x}, x_3) = \int_{x_3}^0 \nabla_H \cdot \mathbf{w}(\mathbf{x}, s) ds. \text{ Thus,}$$

$$\|w_3\|_{0,\Omega} + \|\partial_3 w_3\|_{0,\Omega} \leq C_1 |\mathbf{w}|_{1,\Omega}, \quad (56)$$

for some constant $C_1 > 0$.

Now, if \mathbf{W} is smooth, we see by integrations by parts that for $\mathbf{u} \in V_2$ and $\mathbf{v} \in V_k$,

$$\int_{\Omega} (\mathbf{W} \cdot \nabla \mathbf{u}) \cdot \mathbf{v} dx dx_3 = \int_{\Omega} [(\mathbf{w} \cdot \nabla_H \mathbf{u}) \cdot \mathbf{v} - \partial_3 w_3 \mathbf{u} \cdot \mathbf{v} - w_3 \mathbf{u} \cdot \partial_3 \mathbf{v}] dx dx_3.$$

Then, we may define the duality $\langle \mathbf{W} \cdot \nabla \mathbf{u}, \mathbf{v} \rangle$ by

$$\langle \mathbf{W} \cdot \nabla \mathbf{u}, \mathbf{v} \rangle = \int_{\Omega} [(\mathbf{w} \cdot \nabla_H \mathbf{u}) \cdot \mathbf{v} - \partial_3 w_3 \mathbf{u} \cdot \mathbf{v} - w_3 \mathbf{u} \cdot \partial_3 \mathbf{v}] dx dx_3. \quad (57)$$

Using (56),

$$\begin{aligned} |\langle \mathbf{W} \cdot \nabla \mathbf{u}, \mathbf{v} \rangle| &\leq C_2 (\|\mathbf{w}\|_{0,4;\Omega} |\mathbf{u}|_{1,\Omega} \|\mathbf{v}\|_{0,4;\Omega} + |\mathbf{w}|_{1,\Omega} \|\mathbf{u}\|_{0,4;\Omega} \|\mathbf{v}\|_{0,4;\Omega} \\ &\quad + \|\mathbf{w}\|_{0,\Omega} \|\mathbf{u}\|_{0,6;\Omega} \|\partial_3 \mathbf{v}\|_{0,3;\Omega}) \leq \hat{C}_k |\mathbf{w}|_{1,\Omega} |\mathbf{u}|_{1,\Omega} |\mathbf{v}|_{1,k;\Omega}. \end{aligned} \quad (58)$$

This proves that $\mathbf{W} \cdot \nabla \mathbf{u} \in V_k'$. Next, consider a field $\mathbf{W} = (\mathbf{w}, w_3) \in V_2 \times L^2(\Omega)$ with $\nabla \cdot \mathbf{W} = 0$, $w_3|_{\Gamma_s} = 0$. Then, there exists a sequence $\{\mathbf{w}_n\}_{n \geq 1} \subset [\mathcal{D}(\bar{\Omega})]^2$ such that $\mathbf{w}_n = 0$ on Γ_b , which converges to \mathbf{w} in V_2 . This is proved by a standard argument (for instance, by symmetrization with respect to Γ_s) using that $\partial \Omega$ is Lipschitz-continuous. Let $\mathbf{W}_n = (\mathbf{w}_n, w_{3n})$, with $w_{3n}(\mathbf{x}, x_3) = \int_{x_3}^0 \nabla_H \cdot \mathbf{w}_n(\mathbf{x}, s) ds$.

Following Dautray and Lions [18], Chapter XXI, we may ensure that if a function $z \in L^2(\Omega)$ is such that $\partial_3 z \in L^2(\Omega)$, then the trace of z on Γ_s belongs to $H^{1/2}(\Gamma_s)$. Moreover, a Poincaré inequality holds if $z|_{\Gamma_s} = 0$:

$$\|z\|_{0,\Omega} \leq C_3 \|\partial_3 z\|_{0,\Omega},$$

for some constant $C_3 > 0$. Therefore,

$$\|w_3 - w_{3n}\|_{0,\Omega} \leq C_3 \|\nabla_H \cdot (\mathbf{w} - \mathbf{w}_n)\|_{0,\Omega},$$

and w_{3n} converges to w_3 in $L^2(\Omega)$. Thus, we may pass to the limit in the r.h.s. of (57), and define $\mathbf{W} \cdot \nabla \mathbf{u}$ as a linear form on V_k . Now, passing to the limit in (58) we deduce $\mathbf{W} \cdot \nabla \mathbf{u} \in V_k'$ and estimate (55).

ii) Consider $\mathbf{w} \in V_k$. As $\mathbf{w} \in [L^k(\Omega)]^2$, one readily proves $\langle \mathbf{w} \rangle \in [L^k(\omega)]^2$. Also, let $\varphi \in \mathcal{D}(\omega)$. Then, if \mathbf{w} is smooth, for $i = 1, 2$,

$$\int_{\omega} \langle \partial_i \mathbf{w} \rangle(\mathbf{x}) \varphi(\mathbf{x}) \, d\mathbf{x} = \int_{\Omega} \partial_i \mathbf{w}(\mathbf{x}, x_3) \varphi(\mathbf{x}) \, d\mathbf{x} \, dx_3 = \quad (59)$$

$$= \int_{\partial\Omega} n_i \mathbf{w} \varphi \, d(\partial\Omega) - \int_{\Omega} \mathbf{w}(\mathbf{x}, x_3) \partial_i \varphi(\mathbf{x}) \, d\mathbf{x} \, dx_3 = \quad (60)$$

$$= - \int_{\omega} \langle \mathbf{w} \rangle(\mathbf{x}) \partial_i \varphi(\mathbf{x}) \, d\mathbf{x}, = - \int_{\omega} \partial_i \langle \mathbf{w} \rangle(\mathbf{x}) \varphi(\mathbf{x}) \, d\mathbf{x}, \quad (61)$$

as $n_i = 0$ on Γ_s and $\mathbf{w} = 0$ on Γ_b . Thus, $\partial_i \langle \mathbf{w} \rangle = \langle \partial_i \mathbf{w} \rangle$ and $\mathbf{w} \in [W^{1,k}(\omega)]^2$.

If \mathbf{w} is any element of V_k , the same results follows from a density argument similar to that of the proof of statement i) above. \square

Remark 7. Any solution (\mathbf{y}, p) of problem (54) is a weak solution of problem (50) in the distribution sense. Furthermore, if we recover the vertical velocity y_3 by (52), and the physical pressure P by (53), then the couple $((\mathbf{y}, y_3), P)$ is a solution of problem (51) in the distribution sense.

We shall discretize problem (54) by a penalty stabilized method, of Brezzi and Pitkäranta's kind (*cf.* [9]). Consider a triangulation \mathcal{C}_h of $\bar{\omega}$ such that D is affine on each triangle $T \in \mathcal{C}_h$. Consider also a partition \mathcal{P}_h of $\bar{\Omega}$ by sets of the form

$$P_T = \{(\mathbf{x}, x_3) \in \mathbf{R}^3, \text{ such that } \mathbf{x} \in T, -D(\mathbf{x}) \leq x_3 \leq 0\} \text{ for some triangle } T \in \mathcal{C}_h.$$

Notice that if a triangle $T \in \mathcal{C}_h$ is not adjacent to $\partial\omega$, or if it is adjacent to $\partial\omega$ and $D > 0$ on \bar{T} , then its associated set P_T is a triangular prism with upper base $T \times \{0\}$ and possibly non-horizontal lower base. However, if T is adjacent to $\partial\omega$ and $D = 0$ on a part of ∂T , then P_T is a non-prismatic polyhedron.

We shall consider a triangulation \mathcal{T}_h of Ω constructed by subdividing each element \mathcal{P}_h into tetrahedra. Let us define the finite element spaces,

$$V_h = \{v_h \in C^0(\bar{\Omega}) \mid v_h|_K \in \mathbf{P}_1(K), \forall K \in \mathcal{T}_h\}; \quad (62)$$

$$Y_h = \{\mathbf{v}_h \in V_h^2 \mid v_h|_{\Gamma_b} = 0\};$$

$$\tilde{N}_h = \{q_h \in C^0(\bar{\omega}) \mid q_h|_T \in \mathbf{P}_1(T), \forall T \in \mathcal{C}_h\}; \quad N_h = \tilde{N}_h/R.$$

We introduce the following discretization of (54):

$$\left\{ \begin{array}{l} \text{Obtain } (\mathbf{y}_h, p_h) \in Y_h \times N_h \text{ such that} \\ B_h^{(PE)}(\mathbf{y}_h, p_h; \mathbf{v}_h, q_h) = F(\mathbf{v}_h); \quad \forall (\mathbf{v}_h, q_h) \in Y_h \times N_h; \end{array} \right. \quad (63)$$

where

$$\begin{aligned} B_h^{(PE)}(\mathbf{u}_h, r_h; \mathbf{v}_h, q_h) &= B^{(PE)}(\mathbf{u}_h, r_h; \mathbf{v}_h, q_h) + \sum_{K \in \mathcal{T}_h} \tau_K^{(c)} (\mathbf{W}_h \cdot \nabla \mathbf{u}_h, \mathbf{W}_h \cdot \nabla \mathbf{v}_h)_K \\ &\quad - \sum_{T \in \mathcal{C}_h} \tau_T^{(p)} (\nabla_H r_h, \nabla_H q_h)_T. \end{aligned}$$

The stabilizing coefficients for convection $\tau_K^{(c)}$ are assumed to be still given by (11). This will provide some stabilization of the convective derivative. Also, to ensure the stability of the pressure discretization we shall assume that the stabilizing coefficients for pressure $\tau_T^{(p)}$ satisfy the following condition: There exist two constants $\beta_1 > 0$, $\beta_2 > 0$ such that

$$\beta_1 h_T^2 \frac{\int_T D \, d\mathbf{x}}{|T|} \leq \tau_T^{(p)} \leq \beta_2 h_T^2 \frac{\int_T D \, d\mathbf{x}}{|T|}, \quad \forall T \in \mathcal{C}_h. \quad (64)$$

Observe that these inequalities make sense as we assume $D > 0$ on ω . In the stabilizing terms of (63), we replace the convection velocity $\mathbf{W} = (\mathbf{w}, w_3)$ by some interpolate $\mathbf{W}_h = (\mathbf{w}_h, w_{3h}) \in Y_h \times V_h$, satisfying for some constant $C > 0$,

$$|\mathbf{W}_h|_1 \leq C |\mathbf{w}|_1. \quad (65)$$

We now state the main result of this section.

Theorem 6. *Assume the convection velocity $\mathbf{W} = (\mathbf{w}, w_3)$ lies in the space $V_2 \times L^2(\Omega)$ and verifies $\nabla \cdot \mathbf{W} = 0$, $w_3|_{\Gamma_s} = 0$. Assume the triangulations $\{\mathcal{T}_h\}_{h>0}$ are regular. Then, the following statements hold.*

- i) *Problem (63) admits a unique solution $(\mathbf{y}_h, p_h) \in Y_h \times N_h$ which is bounded in $V_2 \times L_{D,0}^{3/2}(\omega)$.*
- ii) *The sequence $\{(\mathbf{y}_h, p_h)\}_{h>0}$ contains a subsequence which is weakly convergent in $V_2 \times L_{D,0}^{3/2}(\omega)$ to a solution of (54) satisfying the estimate*

$$|\mathbf{y}|_1 + \|p\|_{L_{D,0}^{3/2}(\omega)} \leq C (\|\mathbf{f}\|_{V_2'} + \|\mathbf{g}\|_{-1/2, \Gamma_s}) (1 + |\mathbf{w}|_{1, \Omega}), \quad (66)$$

for some constant $C > 0$ independent of h .

Proof. We proceed by steps.

Step 1: Embedding of method (63) in abstract method.

Given an element $T \in \mathcal{C}_h$, let us define $\tau_K^{(p)} = \frac{|T|}{\int_T D \, d\mathbf{x}} \tau_T^{(p)}$, for any element $K \in \mathcal{T}_h$ that be in the prism P_T that lies on T . We assume that the pressures of N_h are defined on the whole Ω , as constant functions in the x_3

variable. Then,

$$\begin{aligned}
\sum_{K \in \mathcal{T}_h} \tau_K^{(p)} (\nabla r_h, \nabla q_h)_K &= \sum_{T \in \mathcal{C}_h} \frac{|T|}{\int_T D \, d\mathbf{x}} \tau_T^{(p)} \int_{P_T} \nabla_H r_h \cdot \nabla_H q_h \, d\mathbf{x} \, dx_3 \\
&= \sum_{T \in \mathcal{C}_h} \frac{|T|}{\int_T D \, d\mathbf{x}} \tau_T^{(p)} (\nabla_H r_h)|_T \cdot (\nabla_H q_h)|_T \int_{P_T} d\mathbf{x} \, dx_3 \\
&= \sum_{T \in \mathcal{C}_h} \tau_T^{(p)} (\nabla_H r_h, \nabla_H q_h)_T, \quad \forall r_h, q_h \in N_h.
\end{aligned} \tag{67}$$

Let us define $M_h = V_h/\mathbf{R}$, where V_h is given by (62). We now apply Lemma 1: There exists a bubble finite element space B_{1h} , generated on \mathcal{T}_h by a reference element $B_1^* \subset [H_0^1(K^*)]^3$, and a bilinear coercive form \mathbf{S}_{1h} on $[H_0^1(\Omega)]^3$, such that

$$\sum_{K \in \mathcal{T}_h} \tau_K^{(p)} (\nabla r_h, \nabla q_h)_K = \mathbf{S}_{1h}(\mathcal{R}_{1h}(\nabla r_h), \mathcal{R}_{1h}(\nabla q_h)), \quad \forall r_h, q_h \in M_h; \tag{68}$$

where \mathcal{R}_{1h} is the static condensation operator on B_{1h} with respect to form \mathbf{S}_{1h} . We may identify N_h with the subspace of M_h defined by $\{q_h \in V_h \mid \partial_3 q_h = 0\}$. Then, from (67) and (68) we deduce

$$\sum_{T \in \mathcal{C}_h} \tau_T^{(p)} (\nabla_H r_h, \nabla_H q_h)_T = \mathbf{S}_{1h}(\mathcal{R}_{1h}(\nabla r_h), \mathcal{R}_{1h}(\nabla q_h)), \quad \forall r_h, q_h \in N_h. \tag{69}$$

Also, again by Lemma 1, there exists a bubble finite element space B_{2h} , generated on \mathcal{T}_h by a reference element $B_2^* \subset [H_0^1(K^*)]^2$, and a bilinear coercive form \mathbf{S}_{2h} on $[H_0^1(\Omega)]^2$, such that $\forall \mathbf{u}_h, \mathbf{v}_h \in Y_h$,

$$\sum_{K \in \mathcal{T}_h} \tau_K^{(c)} (\mathbf{W}_h \cdot \nabla \mathbf{u}_h, \mathbf{W}_h \cdot \nabla \mathbf{v}_h)_K = \mathbf{S}_{2h}(\mathcal{R}_{2h}(\mathbf{W}_h \cdot \nabla \mathbf{u}_h), \mathcal{R}_{2h}(\mathbf{W}_h \cdot \nabla \mathbf{v}_h)); \tag{70}$$

where \mathcal{R}_{2h} is the static condensation operator on B_{2h} with respect to form \mathbf{S}_{2h} . Then,

$$\begin{aligned}
B_h^{(PE)}(\mathbf{u}_h, r_h; \mathbf{v}_h, q_h) &= B^{(PE)}(\mathbf{u}_h, r_h; \mathbf{v}_h, q_h) \\
&\quad + \mathbf{S}_{2h}(\mathcal{R}_{2h}(\mathbf{W}_h \cdot \nabla \mathbf{u}_h), \mathcal{R}_{2h}(\mathbf{W}_h \cdot \nabla \mathbf{v}_h)) \\
&\quad - \mathbf{S}_{1h}(\mathcal{R}_{1h}(\nabla r_h), \mathcal{R}_{1h}(\nabla q_h)), \quad \forall \mathbf{u}_h, \mathbf{v}_h \in Y_h, \forall r_h, q_h \in N_h.
\end{aligned}$$

We recall that by Theorem (2) (Step 3), the forms $\{\mathbf{S}_{2h}\}_{h>0}$ are uniformly continuous and coercive in H^1 norm. Also, due to (64) and the regularity of triangulations \mathcal{T}_h , the coefficients $\tau_K^{(p)}$ are of order h_K^2 . Then, the forms $\{\mathbf{S}_{1h}\}_{h>0}$ also are uniformly continuous and coercive.

Step 2: Discrete inf-sup condition.

We state the following:

Lemma 6. *Given $\alpha \in (1, 2]$, there exists a constant $C_\alpha > 0$ such that $\forall q_h \in N_h$,*

$$C_\alpha \|q_h\|_{L_{D,0}^\alpha(\omega)} \leq \sup_{\mathbf{v}_h \in Y_h - \{0\}} \frac{(\nabla_H \cdot \langle \mathbf{v}_h \rangle, q_h)_\omega}{|\mathbf{v}_h|_{1,\alpha',\Omega}} + [\mathbf{S}_{1h}(\mathcal{R}_{1h}(\nabla q_h), \mathcal{R}_{1h}(\nabla q_h))]^{1/2}, \tag{71}$$

where α' is the conjugate exponent of α .

Proof. Define the space $W_h = \{(\mathbf{v}_h, v_{3h}) \in V_h^3 \mid (\mathbf{v}_h, v_{3h})|_{\partial\Omega} = 0\}$. It is enough to prove that

$$C_\alpha \|q_h\|_{L_0^\alpha(\Omega)} \leq \sup_{(\mathbf{v}_h, v_{3h}) \in W_h - \{0\}} \frac{(\nabla \cdot (\mathbf{v}_h, v_{3h}), q_h)_\Omega}{|(\mathbf{v}_h, v_{3h})|_{1, \alpha', \Omega}} + [\mathbf{S}_{1h}(\mathcal{R}_{1h}(\nabla q_h), \mathcal{R}_{1h}(\nabla q_h))]^{1/2}, \quad \forall q_h \in V_h. \quad (72)$$

Indeed, if $q_h \in N_h$, $(\mathbf{v}_h, v_{3h}) \in W_h$,

$$(\nabla \cdot (\mathbf{v}_h, v_{3h}), q_h)_\Omega = (\nabla_H \cdot \mathbf{v}_h, q_h)_\Omega - (v_{3h}, \partial_3 q_h)_\Omega = (\nabla_H \cdot \langle \mathbf{v}_h \rangle, q_h)_\omega.$$

Then,

$$\begin{aligned} \sup_{(\mathbf{v}_h, v_{3h}) \in W_h - \{0\}} \frac{(\nabla \cdot (\mathbf{v}_h, v_{3h}), q_h)_\Omega}{|(\mathbf{v}_h, v_{3h})|_{1, \alpha', \Omega}} &= \sup_{(\mathbf{v}_h, v_{3h}) \in W_h - \{0\}} \frac{(\nabla_H \cdot \langle \mathbf{v}_h \rangle, q_h)_\omega}{|(\mathbf{v}_h, v_{3h})|_{1, \alpha', \Omega}} \\ &\leq \sup_{\mathbf{v}_h \in Y_h - \{0\}} \frac{(\nabla_H \cdot \langle \mathbf{v}_h \rangle, q_h)_\omega}{|\mathbf{v}_h|_{1, \alpha', \Omega}}. \end{aligned}$$

Also, $\|q_h\|_{L_0^\alpha(\Omega)} = \|q_h\|_{L_{D,0}^\alpha(\omega)}$ if $q_h \in N_h$. Thus, (71) follows from (72).

To prove (72), consider $q_h \in V_h$. As Ω is polyhedric, then $\partial\Omega$ is Lipschitz, and the continuous inf-sup condition in $L^\alpha(\Omega)$ norm is satisfied (*cf.* Amrouche and Girault [1]): There exists a constant $D_\alpha > 0$ such that

$$D_\alpha \|q_h\|_{L_0^\alpha(\Omega)} \leq \sup_{\mathbf{v} \in [W_0^{1, \alpha'}(\Omega)]^3 - \{0\}} \frac{(\nabla \cdot \mathbf{v}, q_h)_\Omega}{|\mathbf{v}|_{1, \alpha', \Omega}}, \quad \forall q \in L_0^\alpha(\Omega).$$

As $[\mathcal{D}(\Omega)]^3$ is dense in $[W_0^{1, \alpha'}(\Omega)]^3$, there exists a $\mathbf{v}_0 \in [\mathcal{D}(\Omega)]^3$ such that

$$\frac{1}{2} D_\alpha \|q_h\|_{L_0^\alpha(\Omega)} \leq (\nabla \cdot \mathbf{v}_0, q_h)_\Omega, \quad |\mathbf{v}_0|_{1, \alpha', \Omega} = 1.$$

Following the standard finite elements interpolation theory (*cf.* Ciarlet [13]), there exists an interpolate $\mathbf{v}_{0h} \in W_h$ such that

$$|\mathbf{v}_{0h}|_{1, \alpha', \Omega} \leq C_1 |\mathbf{v}_0|_{1, \alpha', \Omega}; \quad (73)$$

$$\|\mathbf{v}_{0h} - \mathbf{v}_0\|_{0, K} \leq C_1 h_K |\mathbf{v}_0|_{1, K}, \quad \forall K \in \mathcal{T}_h; \quad (74)$$

for some constant $C_1 > 0$ independent of h . Then, as q_h is continuous,

$$\begin{aligned} \frac{1}{2} D_\alpha \|q_h\|_{L_0^\alpha(\Omega)} &\leq (\nabla \cdot \mathbf{v}_{0h}, q_h)_\Omega + (\mathbf{v}_{0h} - \mathbf{v}_0, \nabla q_h)_\Omega \\ &\leq C_1 \frac{|(\nabla \cdot \mathbf{v}_{0h}, q_h)_\Omega|}{|\mathbf{v}_{0h}|_{1, \alpha', \Omega}} \\ &\quad + \left[\sum_{K \in \mathcal{T}_h} h_K^{-2} \|\mathbf{v}_{0h} - \mathbf{v}_0\|_{0, K}^2 \right]^{1/2} \left[\sum_{K \in \mathcal{T}_h} h_K^2 \|\nabla q_h\|_{0, K}^2 \right]^{1/2}. \end{aligned}$$

As $\alpha' \geq 2$, then (74) yields

$$\left[\sum_{K \in \mathcal{T}_h} h_K^{-2} \|\mathbf{v}_{0h} - \mathbf{v}_0\|_{0, K}^2 \right]^{1/2} \leq C_2 |\mathbf{v}_0|_{1, \alpha', \Omega}.$$

Also, from the representation formula (68), hypothesis (64) and the regularity of the triangulations, we obtain

$$\sum_{K \in \mathcal{T}_h} h_K^2 \|\nabla q_h\|_{0,K}^2 \leq C_3 \mathbf{S}_{1h}(\mathcal{R}_{1h}(\nabla q_h), \mathcal{R}_{1h}(\nabla q_h)).$$

Thus, estimate (72) follows.

Step 3: Existence of solution of discrete problem.

Problem (63) is equivalent to a square linear system of dimension $\dim(Y_h) + \dim(N_h)$. Then, the existence of solution follows from its uniqueness. If we prove that any solution is bounded by a norm of the data, the uniqueness follows. Let us then consider a solution $(\mathbf{y}_h, p_h) \in Y_h \times N_h$ of problem (63).

Velocity estimate. Denote $\mathbf{c}_h = \mathcal{R}_{1h}(\nabla p_h) \in B_{1h}$, $\mathbf{d}_h = \mathcal{R}_{2h}(\mathbf{W}_h \cdot \nabla \mathbf{y}_h) \in B_{2h}$. As $\nabla \cdot \mathbf{W} = 0$, $w_3|_{\Gamma_s} = 0$, $\mathbf{y}_h \in [W^{1,\infty}(\Omega)]^2$, using (57) we have

$$\langle \mathbf{W} \cdot \nabla \mathbf{y}, \mathbf{y} \rangle_{V'_4 - V_4} = \frac{1}{2} \int_{\partial\Omega} \mathbf{W} \cdot \mathbf{n} |\mathbf{y}|^2 d(\partial\Omega) - \frac{1}{2} \int_{\Omega} \nabla \cdot \mathbf{W} |\mathbf{y}|^2 dx dx_3 = 0.$$

Then,

$$\nu |\mathbf{y}_h|_{1,\Omega}^2 + \mathbf{S}_{1h}(\mathbf{c}_h, \mathbf{c}_h) + \mathbf{S}_{2h}(\mathbf{d}_h, \mathbf{d}_h) = B_h^{(PE)}(\mathbf{y}_h, p_h; \mathbf{y}_h, -p_h) = F(\mathbf{y}_h) \leq \|F\|_{V'_2} |\mathbf{y}_h|_{1,\Omega},$$

where we assume V_2 to be endowed with the $|\cdot|_{1,\Omega}$ norm. Then,

$$|\mathbf{y}_h|_{1,\Omega} \leq \nu^{-1} \|F\|_{V'_2}, \quad \mathbf{S}_{1h}(\mathbf{c}_h, \mathbf{c}_h) \leq \nu^{-1} \|F\|_{V'_2}^2, \quad |\mathbf{d}_h|_{1,\Omega} \leq (\nu\nu_2)^{-1/2} \|F\|_{V'_2}, \quad (75)$$

where ν_2 is the uniform coerciveness constant of the forms \mathbf{S}_{2h} .

Pressure estimate. Take $\mathbf{v}_h \in Y_h$. Then,

$$\begin{aligned} (\nabla_H \cdot \langle \mathbf{v}_h \rangle, p_h)_\omega &= -F(\mathbf{v}_h) + \langle \mathbf{W} \cdot \nabla \mathbf{y}_h, \mathbf{v}_h \rangle_{V'_4 - V_4} + \nu (\nabla \mathbf{y}_h, \nabla \mathbf{v}_h)_\Omega \\ &\quad + \mathbf{S}_{2h}(\mathbf{d}_h, \mathcal{R}_{2h}(\mathbf{W}_h \cdot \nabla \mathbf{v}_h)). \end{aligned}$$

Observe that, due to Sobolev's injections and the stability interpolate (65),

$$\begin{aligned} \mathbf{S}_{2h}(\mathbf{d}_h, \mathcal{R}_{2h}(\mathbf{W}_h \cdot \nabla \mathbf{v}_h)) &= (\mathbf{W}_h \cdot \nabla \mathbf{v}_h, \mathbf{d}_h)_\Omega \leq C_1 \|\mathbf{w}_h\|_{0,4;\Omega} |\mathbf{v}_h|_{1,\Omega} \|\mathbf{d}_h\|_{0,4;\Omega} \\ &\quad + C_2 \|w_3\|_{0,\Omega} \|\partial_3 \mathbf{v}_h\|_{0,3;\Omega} \|\mathbf{d}_h\|_{0,6;\Omega} \\ &\leq C_3 |\mathbf{w}|_{1,\Omega} |\mathbf{v}_h|_{1,3;\Omega} \|\mathbf{d}_h\|_{0,6;\Omega} \end{aligned}$$

Then, due to (55),

$$\begin{aligned} (\nabla_H \cdot \langle \mathbf{v}_h \rangle, p_h)_\omega &\leq C_4 (\|F\|_{V'_2} + |\mathbf{w}|_{1,\Omega} |\mathbf{y}_h|_{1,\Omega} + \nu |\mathbf{y}_h|_{1,\Omega} \\ &\quad + |\mathbf{w}|_{1,\Omega} |\mathbf{d}_h|_{1,\Omega}) |\mathbf{v}_h|_{1,3;\Omega}. \end{aligned}$$

Then, using (75),

$$\sup_{\mathbf{v}_h \in Y_h - \{0\}} \frac{(\nabla_H \cdot \langle \mathbf{v}_h \rangle, q_h)_\omega}{|\mathbf{v}_h|_{1,3;\Omega}} + [\mathbf{S}_{1h}(\mathbf{c}_h, \mathbf{c}_h)]^{1/2} \leq C_5 \|F\|_{V'_2} (1 + |\mathbf{w}|_{1,\Omega}).$$

Due to Lemma 6, we deduce that

$$\|p_h\|_{L_{D,0}^{3/2}(\omega)} \leq C_6 \|F\|_{V'_2} (1 + |\mathbf{w}|_{1,\Omega}). \quad (76)$$

Thus, the discrete problem (54) admits a unique solution satisfying

$$|\mathbf{y}_h|_1 + \|p_h\|_{L_{D,0}^{3/2}(\omega)} \leq C_7 \|F\|_{V_2'} (1 + |\mathbf{w}|_{1,\Omega}). \quad (77)$$

Step 4: Conclusion.

Due to estimates (77), the sequence $\{(\mathbf{y}_h, p_h)\}_{h>0}$ is bounded in $V_2 \times L_{D,0}^{3/2}(\omega)$, which is a reflexive space. Then, it contains a subsequence, that we still denote in the same way, weakly convergent in that space to a pair (\mathbf{y}, p) . Let us prove that this pair is a solution of problem (50).

Consider a pair $(\mathbf{v}, q) \in V_4 \times L_{D,0}^2(\omega)$. There exists a sequence $\{(\mathbf{v}_h, q_h)\}_{h>0}$ with $(\mathbf{v}_h, q_h) \in Y_h \times N_h$ which is strongly convergent to (\mathbf{v}, q) in $V_4 \times L_{D,0}^2(\omega)$. Indeed, the fact that $\mathbf{v}_h \rightarrow \mathbf{v}$ in V_4 is proved by the standard interpolation estimates by piecewise affine finite elements. Also, by the same theory, there exists a sequence $\{q_h\}_{h>0}$ with $q_h \in \tilde{N}_h$ which converges to q in $L^2(\omega)$. But

$$\|q_h - q\|_{L_{D,0}^2(\omega)} \leq \|q_h - q\|_{L_D^2(\omega)} \leq \|D\|_{0,\infty;\omega}^{1/2} \|q_h - q\|_{0,2;\omega}.$$

Thus, $\lim_{h \rightarrow 0} \|q_h - q\|_{L_{D,0}^2(\omega)} = 0$.

From (57) we deduce

$$\lim_{h \rightarrow 0} \langle \mathbf{W} \cdot \nabla \mathbf{y}_h, \mathbf{v}_h \rangle_{V_4' - V_4} = \langle \mathbf{W} \cdot \nabla \mathbf{y}, \mathbf{v} \rangle_{V_4' - V_4}.$$

Also,

$$\lim_{h \rightarrow 0} (p_h, \nabla_H \cdot \langle \mathbf{v}_h \rangle)_\omega = \lim_{h \rightarrow 0} (p_h, \nabla_H \cdot \mathbf{v}_h)_\Omega = (p, \nabla_H \cdot \mathbf{v})_\Omega = (p, \nabla_H \cdot \langle \mathbf{v} \rangle)_\omega,$$

and similarly

$$\lim_{h \rightarrow 0} (\nabla_H \cdot \langle \mathbf{y}_h \rangle, q_h)_\omega = (\nabla_H \cdot \langle \mathbf{y} \rangle, q)_\omega.$$

In a standard way, we have

$$\lim_{h \rightarrow 0} (\nabla \mathbf{y}_h, \nabla \mathbf{v}_h)_\Omega = (\nabla \mathbf{y}, \nabla \mathbf{v})_\Omega.$$

To pass to the limit in the stabilizing terms, we need the following property of the bubble finite element spaces:

Lemma 7. *Consider a family $\{Z_h\}_{h>0}$ of finite element subspaces of $[H_0^1(\Omega)]^d$ generated by a reference space Z^* . Assume that Z^* does not contain the constant functions. Then, the following statements hold.*

i) *For any $q \in [2, 6]$, there exists a constant $C_q > 0$ such that*

$$\|\mathbf{z}_h\|_{0,q} \leq C_q h^\beta |\mathbf{z}_h|_1, \quad \forall \mathbf{z}_h \in Z_h; \quad \text{where } \frac{\beta}{2} + \frac{1-\beta}{6} = \frac{1}{q}. \quad (78)$$

ii) *If a sequence $\{\mathbf{z}_h\}_{h>0}$, with $\mathbf{z}_h \in Z_h$, $\forall h > 0$, is bounded in $[H_0^1(\Omega)]^d$, then it converges weakly to zero in $[H_0^1(\Omega)]^d$.*

This Lemma is proved in the Appendix.

Due to (75) and to the uniform coerciveness of the forms \mathbf{S}_{1h} , the sequence $\{\mathbf{c}_h\}_{h>0}$ is bounded in $[H_0^1(\Omega)]^3$. Then, it is weakly convergent to zero in that space. Thus,

$$\lim_{h \rightarrow 0} \mathbf{S}_{1h}(\mathcal{R}_{1h}(\nabla q_h), \mathbf{c}_h) = \lim_{h \rightarrow 0} (-q_h, \nabla \cdot \mathbf{c}_h)_\Omega = 0.$$

Also

$$\begin{aligned}
|\mathbf{S}_{2h}(\mathbf{d}_h, \mathcal{R}_{2h}(\mathbf{W}_h \cdot \nabla \mathbf{v}_h))| &= |\langle \mathbf{W}_h \cdot \nabla \mathbf{v}_h, \mathbf{d}_h \rangle_{H^{-1}-H_0^1}| = \left| \int_{\Omega} (\mathbf{W}_h \cdot \nabla \mathbf{v}_h) \cdot \mathbf{d}_h \, d\mathbf{x} \, dx_3 \right| \\
&\leq C_7 (\|\mathbf{w}_h\|_{0,4;\Omega} \|\mathbf{v}_h\|_{1,\Omega} \|\mathbf{d}_h\|_{0,4;\Omega} + \|w_{3h}\|_{0,\Omega} \|\partial_3 \mathbf{v}_h\|_{0,4;\Omega} \|\mathbf{d}_h\|_{0,4;\Omega}) \\
&\leq C_8 |\mathbf{w}|_{1,\Omega} |\mathbf{v}_h|_{1,4;\Omega} \|\mathbf{d}_h\|_{0,4;\Omega} \\
&\leq C_9 h^{1/4} |\mathbf{w}|_{1,\Omega} |\mathbf{v}_h|_{1,4;\Omega} \|\mathbf{d}_h\|_{1,\Omega} \\
&\leq C_{10} h^{1/4} |\mathbf{w}|_{1,\Omega} |\mathbf{v}_h|_{1,4;\Omega} \|F\|_{V_2'}.
\end{aligned}$$

Thus,

$$\lim_{h \rightarrow 0} \mathbf{S}_{2h}(\mathbf{d}_h, \mathcal{R}_{2h}(\mathbf{W}_h \cdot \nabla \mathbf{v}_h)) = 0.$$

To complete the proof, we combine the weak lower semicontinuity of the norm on reflexive Banach spaces with estimate (77) to derive estimate (66). \square

Remark 8. This result shows the adaptivity of the general formulation provided by the abstract method (10), and of its analysis technique in Sections 3 and 4. The two main contributions in the actual application are

- To derive a discrete inf-sup condition in L^α norm ($1 < \alpha < 2$), and
- To prove that the stabilizing terms vanish in the limit $h \rightarrow 0$.

We have chosen piecewise affine elements for simplicity, but the same analysis applies to the general finite element spaces introduced in Section 2.

7. CONCLUSION

We have developed in this paper a systematic way to extend the standard stability analysis of mixed methods to stabilized methods. The stability of pressure discretization follows from an underlying discrete inf-sup condition. The stability of velocity discretization follows from a uniform separation property between standard finite element spaces and bubble finite element spaces.

We have proved the adaptivity of this technique by analyzing two non-standard situations (spectral element method and primitive equations of the ocean) by means of the same essential analysis.

Roughly speaking, we have found a way to extend to stabilized methods any stability-related property that one could prove for stable mixed methods, as there is always an underlying discrete inf-sup condition to each actual stabilized method.

APPENDIX

Proof of Lemma 1. This proof is based upon the following:

Lemma 8. Denote by $B_j(K)$, $W_j(K)$, respectively, the affine-transformed of spaces B_j^* , W_j^* on element K .

Then, $B_j(K)$ is a space of virtual bubbles of $[H_0^1(K)]^d$ reproducing the operator $\tau_K I_K$ on $W_j(K)$, with respect to the inner product \mathbf{S}_K .

Proof. We shall respectively denote by $(\cdot, \cdot)_*$ and $((\cdot, \cdot))_*$ the standard inner products in $[L^2(K^*)]^d$ and $[H_0^1(K^*)]^d$. Consider an element $K \in \mathcal{T}_h$. Given a function \mathbf{v}^* defined on the reference element K^* , let us denote by $\hat{\mathbf{v}}$ its affine-transformed on K , $\hat{\mathbf{v}} = \mathbf{v}^* \circ (F_K)^{-1}$, where F_K is the affine mapping that transforms K^* on to K , i.e., $F_K(x^*) = A_K x^* + b_K$.

If $\mathbf{w}^*, \mathbf{v}^* \in [H_0^1(K^*)]^d$, then $\hat{\mathbf{w}}, \hat{\mathbf{v}} \in [H_0^1(K)]^d$ and

$$\mathbf{S}_K(\hat{\mathbf{w}}, \hat{\mathbf{v}})_K = |\det A_K| \beta_K h_K^{-2} ((\mathbf{w}^*, \mathbf{v}^*))_* \tag{79}$$

Denote by \mathcal{R}^* the static condensation operator acting on space B_j^* , associated to the inner product $((\cdot, \cdot))_*$. Given $\mathbf{v}^* \in W_j^*$, $\mathcal{R}^*(\mathbf{v}^*)$ satisfies

$$((\mathcal{R}^*(\mathbf{v}^*), \zeta^*))_* = (\mathbf{v}^*, \zeta^*)_*, \quad \forall \zeta^* \in B_j^*.$$

From (79) we deduce that for any $\mathbf{w} \in W_j(K)$,

$$h_K^2 \beta_K^{-1} \mathbf{S}_K(\widehat{\mathcal{R}^*(\mathbf{w}^*)}, \zeta) = (\mathbf{w}, \zeta)_K, \quad \forall \zeta \in B_j(K);$$

where $\mathbf{w}^* = \mathbf{w} \circ F_K$. Then, $\mathcal{R}_K(\mathbf{w}) = h_K^2 \beta_K^{-1} \widehat{\mathcal{R}^*(\mathbf{w}^*)}$.

Furthermore, by hypothesis,

$$((\mathcal{R}^*(\mathbf{w}_1^*), \mathcal{R}^*(\mathbf{w}_2^*)))_* = \tau^*(\mathbf{w}_1^*, \mathbf{w}_2^*)_*, \quad \forall \mathbf{w}_1^*, \mathbf{w}_2^* \in W_j^*.$$

From (79), this implies

$$h_K^2 \beta_K^{-1} \mathbf{S}_K(\widehat{\mathcal{R}^*(\mathbf{w}_1^*)}, \widehat{\mathcal{R}^*(\mathbf{w}_2^*)}) = \tau^*(\mathbf{w}_1, \mathbf{w}_2)_K, \quad \forall \mathbf{w}_1, \mathbf{w}_2 \in W_j(K).$$

Thus,

$$\mathbf{S}_K(\mathcal{R}_K(\mathbf{w}_1), \mathcal{R}_K(\mathbf{w}_2)) = \tau_K(\mathbf{w}_1, \mathbf{w}_2)_K, \quad \forall \mathbf{w}_1, \mathbf{w}_2 \in W_j(K).$$

□

Consider now an element $\varphi \in [H^{-1}(\Omega)]^d$. Then, $\mathcal{R}_h(\varphi)|_K = \mathcal{R}_K(\varphi|_K)$. This occurs because $\text{int}(K) \cap \text{int}(K') = \emptyset$ if $K \neq K'$.

Then, if $\mathbf{w}_1, \mathbf{w}_2 \in W_h^{(j)}$,

$$\mathbf{S}_h(\mathcal{R}_h^{(j)}(\mathbf{w}_1), \mathcal{R}_h^{(j)}(\mathbf{w}_2)) = \sum_{K \in \mathcal{T}_h} \mathbf{S}_K(\mathcal{R}_h^{(j)}(\mathbf{w}_1|_K), \mathcal{R}_h^{(j)}(\mathbf{w}_2|_K)) = \sum_{K \in \mathcal{T}_h} \tau_K(\mathbf{w}_2, \mathbf{w}_1)_K.$$

This last equality holds because of Lemma 8. To finish the proof, we observe that the orthogonal projection of $\mathbf{f}|_K$ on $W(K)$ with respect to the L^2 inner product is just $\mathbf{f}_h|_K$. Then,

$$\sum_{K \in \mathcal{T}_h} \tau_K(\mathbf{f}, \mathbf{w}_1)_K = \sum_{K \in \mathcal{T}_h} \tau_K(\mathbf{f}_h, \mathbf{w}_1)_K = \mathbf{S}_h(\mathcal{R}_h^{(j)}(\mathbf{w}_1), \mathcal{R}_h^{(j)}(\mathbf{f}_h)).$$

□

Proof of Lemma 2. Consider a triangulation \mathcal{T}_h of the family. Given an element $K \in \mathcal{T}_h$, denote by h_K the diameter of element K . Let us consider the following bilinear form on $[H_0^1(\Omega)]^d$,

$$((\mathbf{w}, \mathbf{v}))_h = \sum_{K \in \mathcal{T}_h} \int_K (C_K \nabla \mathbf{w}) : \nabla \mathbf{v} \, dx, \quad \forall \mathbf{w}, \mathbf{v} \in [H_0^1(\Omega)]^d.$$

Here, matrix C_K is defined as in (34). The form $((\cdot, \cdot))_h$ is an inner product on $[H_0^1(\Omega)]^d$, as each matrix C_K is symmetric and positive definite.

Some standard estimates yield

$$|((\mathbf{w}, \mathbf{v}))_h| \leq M_h |\mathbf{w}|_1 |\mathbf{v}|_1, \quad ((\mathbf{w}, \mathbf{w}))_h \geq \Lambda_h |\mathbf{w}|_1^2,$$

where $M_h = \frac{1}{h_K} \max_{K,l} \sigma_l^K$, $\Lambda_h = \frac{1}{h_K} \min_{K,l} \sigma_l^K$, $\sigma_1^K, \dots, \sigma_d^K$ denoting the singular values of A_K .

As the family of triangulations is regular, there exist two constants $C_1 > 0$, $C_2 > 0$ such that (cf. Ciarlet [13]),

$$C_1 h_K \leq \sigma_l^K \leq C_2 h_K, \quad \forall K \in \mathcal{T}_h, \forall h > 0, \forall l = 1, \dots, d. \quad (80)$$

Consequently, the forms $((\cdot, \cdot))_h$ define inner products on $[H_0^1(\Omega)]^d$ which are uniformly (in h) equivalent to the standard one. If we denote by $\|\cdot\|_h$ the norm generated by the inner product $((\cdot, \cdot))_h$, we have

$$\Lambda |\mathbf{w}|_1 \leq \|\mathbf{w}\|_h \leq M |\mathbf{w}|_1, \quad \forall \mathbf{w} \in [H_0^1(\Omega)]^d; \quad \text{where } \Lambda = C_1^2 > 0, \quad M = C_2^2 > 0. \quad (81)$$

Let us now make the change of variable $x = A_K x^* + b_K$. Given $\mathbf{w}, \mathbf{v} \in [H_0^1(\Omega)]^d$, define $\mathbf{w}^K(x^*) = \mathbf{w}(x)$, $\mathbf{v}^K(x^*) = \mathbf{v}(x)$, $\forall x^* \in K^*$. Then,

$$\int_K (C_K \nabla \mathbf{w}) : \nabla \mathbf{v} \, dx = h_K^{-2} |\det A_K| \int_{K^*} \nabla \mathbf{w}^K : \nabla \mathbf{v}^K \, dx^*.$$

Thus, given $\mathbf{y}_h \in Y_h$, $\mathbf{z}_h \in Z_h$, we have

$$((\mathbf{y}_h, \mathbf{z}_h))_h = \sum_{K \in \mathcal{T}_h} h_K^{-2} |\det A_K| \int_{K^*} \nabla \mathbf{y}_h^K : \nabla \mathbf{z}_h^K \, dx^*. \quad (82)$$

Notice that we always have $\mathbf{y}_h^K \in Y^*$, $\mathbf{z}_h^K \in Z^*$.

Consider now the bilinear form on $[H^1(K^*)]^d$,

$$((\tilde{\mathbf{w}}, \tilde{\mathbf{v}}))_* = \int_{K^*} \nabla \tilde{\mathbf{w}} : \nabla \tilde{\mathbf{v}} \, dx^*.$$

This is an inner product on the quotient space $\tilde{H} = [H^1(K^*)]^d / \mathbf{R}$. The spaces $\tilde{Y}^* = Y^* / \mathbf{R}$ and $\tilde{Z}^* = Z^* / \mathbf{R}$ are subspaces of \tilde{H} satisfying $\tilde{Y}^* \cap \tilde{Z}^* = \{0\}$.

Consider now a fixed nonzero $\tilde{\mathbf{z}} \in \tilde{Z}^*$. Denote by Π the orthogonal projection from \tilde{H} on to \tilde{Y}^* with respect to the inner product $((\cdot, \cdot))_*$. Then,

$$-\|\tilde{\mathbf{y}}\|_* \|\Pi \tilde{\mathbf{z}}\|_* \leq ((\tilde{\mathbf{y}}, \tilde{\mathbf{z}}))_* \leq \|\tilde{\mathbf{y}}\|_* \|\Pi \tilde{\mathbf{z}}\|_*, \quad \forall \tilde{\mathbf{y}} \in \tilde{Y}^*;$$

where $\|\cdot\|_*$ denotes the norm associated to the inner product $((\cdot, \cdot))_*$. As $\tilde{Y}^* \cap \tilde{Z}^* = \{0\}$, $\tilde{\mathbf{z}}$ cannot belong to \tilde{Y}^* and $\|\Pi \tilde{\mathbf{z}}\|_* < \|\tilde{\mathbf{z}}\|_*$. Thus, there exists a constant $\delta > 0$ such that

$$|((\tilde{\mathbf{y}}, \tilde{\mathbf{z}}))_*| \leq (1 - \delta) \|\tilde{\mathbf{y}}\|_* \|\tilde{\mathbf{z}}\|_*, \quad \forall \tilde{\mathbf{y}} \in \tilde{Y}^*. \quad (83)$$

Define $N = \sup \{|((\tilde{\mathbf{y}}, \tilde{\mathbf{z}}))_*| \text{ for } \tilde{\mathbf{y}} \in \tilde{Y}^*, \tilde{\mathbf{z}} \in \tilde{Z}^* \text{ with } \|\tilde{\mathbf{y}}\|_* = \|\tilde{\mathbf{z}}\|_* = 1\}$. As both \tilde{Y}^* and \tilde{Z}^* are spaces of finite dimension, in fact this supremum is achieved. Due to (83), we should have $N < 1$. Thus, we may assume that (83) holds for all $\tilde{\mathbf{y}} \in \tilde{Y}^*$ and for all $\tilde{\mathbf{z}} \in \tilde{Z}^*$.

From (82) we now obtain

$$\begin{aligned} |((\mathbf{y}_h, \mathbf{z}_h))_h| &\leq (1 - \delta) \sum_{K \in \mathcal{T}_h} h_K^{-2} |\det A_K| \left(\int_{K^*} |\nabla \mathbf{y}_h^K|^2 \, dx^* \right)^{1/2} \left(\int_{K^*} |\nabla \mathbf{z}_h^K|^2 \, dx^* \right)^{1/2} \\ &\leq (1 - \delta) \|\mathbf{y}_h\|_h \|\mathbf{z}_h\|_h. \end{aligned}$$

Then,

$$\begin{aligned} \|\mathbf{y}_h + \mathbf{z}_h\|_h^2 &\geq \|\mathbf{y}_h\|_h^2 + \|\mathbf{z}_h\|_h^2 - 2(1 - \delta) \|\mathbf{y}_h\|_h \|\mathbf{z}_h\|_h \\ &\geq \delta (\|\mathbf{y}_h\|_h^2 + \|\mathbf{z}_h\|_h^2). \end{aligned}$$

From this last inequality,

$$\|\mathbf{y}_h\|_h + \|\mathbf{z}_h\|_h \leq \sqrt{\frac{2}{\delta}} \|\mathbf{y}_h + \mathbf{z}_h\|_h.$$

Now, it is enough to use the uniform coerciveness and boundedness of the inner products $((\cdot, \cdot))_h$ to conclude the proof. \square

Proof of Lemma 4. The elements of $[Q_N(K^*)]^d$ are elements of $[H^{-1}(K^*)]^d$ if we identify the $H_0^1 - H^{-1}$ duality with the L^2 inner product. Observe that the bilinear form $(\cdot, \cdot)_{N, K^*}$ is an inner product on $[Q_N(K^*)]^d$. Indeed, if $(\hat{q}_N, \hat{q}_N)_{N, K^*} = 0$ for some $\hat{q}_N \in Q_N(K^*)$, then $\hat{q}_N(\xi_i, \xi_j, \xi_k) = 0$, for $i, j, k = 1, \dots, N+1$. Then, $\hat{q}_N = 0$.

Thus, similarly to the Step 1 in the proof of Theorem 2, by Theorem 4, there exists a $\hat{\mu} > 0$ such that if $0 < \tau^* \leq \hat{\mu}$, there exists a finite-dimensional subspace B^* of $[H_0^1(K^*)]^d$ that reproduces the operator $\tau^* I$ on $([Q_N(K^*)]^d, (\cdot, \cdot)_{N, K^*})$ with respect to the standard inner product of $[H_0^1(K^*)]^d$.

Consider now an element $K \in \mathcal{T}_h$, and two functions $\mathbf{v}, \mathbf{w} \in [C^0(\Omega)]^d$. Define $\hat{\mathbf{v}} = \mathbf{v} \circ F_K$, $\hat{\mathbf{w}} = \mathbf{w} \circ F_K$, where F_K is a bijective affine transformation from K^* on to K . Then,

$$(\mathbf{v}, \mathbf{w})_{N, K} = |\det A_K| (\hat{\mathbf{v}}, \hat{\mathbf{w}})_{N, K^*}, \quad A_K = \nabla F_K.$$

Denote by $B_j(K)$ the affine-transformed of B^* by F_K^{-1} : $B_j(K) = B^* \circ F_K^{-1}$. Similarly to Lemma 8, we deduce that $B_j(K)$ is a space of virtual bubbles that reproduces the operator $\tau_K I$ on $([Q_N(K)]^d, (\cdot, \cdot)_{N, K})$ with respect to the inner product \mathbf{S}_K defined by (34) with $\beta_K = \tau^* \tau_K^{-1} h_K^2$.

Denote also by $B_{\mathcal{H}}$ the finite element subspace of $[H_0^1(\Omega)]^d$ generated by the reference element B^* on triangulation \mathcal{T}_h . Similarly to Lemma 1, we deduce that

$$\mathbf{S}_h(\mathcal{R}_{\mathcal{H}}(\mathbf{w}_1), \mathcal{R}_{\mathcal{H}}(\mathbf{w}_2)) = \sum_{K \in \mathcal{T}_h} \tau_K (\mathbf{w}_1, \mathbf{w}_2)_{N, K}, \quad \forall \mathbf{w}_1, \mathbf{w}_2 \in W_{\mathcal{H}},$$

where $\mathcal{R}_{\mathcal{H}}$ denotes the static condensation operator on $B_{\mathcal{H}}$ with respect to form \mathbf{S}_h .

Consider now $\mathbf{v}_1, \mathbf{v}_2 \in [C_{p, h}(\Omega)]^d$. Then, $I_{\mathcal{H}} \mathbf{v}_i \in W_{\mathcal{H}}$, $i = 1, 2$ and $(I_{\mathcal{H}} \mathbf{v}_1, I_{\mathcal{H}} \mathbf{v}_2)_{N, K} = (\mathbf{v}_1, \mathbf{v}_2)_{N, K}$. Consequently, (45) holds. \square

Proof of Lemma 7. i) As Z^* does not contain the constant functions, then the H^1 seminorm is a norm on Z^* . Therefore, there exists a constant $C_* > 0$ such that

$$\|\mathbf{z}^*\|_0 \leq C_* |\mathbf{z}^*|_1, \quad \forall \mathbf{z}^* \in Z^*.$$

Consider now $\mathbf{z}_h \in Z_h$. Then, using the notation introduced in Lemma 2,

$$\|\mathbf{z}_h\|_0^2 = \sum_{K \in \mathcal{T}_h} |\det A_K| \|\mathbf{z}_h^K\|_{0, K^*}^2 \leq C_* \sum_{K \in \mathcal{T}_h} |\det A_K| |\mathbf{z}_h^K|_{1, K^*}^2.$$

Following Girault and Raviart [17], Lemma A.1, there exists a constant $\gamma > 0$ such that

$$|\det A_K|^{1/2} |\mathbf{z}_h^K|_{1, K^*} \leq \gamma \|A_K\| |\mathbf{z}_h|_{1, K},$$

where $\|A_K\|$ denotes the spectral matrix norm. Then,

$$\|\mathbf{z}_h\|_0^2 \leq C_* \gamma^2 \sum_{K \in \mathcal{T}_h} \|A_K\|^2 |\mathbf{z}_h|_{1,K}^2 \leq C_1 h^2 |\mathbf{z}_h|_1^2,$$

for some constant $C_1 > 0$.

Let us now consider the interpolation estimate (*cf.* Brézis [6]),

$$\|\mathbf{w}\|_{0,q} \leq C_2 \|\mathbf{w}\|_0^\beta \|\mathbf{w}\|_{0,6}^{1-\beta}, \quad \forall \mathbf{w} \in [L^6(\Omega)]^d,$$

if $2 \leq q \leq 6$, with β given in (78). As $[H_0^1(\Omega)]^d$ is continuously embedded in $[L^6(\Omega)]^d$ if $d = 2$ or $d = 3$, then (78) follows.

ii) Consider a sequence $\{\mathbf{z}_h\}_{h>0}$, with $\mathbf{z}_h \in Z_h$. This sequence contains a subsequence, that we still denote in the same way, weakly convergent to some element \mathbf{z} in $[H_0^1(\Omega)]^d$. As $[H_0^1(\Omega)]^d$ is compactly embedded in $[L^2(\Omega)]^d$, we may assume that this sequence converges strongly in $[L^2(\Omega)]^d$.

Recall that space $Y_h^{(0)}$ is defined by

$$Y_h^{(0)} = \left\{ \mathbf{v} \in [L^2(\Omega)]^d \mid \mathbf{v}|_K \text{ is constant}, \quad \forall K \in \mathcal{T}_h \right\}. \quad (84)$$

Due to standard finite element interpolation analysis, there exists a sequence $\{\mathbf{y}_h\}_{h>0}$, with $\mathbf{y}_h \in Y_h^{(0)}$, strongly convergent to \mathbf{z} in $[L^2(\Omega)]^d$ (even if the family of triangulations is not regular).

Denote by Y^* the reference space that generates space $Y_h^{(0)}$. By hypothesis, $Y^* \cap Z^* = \{0\}$. Then, there exists $\rho > 0$ such that

$$|(\mathbf{z}^*, \mathbf{y}^*)_{K^*}| \leq (1 - \rho) \|\mathbf{z}^*\|_{0,K^*} \|\mathbf{y}^*\|_{0,K^*}, \quad \forall \mathbf{z}^* \in Z^*, \mathbf{y}^* \in Y^*.$$

This is proved similarly to estimate (83) in Lemma 2. Thus,

$$\begin{aligned} |(\mathbf{z}_h, \mathbf{y}_h)| &= \sum_{K \in \mathcal{T}_h} |\det A_K| |(\mathbf{z}_h^K, \mathbf{y}_h^K)_{K^*}| \leq (1 - \rho) \sum_{K \in \mathcal{T}_h} |\det A_K| \|\mathbf{z}_h^K\|_{0,K^*} \|\mathbf{y}_h^K\|_{0,K^*} \\ &\leq (1 - \rho) \|\mathbf{z}_h\|_0 \|\mathbf{y}_h\|_0. \end{aligned}$$

Consequently, $\mathbf{z} = 0$ as $\|\mathbf{z}\|_0^2 = \lim_{h \rightarrow 0} |(\mathbf{z}_h, \mathbf{y}_h)| \leq (1 - \rho) \|\mathbf{z}\|_0^2$.

As the limit of any weakly convergent subsequence is necessarily zero, the the whole sequence $\{\mathbf{z}_h\}_{h>0}$ converges weakly to zero. \square

REFERENCES

- [1] C. Amrouche and V. Girault, Decomposition of Vector spaces and application to the Stokes problem in arbitrary dimensions. *Czechoslovak Math. J.* **44** (1994) 109–140.
- [2] O. Besson and M. R. Laydi, Some estimates for the anisotropic Navier–Stokes equations and for the hydrostatic approximation. *RAIRO-Modél. Math. Anal. Numér.* **26** (1992) 855–865.
- [3] I. Babuška, The Finite Element Method with Lagrange multipliers. *Numer. Math.* **20** (1973) 179–192.
- [4] C. Baiocchi, F. Brezzi and L. P. Franca, Virtual Bubbles and Galerkin-least-squares type methods (Ga.L.S.). *Comput. Methods Appl. Mech. Engrg.* **105** (1993) 125–141.
- [5] C. Bernardi and Y. Maday, *Approximations spectrales de problèmes aux limites elliptiques*. Springer-Verlag, Berlin (1992).
- [6] H. Brézis, *Analyse Fonctionnelle*. Masson, Paris (1983).
- [7] F. Brezzi, On the existence, uniqueness and approximation of saddle-point problems arising from Lagrange Multipliers. *RAIRO-Anal. Numér.* **R2** (1974) 129–151.
- [8] F. Brezzi and J. Douglas, Stabilized mixed methods for the Stokes problem. *Numer. Math.* **53** (1988) 225–236.

- [9] F. Brezzi and J. Pitkäranta, On the stabilization of Finite Element approximations of the Stokes problem, in *Efficient Solutions for Elliptic Systems. Notes on Numerical Fluid Mechanics* **10**, W. Hackbusch Ed., Springer-Verlag, Berlin (1984) 11–19.
- [10] T. Chacón Rebollo, A term by term Stabilization Algorithm for Finite Element solution of incompressible flow problems. *Numer. Math.* **79** (1998) 283–319.
- [11] T. Chacón Rebollo and A. Domínguez Delgado, A unified analysis of Mixed and Stabilized Finite Element Solutions of Navier-Stokes equations. *Comput. Methods Appl. Mech. Engrg.* **182** (2000) 301–331.
- [12] T. Chacón Rebollo and F. Guillén González, An intrinsic analysis of existence of solutions for the hydrostatic approximation of Navier-Stokes equations. *C. R. Acad. Sci. Paris, Série I* **330** (2000) 841–846.
- [13] P.G. Ciarlet, *The Finite Element Method for Elliptic Problems*. North-Holland, Amsterdam (1978).
- [14] L.P. Franca and S.L. Frey, Stabilized Finite Elements: II. The incompressible Navier-Stokes equations. *Comput. Methods Appl. Mech. Engrg.* **99** (1992) 209–233.
- [15] L.P. Franca and R. Stenberg, Error analysis for some Galerkin-Least-Squares methods for the elasticity equations. *SIAM J. Numer. Anal.* **28** (1991) 1680–1697.
- [16] L.P. Franca, T.J.R. Hughes and R. Stenberg, Stabilized Finite Element Methods, in *Incompressible Computational Fluid Dynamics*, M.D. Gunzburger and R.A. Nicolaides Eds., Cambridge Univ. Press, New York (1993).
- [17] V. Girault and P.A. Raviart, *Finite Element Methods for Navier-Stokes equations*. Springer-Verlag, Berlin (1988).
- [18] R. Dautray and L.L. Lions, *Analyse Mathématique et Calcul Numérique pour les Sciences et les Techniques*. Masson, Paris (2000).
- [19] P. Gervasio and F. Saleri, Stabilized Spectral Element approximation for the Navier-Stokes equations. *Numer. Methods Partial Differential Eq.* **14** (1988) 115–141.
- [20] T.J.R. Hughes and L.P. Franca, A new Finite Element formulation for CFD: VII. The Stokes problem with various well-posed boundary conditions: Symmetric formulations that converge for all velocity/pressure spaces. *Comput. Methods Appl. Mech. Engrg.* **65** (1987) 85–96.
- [21] T.J.R. Hughes, L.P. Franca and M. Balestra, A new Finite Element formulation for CFD: V. Circumventing the Brezzi-Babuška condition: A stable Petrov-Galerkin formulation of the Stokes problem accommodating equal-order interpolations. *Comput. Methods Appl. Mech. Engrg.* **59** (1986) 85–99.
- [22] P. Knobloch and L. Tobiska, Stabilization methods of Bubble type for the $Q_1 - Q_1$ -Element applied to the incompressible Navier-Stokes equations. *ESAIM: M2AN* **34** (2000) 85–107.
- [23] R. Lewandowski, *Analyse Mathématique et Océanographie*. Masson, Paris (1997).
- [24] J.L. Lions, R. Temam and S. Wang, New formulation of the primitive equations of the atmosphere and applications. *Nonlinearity* **5** (1992) 237–288.
- [25] R. Pierre, Simple C^0 -approximations for the computation of incompressible flows. *Comput. Methods Appl. Mech. Engrg.* **68** (1989) 205–228.
- [26] G. Russo, Bubble stabilization of Finite Element Methods for the linearized incompressible Navier-Stokes equations. *Comput. Methods Appl. Mech. Engrg.* **132** (1996) 335–343.
- [27] L. Tobishka and R. Verfürth, Analysis of a Streamline Diffusion finite element method for the Stokes and Navier-Stokes equations. *SIAM J. Numer. Anal.* **33** (1996) 107–127.
- [28] R. Verfürth, Analysis of some Finite Element solutions for the Stokes Problem. *RAIRO-Anal. Numér.* **18** (1984) 175–182.