

COARSE-GRAINING SCHEMES AND *A POSTERIORI* ERROR ESTIMATES FOR STOCHASTIC LATTICE SYSTEMS

MARKOS A. KATSOUKAKIS¹, PETR PLECHÁČ², LUC REY-BELLET¹
AND DIMITRIOS K. TSAGKAROGIANNIS³

Abstract. The primary objective of this work is to develop coarse-graining schemes for stochastic many-body microscopic models and quantify their effectiveness in terms of *a priori* and *a posteriori* error analysis. In this paper we focus on stochastic lattice systems of interacting particles at equilibrium. The proposed algorithms are derived from an initial coarse-grained approximation that is directly computable by Monte Carlo simulations, and the corresponding numerical error is calculated using the specific relative entropy between the exact and approximate coarse-grained equilibrium measures. Subsequently we carry out a cluster expansion around this first – and often inadequate – approximation and obtain more accurate coarse-graining schemes. The cluster expansions yield also sharp *a posteriori* error estimates for the coarse-grained approximations that can be used for the construction of adaptive coarse-graining methods. We present a number of numerical examples that demonstrate that the coarse-graining schemes developed here allow for accurate predictions of critical behavior and hysteresis in systems with intermediate and long-range interactions. We also present examples where they substantially improve predictions of earlier coarse-graining schemes for short-range interactions.

Mathematics Subject Classification. 65C05, 65C20, 82B20, 82B80, 82-08.

Received August 1st, 2006. Revised February 1st, 2007.

INTRODUCTION

In the recent years there has been a growing interest in developing and analyzing hierarchical coarse-graining methods for the purpose of modeling and simulation across scales for systems arising in a broad spectrum of scientific disciplines ranging from materials science to biology and atmosphere/ocean science. Typically in microscopic simulations of complex systems the model is formulated in terms of simple rules describing interactions between small-scale degrees of freedom such as individual particles or spin variables. On the other hand computational difficulties arise immediately from evaluating their interactions for any realistic size spatio-temporal scales. When a coarse-grained model becomes available, it has fewer observables than the original microscopic system making it computationally more efficient than the direct numerical simulations. At the same time it is expected that it can describe accurately the unresolved degrees of freedom. The coarse-graining

Keywords and phrases. Coarse-graining, *a posteriori* error estimate, relative entropy, lattice spin systems, Monte Carlo method, Gibbs measure, cluster expansion, renormalization group map.

¹ Department of Mathematics, University of Massachusetts, USA. markos@math.umass.edu; lr7q@math.umass.edu

² Department of Mathematics, University of Tennessee, USA. plechac@math.utk.edu

³ Max Planck Institute for Mathematics in the Sciences, Germany. tsagkaro@mis.mpg.de

strategy we are pursuing here is attempting to address these goals in the context of equilibrium sampling of stochastic lattice systems by combining methods from statistical mechanics and perturbation analysis.

In this paper we consider stochastic lattice systems such as Ising-type models as a paradigm of hierarchical coarse-graining that can provide an explicit numerical method with prescribed error tolerance. Such lattice systems for N particles are defined in terms of a microscopic lattice Hamiltonian $H_N(\sigma)$ with σ being the microscopic configuration and an *a-priori* Bernoulli measure $P_N(d\sigma)$. We perform a coarse-graining by subdividing the lattice into coarse cells and defining variables η on each coarse cell to be the total magnetization in the cell. The corresponding renormalization group map (known as the Kadanoff transform) [7, 12] is defined by the formula

$$e^{-\beta\bar{H}_M(\eta)} = \int e^{-\beta H_N(\sigma)} P_N(d\sigma|\eta),$$

where $\bar{H}_M(\eta)$ is the Hamiltonian at the coarse level and $P_N(d\sigma|\eta)$ is the conditional probability of having a microscopic configuration σ given a configuration η at the coarse level.

Such a Hamiltonian defined on a coarser level than the microscopic, is an exact equivalent of the microscopic Hamiltonian H_N , in the sense that the finer degrees of freedom have been averaged. However, it cannot be easily calculated explicitly and hence used in numerical simulations. Our perspective is to approximate it by viewing it as a perturbation of a coarse-grained approximating Hamiltonian $\bar{H}_M^{(0)}$ suggested in [14, 15] and defined by

$$\bar{H}_M^{(0)}(\eta) = \int H_N(\sigma) P_N(d\sigma|\eta).$$

A closely related coarse-grained Hamiltonian was suggested independently in [10, 11], where it was constructed in an equilibrium context using a wavelet expansion.

Using this first approximation we have

$$\bar{H}_M(\eta) = \bar{H}_M^{(0)}(\eta) - \frac{1}{\beta} \log \int e^{-\beta(H_N(\sigma) - \bar{H}_M^{(0)}(\eta))} P_N(d\sigma|\eta).$$

The fact that the conditional probability $P_N(d\sigma|\eta)$ factorizes at the level of the coarse cells allows us to use cluster expansion techniques to write a series expansion for $\bar{H}_M(\eta)$ around $\bar{H}_M^{(0)}$. We obtain the following series

$$\bar{H}_M(\eta) = \bar{H}_M^{(0)}(\eta) + \bar{H}_M^{(1)}(\eta) + \dots + \bar{H}_M^{(p)}(\eta) + \mathcal{O}(\epsilon^{p+1}), \quad p = 2, \dots \quad (1)$$

uniformly in η , where the correction terms $\bar{H}_M^{(1)}(\eta)$, $\bar{H}_M^{(2)}(\eta)$, *etc.*, can be calculated explicitly with the relevant errors and ϵ is a small parameter depending on the characteristics of the coarse-graining, the potential and the inverse temperature. In this paper we first show that this strategy works well provided that the interactions have a range which is long compared to the size of the coarse cells and that they vary slowly over the size of a coarse cell. The parameter ϵ (given explicitly in (20)) encapsulates these conditions.

Notice that we do *not* perform the usual high-temperature expansion using the Bernoulli product measure $P_N(d\sigma)$ but rather expand the Hamiltonian around a well-chosen first approximation using the product structure at the coarse level of the conditional probability $P_N(d\sigma|\eta)$. This allows us to construct these approximating Hamiltonians well-beyond the temperature range allowed in a standard high-temperature expansion. The basic tool for cluster expansions is the reformulation of the system in terms of what is called the polymer model. This technique originates from Mayer [19] and Peierls [20] for the case of high/low temperature respectively. For the high temperature case, a first rigorous proof was given in [6], while our approach is based on the polymer system introduced in [8]. For an overview of these methods and references we refer to [23].

Clearly the choice of the first approximation $\bar{H}_M^{(0)}$ is crucial to our method and it should be such that (i) it is explicitly computable as the constructions in [10, 14], and (ii) it provides a good estimate between the microscopic and the 2nd-order coarse-graining. In a subsequent paper [16] we will show how to apply these

ideas and techniques to systems with both short and long-range interactions. In this case the choice of the first approximation $\bar{H}_M^{(0)}$ will not be given in a closed form but it will be computed numerically in an efficient way.

All error estimates are calculated in terms of the specific relative entropy of the corresponding equilibrium Gibbs measures. Specific relative entropy represents the loss of information in the transition from the microscopic to the coarse-grained models and here is used to assess the information compression for the same level of coarse graining in schemes that are of a higher or lower order, determined by the truncation level p in (1). The primary practical purpose of the higher-order corrections is in allowing us to extend the regime of validity of the expansion and to obtain very accurate coarse grainings of the Gibbs measure even if the parameter ϵ in (1) is not necessarily much smaller than one. We refer to examples in Section 4 where even for $\epsilon = \mathcal{O}(1)$, we obtain an accurate prediction of the hysteresis when we include the higher-order correction terms.

The error analysis developed in this paper also provides an error expansion that can be computed and tracked in the process of a simulation from the numerical data. Such an *a posteriori* error cannot be numerically computed directly from the relative entropy formula, since it involves the calculation of the probability density functions of the microscopic and the coarse-grained measure. However, the expansion in (1) shows a constructive way of calculating the error made, for instance, by the 2nd-order coarse-graining $\bar{H}_M^{(0)}(\eta)$ in terms of other coarse observables given by the higher-order correction terms $\bar{H}_M^{(p)}$ plus a controlled error of order $\mathcal{O}(\epsilon^{p+1})$. Similarly to the numerical analysis of approximations for PDEs, we derive *a priori* and *a posteriori* error estimates between the exact microscopic solution and the approximating coarse-grained one. In contrast to the PDE setting where such error analysis is calculated in a suitable norm, here the error is measured in terms of the specific relative entropy. As in the case of PDEs such *a posteriori* errors are also useful for constructing adaptive methods. For some earlier work on adaptivity for stochastic systems see, *e.g.*, [3, 4, 24].

The paper is organized as follows: in Section 1 we present the main results and an outline of our methods. In Section 2 we discuss cluster expansions in the context of coarse-graining and derive the effective coarse-grained Hamiltonian as an abstract expansion. In Section 3 we calculate the first terms in the expansion of the Hamiltonian and formulate two concrete numerical schemes; Scheme 3.1 is second-order accurate, while Scheme 3.2 is third-order accurate. In Section 4 we present simulations with these schemes in a demanding phase transition regime. Appendix A includes the details of our analytical calculations in Section 3. Appendix B contains a brief description of computational background for the Monte Carlo algorithms used in the sampling of microscopic and coarse-grained Gibbs states carried out in Section 4.

1. MAIN RESULTS AND OUTLINE OF THE METHOD

1.1. Microscopic models

We consider as the physical domain for the system the d -dimensional torus $\mathbb{T}_d := [0, 1)^d$ with periodic boundary conditions. The microscopic system consists of a uniform lattice $\Lambda_N := (\frac{1}{n}\mathbb{Z})^d \cap \mathbb{T}_d$. The number of lattice sites $N = n^d$ is fixed, but arbitrary and finite. We consider here periodic boundary conditions, but other boundary conditions can be accommodated easily.

The spin (or order parameter) $\sigma(x)$ takes values in $\{+1, -1\}$ at each lattice site $x \in \Lambda_N$. A spin configuration $\sigma = \{\sigma(x)\}_{x \in \Lambda_N}$ is an element of the configuration space $\mathcal{S}_N := \{+1, -1\}^{\Lambda_N}$. The energy of the configuration σ is given by the Hamiltonian

$$H_N(\sigma) = -\frac{1}{2} \sum_{x \in \Lambda_N} \sum_{y \neq x} J(x-y) \sigma(x) \sigma(y) + h \sum_{x \in \Lambda_N} \sigma(x), \quad (2)$$

where the two-body inter-particle potential J describes the interaction between individual spins and h is an external field.

The strength of the potential is measured by $\|J\| \equiv \sum_{x \neq 0} |J(x)|$, *i.e.* we assume that the two-body potential is summable. As we will scale later the Hamiltonian with the inverse temperature β we can assume, without loss of generality, that $\|J\| = 1$.

Example 1.1 (Nearest-neighbor interaction). In this case the spin at site x interacts only with its nearest neighbors on the lattice Λ_N , *i.e.*

$$J(x-y) = \begin{cases} J & \text{if } |x-y| = \frac{1}{n}; \\ 0 & \text{otherwise.} \end{cases}$$

Example 1.2 (Finite-range interactions). A spin at site x interacts with its neighbors which are at most L lattice points away from x . It will be useful to consider the range of the interaction L as a parameter of the model. To do this let

$$V: \mathbb{R}^+ \rightarrow \mathbb{R}, \quad V(r) = 0, \quad \text{if } |r| \geq 1. \quad (3)$$

Then the potential $J(x-y)$ can be taken to have the form

$$J(x-y) = \frac{1}{L^d} V\left(\frac{n}{L}|x-y|\right), \quad x, y \in \Lambda_N. \quad (4)$$

The factor $1/L^d$ in (4) is a normalization which ensures that the strength of the potential J is essentially independent of L and we have $\|J\| \simeq \int |V(r)| dr$. Note also that Example 1.1 can be obtained from Example 1.2 by setting $L = 1$.

Example 1.3 (Long-range interactions). In this case we assume that a spin interacts with all spins on the lattice Λ_N *via* a summable interaction $J(x)$. Since our goal is to construct coarse-grained approximations of the Hamiltonian suitable for numerical simulations it will be convenient to truncate long-range interactions and control the error term. In order to do this we choose a small parameter θ and choose L such that

$$\sum_{\{x: |x| \geq \frac{L}{n}\}} |J(x)| \leq \theta.$$

The parameter L can be thought as the effective range of the potential and we can truncate the long-range potential $J(x)$ by setting $J(x) = 0$ if $|x| \geq \frac{L}{n}$. If we denote by \hat{J} the truncated potential and by \hat{H}_N the corresponding Hamiltonian it is easy to see that

$$\frac{1}{N} |H_N(\sigma) - \hat{H}_N(\sigma)| = \mathcal{O}(\theta),$$

i.e., the error per unit volume is of order θ . We can then assume that the truncated potential has the form (4).

Example 1.4 (Kac-type interactions). Mean-field interactions are obtained formally by taking a finite range interaction and setting $L = n$ in (4)

$$J(x-y) = \frac{1}{N} V(|x-y|), \quad x, y \in \Lambda_N. \quad (5)$$

Note that this is different from a summable long-range interaction since the potential is scaled with the size of the system.

The potential in Example 1.2 will be central to our analysis. Both Examples 1.1 and 1.4 are obtained in suitable limiting cases ($L = 1$ and $L = n$ respectively) and the potential in Example 1.3 can be approximated by such potentials (see the error estimates below).

The finite-volume equilibrium states of the system are given by the canonical Gibbs measure

$$\mu_{N,\beta}(d\sigma) = \frac{1}{Z_N} e^{-\beta H_N(\sigma)} P_N(d\sigma), \quad (6)$$

where β is the inverse temperature, Z_N is the normalizing partition function, and $P_N(d\sigma)$, the prior distribution on \mathcal{S}_N , is the product measure

$$P_N(d\sigma) = \prod_{x \in \Lambda_N} \rho(d\sigma(x)).$$

A typical choice $\rho(\sigma(x) = +1) = \frac{1}{2}$ and $\rho(\sigma(x) = -1) = \frac{1}{2}$ is the distribution of a Bernoulli random variable for each $x \in \Lambda_N$.

1.2. Specific relative entropy

One of the obvious issues arising in any attempt to coarse-grain microscopic systems is the evaluation of the numerical error as we move from finer to coarser scales. This error essentially involves the comparison of the microscopic and the coarse-grained probability measures. A natural way to compare two probability measures is to compute their relative entropy. Let $\pi_1(\sigma)$ and $\pi_2(\sigma)$ be two probability measures defined on a common probability space, with the state space \mathcal{S} . The relative entropy, or Kullback-Leibler distance, of π_1 with respect to π_2 , is defined as

$$\mathcal{R}(\pi_1 | \pi_2) \equiv \int_{\mathcal{S}} \log \left(\frac{d\pi_1}{d\pi_2} \right) d\pi_1, \quad (7)$$

where $d\pi_1/d\pi_2$ is the Radon-Nikodym derivative of π_1 with respect to π_2 . On a countable state space \mathcal{S} we obtain

$$\mathcal{R}(\pi_1 | \pi_2) = \sum_{\sigma \in \mathcal{S}} \pi_1(\sigma) \log \frac{\pi_1(\sigma)}{\pi_2(\sigma)}.$$

In information theory the relative entropy $\mathcal{R}(\pi_1 | \pi_2)$ provides a measure of “information loss” or “information distance” of π_1 compared to π_2 . In our context we use the relative entropy in order to assess the information compression of different coarse-graining schemes. Basic properties and applications in information theory can be found in [5].

Using Jensen’s inequality it is not difficult to show that

$$\begin{aligned} \mathcal{R}(\pi_1 | \pi_2) &\geq 0 \quad \text{and,} \\ \mathcal{R}(\pi_1 | \pi_2) &= 0 \quad \text{if and only if } \pi_1(\sigma) = \pi_2(\sigma) \text{ for all } \sigma \in \mathcal{S}. \\ \mathcal{R}(\pi_1 | \pi_2) &= \infty \quad \text{if } \pi_1 \text{ is not absolutely continuous with respect to } \pi_2. \end{aligned}$$

The relative entropy is not a metric (it is not symmetric), but one can use the relative entropy to bound the total variation distance of the measures π_1 and π_2 as demonstrated by *Csiszár-Kullback-Pinsker inequality* [5]

$$\mathcal{R}(\pi_1 | \pi_2) \geq \frac{1}{2} \left(\sum_{\sigma \in \mathcal{S}} |\pi_1(\sigma) - \pi_2(\sigma)| \right)^2 \equiv \frac{1}{2} \|\pi_1 - \pi_2\|_{TV}^2, \quad (8)$$

or equivalently using the dual form of the total variation norm

$$\sup_{\|f\|_{\infty}=1} \left| \int f d\pi_1 - \int f d\pi_2 \right| \leq \sqrt{\frac{1}{2} \mathcal{R}(\pi_1 | \pi_2)}, \quad (9)$$

where $\|\cdot\|_{\infty}$ denotes the usual sup-norm.

In statistical mechanics, the entropy, which is the relative entropy with respect to the prior distribution, and the relative entropy play a prominent role, in particular in the variational principles. The relative entropy is, in general, difficult to compute since it requires the sampling of the probability distribution $\pi_1(\sigma)$ and $\pi_2(\sigma)$ which can be prohibitively expensive if the dimension of the state space is large. For Gibbs measures it is, however, a little simpler in the sense that it depends only on the partition functions and the expected values of

the Hamiltonian. Consider, for example, two Gibbs measures $\mu_{N,\beta}^{(1)}$ and $\mu_{N,\beta}^{(2)}$ with Hamiltonians $H_N^{(1)}$ and $H_N^{(2)}$ and partition functions $Z_N^{(1)}$ and $Z_N^{(2)}$. Then we have

$$\mathcal{R} \left(\mu_{N,\beta}^{(1)} \mid \mu_{N,\beta}^{(2)} \right) = \log \left(\frac{Z_N^{(2)}}{Z_N^{(1)}} \right) + \beta \int \left(H_N^{(2)} - H_N^{(1)} \right) d\mu_{N,\beta}^{(1)}. \tag{10}$$

Note that the Hamiltonians and the logarithm of partition functions are extensive quantities: they are proportional to the size of the system N . The formula (10) shows that the entropy of a Gibbs measure and the relative entropy of a Gibbs measure with respect to another Gibbs measure are also extensive quantities. It is therefore natural to compare two Gibbs measures on \mathcal{S}_N by computing their specific relative entropy

$$\frac{1}{N} \mathcal{R} \left(\mu_{N,\beta}^{(1)} \mid \mu_{N,\beta}^{(2)} \right).$$

The specific relative entropy is a measure of the information loss per unit volume and will be our main estimation tool in this paper. A simple example where this interpretation is evident arises if one considers N independent particles and the ensuing scaling of the collective discrepancy if an individual error is committed on the observation of each particle.

1.3. Coarse-grained models

The coarse-graining procedure consists of three steps that we describe separately.

(a) *Coarse graining of the configuration space.* We partition the torus \mathbb{T}_d into $M = m^d$ cells: For $k = (k_1, \dots, k_d) \in \mathbb{Z}^d$ with $0 \leq k_i \leq m - 1$ we define $C_k \equiv [\frac{k_1}{m}, \frac{k_1+1}{m}) \times \dots \times [\frac{k_d}{m}, \frac{k_d+1}{m})$ and we have $\mathbb{T}_d = \cup_k C_k$. We identify each cell C_k with a lattice point of the coarse lattice $\bar{\Lambda}_M = (\frac{1}{m}\mathbb{Z})^d \cap \mathbb{T}_d$. Each coarse cell contains $Q = q^d$ points of the microscopic lattice points with $N \equiv n^d = (mq)^d \equiv MQ$. We will refer to Q as the level of coarse graining ($Q = 1$ corresponds to no coarse graining).

We assign a new spin value $\eta(k)$ for the cell C_k according to the rule

$$\eta(k) = \sum_{x \in C_k} \sigma(x).$$

The spin $\eta(k)$ takes values in $\{-Q, -Q + 2, \dots, Q\}$ and the configuration space for the coarse grained system is $\bar{\mathcal{S}}_M \equiv \{-Q, -Q + 2, \dots, Q\}^{\bar{\Lambda}_M}$. We denote by \mathbf{F} the map

$$\mathbf{F} : \mathcal{S}_N \rightarrow \bar{\mathcal{S}}_M, \quad \sigma \mapsto \left\{ \sum_{x \in C_k} \sigma(x) \right\}_k,$$

which assigns a configuration $\eta = \{\eta(k)\}_{k \in \bar{\Lambda}_M}$ on the coarse lattice given a configuration $\sigma = \{\sigma(x)\}_{x \in \Lambda_N}$. An equivalent coarse grained variable, which we shall also use later, is

$$\alpha(k) := \text{card} \{x \in C_k : \sigma(x) = +1\}$$

which takes values in $\{0, 1, \dots, Q\}$. Both coarse variables are equivalent and they are related by the transformation $\eta = 2\alpha - Q$ or $\alpha = \frac{\eta+Q}{2}$.

(b) *Coarse-graining of the prior distribution.* The prior distribution P_N on \mathcal{S}_N induces a new prior distribution on $\bar{\mathcal{S}}_M$ given by $\bar{P}_M = P_N \circ \mathbf{F}^{-1}$, i.e.,

$$\bar{P}_M(\eta) = P_N(\sigma : \mathbf{F}(\sigma) = \eta).$$

Since $\eta(k)$ depends only on the spin $\sigma(x)$, with $x \in C_k$, the measure \bar{P}_M is a product measure

$$\bar{P}_M(d\eta) = \prod_{k \in \bar{\Lambda}_M} \bar{\rho}(d\eta(k)),$$

with

$$\bar{\rho}(\eta(k)) = \left(\frac{Q}{\eta(k)+Q} \right) \left(\frac{1}{2} \right)^Q.$$

The conditional probability $P_N(d\sigma|\eta)$ plays a crucial role in the sequel. Since $\eta(k)$ depends only on the spin $\sigma(x)$ with $x \in C_k$, the probability $P_N(d\sigma|\eta)$ factorizes over the coarse cells. We have

$$P_N(d\sigma|\eta) = \frac{P_N(\sigma \cap \{\mathbf{F}(\sigma) = \eta\})}{\bar{P}_M(\eta)} = \prod_{k \in \bar{\Lambda}_M} \tilde{\rho}_{k,\eta(k)}(d\sigma) \tag{11}$$

and $\tilde{\rho}_{k,\eta(k)}(d\sigma)$ depends only on $\{\sigma(x)\}_{x \in C_k}$. In particular we have

$$\tilde{\rho}_{k,\eta(k)}(\sigma(x) = 1) = \frac{\eta(k) + Q}{2Q} \text{ and } \tilde{\rho}_{k,\eta(k)}(\sigma(x) = -1) = \frac{Q - \eta(k)}{2Q}. \tag{12}$$

To simplify the notation and because our estimates are uniform in $\eta(k)$ we denote this measure simply by $\tilde{\rho}_k$. For a function $f = f(\sigma)$ we define the conditional expectation

$$\mathbb{E}[f|\eta] = \int f(\sigma) P_N(d\sigma|\eta) = \frac{1}{\bar{P}_M(\eta)} \int_{\{\sigma: \mathbf{F}(\sigma) = \eta\}} f(\sigma) P_N(d\sigma) = \int f(\sigma) \prod_k \tilde{\rho}_k(d\sigma).$$

(c) *Coarse-graining of the Hamiltonian.* We want to construct a Hamiltonian $\bar{H}_M(\eta)$ at the coarse-level. A natural definition of such Hamiltonian, as we explain below, is given by the renormalization group block averaging transformation (also known as Kadanoff transformation).

Definition 1.1. The *exact* coarse grained Hamiltonian $\bar{H}_M(\eta)$ is defined by the formula

$$e^{-\beta \bar{H}_M(\eta)} = \mathbb{E}[e^{-\beta H_N} | \eta]. \tag{13}$$

Given the Hamiltonian \bar{H}_M we define the corresponding Gibbs measure by

$$\bar{\mu}_{M,\beta}(d\eta) = \frac{1}{\bar{Z}_M} e^{-\beta \bar{H}_M(\eta)} \bar{P}_M(d\eta). \tag{14}$$

The factor β in front of $\bar{H}_M(\eta)$ is merely a convention, in general the Hamiltonian $\bar{H}_M(\eta)$ does itself depend on β in a nonlinear, complicated way.

Clearly for a fixed value of N , this transformation is well defined. However, from the point of view of statistical mechanics this is not sufficient. We want to construct the coarse-grained Hamiltonian $\bar{H}_M(\eta)$, for any M , as a sum of (summable) many-body interactions. Our original Hamiltonian $H_N(\sigma)$ involves only one-body and two-body interactions but the nonlinear transformation (13) will not preserve this property. The Hamiltonian we will consider in the sequel will have the form

$$K_M(\eta) = \sum_{X \subset \bar{\Lambda}_M} J(X, \{\eta_k\}_{k \in X}) \tag{15}$$

where the $J(X, \{\eta_k\}_{k \in X})$ are translation-invariant many-body interactions involving $\text{card}(X)$ -many different sites. Such an interaction is said to be summable if

$$\sum_{\{X : 0 \in X\}} \|J(X, \{\eta_k\}_{k \in X})\| < \infty.$$

It is well-known [25] that the Kadanoff transformation suffers from some pathologies. At very low temperature the Kadanoff transformation for the nearest-neighbor Ising model with the zero magnetic field is not well-defined in the sense that the coarse-grained Hamiltonian \bar{H}_M does not correspond to a summable interaction. These pathologies are relatively mild [1] and can be eliminated by extending slightly the concept of Gibbs measures. For a comprehensive presentation of this issue see the review paper [25] or also [1] among others. Furthermore, these pathologies will not play any role in our analysis: the cluster expansion techniques we are using exclude the occurrence of these pathologies for the models and the values of the parameter we consider.

The exactness of the coarse graining in the Kadanoff transform is expressed by the fact that the specific relative entropy of $\bar{\mu}_{M,\beta}(d\eta)$ with respect to the coarse-grained Gibbs measure $\mu_{N,\beta} \circ \mathbf{F}^{-1}$ vanishes. Indeed we have

$$\begin{aligned} Z_N &= \int e^{-\beta H_N(\sigma)} P_N(d\sigma) = \int \mathbb{E}[e^{-\beta H_N} | \eta] \bar{P}_M(d\eta) \\ &= \int e^{-\beta \bar{H}_M(\eta)} \bar{P}_M(d\eta) = \bar{Z}_M, \end{aligned} \tag{16}$$

and consequently from (13),

$$\frac{1}{\bar{Z}_M} e^{-\beta \bar{H}_M(\eta)} \bar{P}_M(\eta) = \frac{1}{Z_N} \int_{\{\sigma | \mathbf{F}(\sigma) = \eta\}} e^{-\beta H_N} P_N(d\sigma),$$

and thus

$$\frac{1}{N} \mathcal{R}(\bar{\mu}_{M,\beta} | \mu_{N,\beta} \circ \mathbf{F}^{-1}) = 0.$$

Even for moderately large values of N the exact computation of $\bar{H}_M(\eta)$ is, in general, impractical. The coarse grained Hamiltonian involves not only two-body interactions, but also many-body interactions of an arbitrary number of spins. Our goal is to present a systematic way of calculating explicit approximations of the coarse-grained Hamiltonian \bar{H}_M , to any given degree of accuracy. In the first step we define an approximate Hamiltonian $\bar{H}_M^{(0)}(\eta)$ and we give an *a priori* bound on the blocking error. The choice of $\bar{H}_M^{(0)}$ we make, following [15], is given in the following definition.

Definition 1.2. The first approximation $\bar{H}_M^{(0)}$ of the coarse-grained Hamiltonian \bar{H}_M is given by the formula

$$\bar{H}_M^{(0)}(\eta) \equiv \mathbb{E}[H_N | \eta]. \tag{17}$$

In order to compute $\bar{H}_M^{(0)}(\eta)$ we note that we have

$$\begin{aligned} \mathbb{E}[\sigma(x) | \eta] &= \frac{\eta(k)}{Q}, \quad x \in C_k, \\ \mathbb{E}[\sigma(x)\sigma(y) | \eta] &= \frac{\eta(k)^2 - Q}{Q(Q-1)}, \quad x \in C_k, y \in C_k, y \neq x. \end{aligned}$$

Then

$$\bar{H}_M^{(0)}(\eta) = -\frac{1}{2} \sum_k \sum_{l \neq k} \bar{J}(k-l) \eta(k) \eta(l) - \frac{1}{2} \sum_k \bar{J}(0) (\eta^2(k) - Q) + h \sum_k \eta(k)$$

where

$$\begin{aligned} \bar{J}(k, l) &= \frac{1}{Q^2} \sum_{x \in C_k, y \in C_l} J(x - y), \quad \text{for } k \neq l, \\ \bar{J}(k, k) &= \frac{1}{Q(Q-1)} \sum_{x, y \in C_k, y \neq x} J(x - y), \quad \text{for } k = l. \end{aligned}$$

By defining $\bar{H}_M^{(0)}(\eta)$ one replaces the potential $J(x - y)$ by its average over a coarse cell. Thus the error for the potential is proportional to

$$E_{kl}(x - y) := J(x - y) - \bar{J}(k, l), \quad x \in C_k, y \in C_l,$$

which measures the variation of the potential $J(x - y)$ over a cell. An estimate on the error is provided by the following lemma

Lemma 1.3. *Assume that J satisfies (3)–(4) and assume that $V(r)$ is C^1 .*

(1) *There exists a constant $C > 0$ such that, if $x \in C_k$ and $y \in C_l$, we have*

$$|J(x - y) - \bar{J}(k, l)| \leq 2 \frac{q}{L^{d+1}} \sup_{\substack{x' \in C_k, \\ y' \in C_l}} \|\nabla V(x' - y')\|. \tag{18}$$

(2) *There exists a constant $C > 0$ such that, if $\mathbf{F}(\sigma) = \eta$, we have*

$$\frac{1}{N} \left| H_N(\sigma) - \bar{H}_M^{(0)}(\eta) \right| \leq C \frac{q}{L} \|\nabla V\|_\infty. \tag{19}$$

The lemma will be proved in Section 2.2. Note that the estimates in Lemma 1.3 are not necessarily optimal and can be improved under suitable assumptions on the potential. The importance of Lemma 1.3 lies in the identification of the small parameter in Theorem 1.4 below

$$\epsilon \equiv C \beta \frac{q}{L} \|\nabla V\|_\infty, \tag{20}$$

where C is some constant. Note that we included the inverse temperature β in the parameter, taking into account that β multiplies the Hamiltonian in the Gibbs measure. The parameter ϵ encapsulates the various factors involved in our coarse-graining method.

- (i) *Coarse-graining cell size.* The factor q/L in (20) indicates how the size of the coarse cell governs the effectiveness of the coarse-graining procedure.
- (ii) *Slow variation of the potential.* The factor $\|\nabla V\|_\infty$ in 20 indicates that the coarse-graining will be particularly effective even for large cell sizes if the potential V is slowly varying. This fact is exemplified by considering the extreme case of the Curie-Weiss model where $J(x - y) = J$ is independent of $x - y$. It is easy to see that in this case the coarse-graining is actually *exact*.
- (iii) *Temperature.* At high temperature, *i.e.*, for small β the coarse-graining procedure will work well even for large values of q/L , *i.e.*, for very large coarse cells or for potentials with very short range. This reflects the fact that at high temperature Gibbs measures are actually very close to product measures and that, trivially, product measures can be coarse-grained exactly. On the other hand our methods clearly do not apply to very-low temperatures.

Let us denote by $\bar{\mu}_{M,\beta}^{(0)}(d\eta)$ the Gibbs measure

$$\bar{\mu}_{M,\beta}^{(0)}(d\eta) = \frac{1}{Z_M^{(0)}} e^{-\beta \bar{H}_M^{(0)}(\eta)} \bar{P}_M(d\eta).$$

From Lemma 1.3 and with the help of calculation in the proof of Proposition 2.4 we obtain the bound

$$\frac{1}{N} \mathcal{R}(\bar{\mu}_{M,\beta}^{(0)} | \mu_{N,\beta} \circ \mathbf{F}^{-1}) = O(\epsilon). \tag{21}$$

Furthermore, Lemma 1.3 allows us to perform a cluster expansion and compute higher-order corrections. The basic idea is to rewrite the exact coarse-graining as

$$e^{-\beta \bar{H}_M(\eta)} = e^{-\beta \bar{H}_M^{(0)}(\eta)} \mathbb{E}[e^{-\beta(H_N(\sigma) - \bar{H}_M^{(0)}(\eta))} | \eta],$$

or

$$\bar{H}_M(\eta) = \bar{H}_M^{(0)}(\eta) - \frac{1}{\beta} \log \mathbb{E}[e^{-\beta(H_N(\sigma) - \bar{H}_M^{(0)}(\eta))} | \eta]. \tag{22}$$

Note that the exponential in the second term is not necessarily small, it is in fact of order $N\epsilon$. Cluster expansions are tools which allows to expand such quantities in convergent power series using the independence properties of product measures. The crucial fact here is that conditional measure $P_N(d\sigma | \eta)$ factorizes over the coarse cells. The main result of our paper, proved in Section 3, is

Theorem 1.4. *Let us assume that J satisfies (3)–(4) and that $V(r)$ is C^1 . Then there exists a constant $\delta_0 > 0$ such that if*

$$\delta = Q\epsilon < \delta_0,$$

the Hamiltonian $\bar{H}_M(\eta)$ can be expanded into a convergent series

$$\bar{H}_M(\eta) = \sum_{p=0}^{\infty} \bar{H}_M^{(p)}(\eta),$$

where each term $\bar{H}_M^{(p)}(\eta)$ is a sum of finite-range translation invariant many-body potentials. The first few terms are explicitly calculated in Schemes 3.1 and 3.2 in Section 3. We have the following error bounds uniformly in η and N

$$\frac{\beta}{N} \left(\bar{H}_M(\eta) - (\bar{H}_M^{(0)}(\eta) + \dots + \bar{H}_M^{(p)}(\eta)) \right) = \mathcal{O}(\epsilon^{p+1}).$$

If we define the Gibbs measures

$$\bar{\mu}_{M,\beta}^{(p)}(d\eta) = \frac{1}{\bar{Z}_M^{(p)}} e^{-\beta(\bar{H}_M^{(0)}(\eta) + \dots + \bar{H}_M^{(p)}(\eta))} \bar{P}_M(d\eta),$$

then the following bounds for the relative entropy per unit volume hold

$$\frac{1}{N} \mathcal{R}(\bar{\mu}_{M,\beta}^{(0)} | \mu_{N,\beta} \circ \mathbf{F}^{-1}) = \mathcal{O}(\epsilon^2), \tag{23}$$

$$\frac{1}{N} \mathcal{R}(\bar{\mu}_{M,\beta}^{(p)} | \mu_{N,\beta} \circ \mathbf{F}^{-1}) = \mathcal{O}(\epsilon^{p+1}), \tag{24}$$

where $p = 2, \dots$ and ϵ is given by (20).

We note that in information theory the relative entropy provides a measure of “information distance” of two probability measures. In our context we use the relative entropy estimate of Theorem 1.4 in order to assess the information compression of different coarse-graining schemes. According to how many correction terms are included in the expansion corresponding to different truncation levels p , such as Schemes 3.1 and 3.2 in Section 3, we quantify the amount of the information loss when the measures are compared at the same level of coarse graining. Furthermore, it would be mathematically interesting and numerically useful to study the implications of the relative entropy error estimates for weak errors of specific observables; for example, expected times for domain switching in regimes that exhibit phase transitions such as those discussed in Section 4.

Remark 1.5.

- (1) The cluster expansion provides an explicit algorithm to compute the higher-order approximations $\bar{H}_M^{(p)}$. For example, the next order correction is given by $\bar{H}_M^{(1)} + \bar{H}_M^{(2)}$ as in Scheme 3.2 in Section 3. The analytical computations required for determining higher-order corrections become quickly very involved, nonetheless they can be carried out with symbolic computational tools.
- (2) The primary practical purpose of the higher-order estimates in Theorem 1.4 is in allowing us to extend the regime of validity of the expansion and obtain very accurate coarse-graining of the Gibbs measure even if the parameter ϵ given by (20) is not necessarily much smaller than one. We refer to the examples in Section 4 where even for $q \geq L$, and $\beta > 1$, *i.e.*, $\epsilon = \mathcal{O}(1)$, we get an accurate prediction of critical behavior when we include the higher-order correction terms.
- (3) The reader might wonder about the appearance of a new small parameter $\delta = Q\epsilon$ in the statement of Theorem 1.4. The reason lies in the details of the cluster expansion. Since the conditional measure $P_N(d\sigma|\eta)$ factorizes over the coarse cells, it is required for the cluster expansion to converge that the error in every coarse cell is small. The error for each site of the microscopic lattice is of order ϵ and thus the error for a coarse cell is of order $\delta = Q\epsilon$.
- (4) Note further that the bound (23) improves on the bound (21) and that shows the first approximation $\bar{H}_M^{(0)}$ is actually already a second-order method. This is due to cancellations which follow directly from the definition of $\bar{H}_M^{(0)}$, see Definition 1.2 and (46) below.

In the previous theorem the errors are calculated with respect to the relative entropy of the corresponding equilibrium measures. Apart from this *a priori* estimate, we would also like to have a formulation of the error that can be explicitly computed, from the data of the simulation. In our case, such an *a posteriori* error cannot be numerically computed directly from the relative entropy formula, since it involves the calculation of the probability density functions of the microscopic and the coarse-grained measure. However, the error estimate of Theorem 1.4 provides us with an explicit way of calculating the error made, for instance, by the 2nd-order coarse-graining. In Section 3 we prove the *a posteriori* error estimate for $\mathcal{R}(\bar{\mu}_{M,\beta}^{(0)}|\mu_{N,\beta} \circ \mathbf{F}^{-1})$ given in the theorem

Theorem 1.6 (*A posteriori error*). *We have*

$$\mathcal{R}(\bar{\mu}_{M,\beta}^{(0)}|\mu_{N,\beta} \circ \mathbf{F}^{-1}) = \beta \mathbb{E}_{\bar{\mu}_{M,\beta}^{(0)}} [R(\eta)] + \log \left(\mathbb{E}_{\bar{\mu}_{M,\beta}^{(0)}} \left[e^{\beta R(\eta)} \right] \right) + \mathcal{O}(\epsilon^3),$$

where the residuum operator $R(\cdot)$ is given by

$$R(\eta) = \bar{H}_M^{(1)}(\eta) + \bar{H}_M^{(2)}(\eta).$$

Note that the quantity $\mathbb{E}_{\bar{\mu}_{M,\beta}^{(0)}} [R(\eta)]$ can be calculated on-the-fly using the coarse simulation. Calculations involving the *a posteriori* error estimation and related adaptive methods will be discussed in a forthcoming publication. Earlier work that uses only an upper bound and not the sharp estimate of Theorem 1.6 can be found in [3, 4].

2. CLUSTER EXPANSION AND EFFECTIVE INTERACTIONS

In this section we expand the term $\mathbb{E}[e^{-\beta(H_N(\sigma) - \bar{H}_M^{(0)}(\eta))}|\eta]$ in (22) into a convergent series using a cluster expansion.

2.1. Introduction to the polymer model

It is convenient to choose an ordering on the d -dimensional lattice $\bar{\Lambda}_M$. For example, the lexicographic ordering defines for $d = 2$ the relation $(k_1, k_2) \leq (l_1, l_2)$ if and only if $k_1 < l_1$ or $k_1 = l_1$ and $k_2 \leq l_2$ and is

then extended recursively for arbitrary dimension d . To set-up the cluster expansion we write the difference $H_N(\sigma) - \bar{H}_M^{(0)}(\eta)$ as

$$H_N(\sigma) - \bar{H}_M^{(0)}(\eta) = \sum_{k \leq l} \Delta_{kl} J(\sigma), \quad \text{where} \tag{25}$$

$$\Delta_{kl} J(\sigma) := -\frac{1}{2} \sum_{\substack{x \in C_k \\ y \in C_l, y \neq x}} (J(x - y) - \bar{J}(k, l)) \sigma(x) \sigma(y) (2 - \delta_{kl}).$$

Note that we have excluded the external field contribution to the Hamiltonian, the role of which will be presented as a remark at the end of this section. In order to expand the exponential $e^{-\beta(H_N(\sigma) - \bar{H}_M^{(0)}(\eta))}$ into a series we set

$$f_{kl}(\sigma) := e^{-\beta \Delta_{kl} J(\sigma)} - 1 \tag{26}$$

and obtain, using the factorization properties of $P_N(d\sigma|\eta)$

$$\mathbb{E}[e^{-\beta(H_N(\sigma) - \bar{H}_M^{(0)}(\eta))} | \eta] = \int \prod_{k \leq l} (1 + f_{kl}) \prod_k \tilde{\rho}_k(d\sigma). \tag{27}$$

We give a short description of the polymer model that we will use in order to organize the cluster expansion. The product $\prod_{k \leq l} (1 + f_{kl}(\sigma))$ can be expressed as $\sum_{G \in \mathcal{G}_M} \prod_{\{k, l\} \in G} f_{kl}$, where \mathcal{G}_M is a collection of all simple graphs on $M = m^d$ vertices, *i.e.*, graphs where each edge appears only once, with additional loops (k, k) on the same vertex. A one-dimensional example of an element G is $G = \{\{1, 2\}, \{2, 2\}, \{2, 5\}, \{3, 6\}\}$ which corresponds to the term $f_{12} f_{22} f_{25} f_{36}$ in the above sum. We observe further that each G can be divided into connected non-intersecting components called polymers, where by non-intersecting we mean that they do not share the same coarse cells. We define the support $\text{supp}(\gamma)$ of a polymer γ the set of the coarse cells participating in it. In the previous example, $G = (\gamma_1, \gamma_2)$ where $\gamma_1 = \{\{1, 2\}, \{2, 2\}, \{2, 5\}\}$ and $\gamma_2 = \{\{3, 6\}\}$. We also say γ_1 is incompatible with γ_2 and we write $\gamma_1 \not\sim \gamma_2$ if $\text{supp}(\gamma_1) \cap \text{supp}(\gamma_2) = \emptyset$. Then (27) is equal to

$$\int \sum_{n \geq 1} \frac{1}{n!} \sum_{\substack{(\gamma_1, \dots, \gamma_n) \\ \gamma_i \not\sim \gamma_j, i \neq j}} \prod_{i=1}^n \prod_{\{k, l\} \in \gamma_i} f_{kl} \prod_k \tilde{\rho}_k(d\sigma) = \sum_{n \geq 1} \frac{1}{n!} \sum_{\substack{(\gamma_1, \dots, \gamma_n) \\ \gamma_i \not\sim \gamma_j, i \neq j}} \prod_{i=1}^n \int \prod_{\{k, l\} \in \gamma_i} f_{kl} \prod_{\{k\} \in \text{supp}(\gamma_i)} \tilde{\rho}_k(d\sigma),$$

since the polymers $\gamma_1, \dots, \gamma_n$ do not share the same coarse cells. Note that the factor $\frac{1}{n!}$ takes into account the fact that by relabeling the connected polymers $\gamma_1, \dots, \gamma_n$ we get the same G .

An equivalent formulation is to consider as our building blocks the more fundamental (than the polymers) clusters $R \subset \bar{\Lambda}_M$ (*i.e.* the set of vertices appearing in each polymer). For example, the corresponding cluster to γ_1 is $R_1 = \{1, 2, 5\}$. Note that to each cluster there are several polymers that correspond to it. To prevent confusion we warn the reader that what we call clusters R , it is called polymers in [2]. We use the name cluster to distinguish it from the previous description of vertices with edges, that we have called polymers. Using clusters the expansion becomes

$$\sum_{n \geq 1} \frac{1}{n!} \sum_{\substack{(R_1, \dots, R_n) \in \mathcal{R}^n \\ R_i \cap R_j = \emptyset, i \neq j}} \prod_{i=1}^n \zeta(R_i), \quad \text{where} \tag{28}$$

$$\zeta(R) = \int \sum_{g \in G_R} \prod_{\{k, l\} \in g} f_{kl}(\sigma) \prod_{\{k\} \in R} \tilde{\rho}_k(d\sigma)$$

is called the *activity* of the cluster R . By \mathcal{R} we denote the space of clusters $R \subset \bar{\Lambda}_M$ with $|R| = \text{card}(R) \geq 1$ and G_R is the set of generalized connected graphs on the set R . A generalized connected graph on R is a collection

of distinct subsets of R , in which case the vertices of the generalized graph are the indices of the enumeration of the subsets, and any two subsets of the vertices of the graph are linked to each other. The edges of this graph are defined whenever the distinct subsets share a common element. For more details we refer to [22]. Thus we are in the context of [2], Theorem 2, and we can formulate our first lemma.

Lemma 2.1 (Expansion of the Hamiltonian). *If for every integer $r \geq 2$, ζ satisfies the condition*

$$\sup_{k \in \bar{\Lambda}_M} \sum_{\substack{R \in \mathcal{R}, R \supset \{k\} \\ |R|=r}} |\zeta(R)| \leq \delta^{r-1} \tag{29}$$

for some δ such that $\sup_{k \in \bar{\Lambda}_M} \sum_{\substack{R \in \mathcal{R}, R \supset \{k\} \\ |R|=1}} |\zeta(R)| \leq \delta$ and $\delta < \frac{1}{6}$, then

$$\bar{H}_M(\eta) = \bar{H}_M^{(0)}(\eta) - \frac{1}{\beta} \sum_{n \geq 1} \frac{1}{n!} \sum_{\substack{(R_1, \dots, R_n) \in \mathcal{R}^n \\ R_i \subset \bar{\Lambda}_M}} \phi(R_1, \dots, R_n) \prod_{i=1}^n \zeta(R_i), \tag{30}$$

where

$$\phi(R_1, \dots, R_n) = \begin{cases} 1, & n = 1 \\ \sum_{g \in G_n} \prod_{\{i,j\} \in g} (1(R_i, R_j) - 1), & n > 1 \end{cases}$$

and G_n is the set of the generalized, connected graphs on $\{1, \dots, n\}$ and

$$1(R_i, R_j) = \begin{cases} 0, & \{R_i \cap R_j \neq \emptyset\}; \\ 1, & \{R_i \cap R_j = \emptyset\}. \end{cases}$$

Proof. The proof is the same as in [2], Theorem 2, given (22) and the discussion preceding Lemma 2.1. For later use we repeat the main steps of the proof. For the series (30) we have

$$\begin{aligned} \sum_{n \geq 1} \frac{1}{n!} \sum_{\substack{(R_1, \dots, R_n) \in \mathcal{R}^n \\ R_i \subset \bar{\Lambda}_M}} \left| \phi(R_1, \dots, R_n) \prod_{i=1}^n \zeta(R_i) \right| = \\ \sum_{R \in \mathcal{R}, R \subset \bar{\Lambda}_M} |\zeta(R)| + \sum_{n \geq 2} \frac{1}{n!} \sum_{\substack{(R_1, \dots, R_n) \in \mathcal{R}^n \\ R_i \subset \bar{\Lambda}_M}} \left| \phi(R_1, \dots, R_n) \prod_{i=1}^n \zeta(R_i) \right| =: S, \end{aligned} \tag{31}$$

where for the first term ($n = 1$) we have

$$\sum_{R \in \mathcal{R}, R \subset \bar{\Lambda}_M} |\zeta(R)| \leq M \sum_{r \geq 1} \sup_{k \in \bar{\Lambda}_M} \sum_{\substack{R \in \mathcal{R}, R \supset \{k\} \\ |R|=r}} |\zeta(R)| \leq M \left(\delta + \sum_{r \geq 2} \delta^{r-1} \right). \tag{32}$$

For the second term ($n \geq 2$) using the estimate from [2], p. 526, (3.6), we obtain

$$\begin{aligned} \sum_{n \geq 2} \frac{1}{n!} \sum_{\substack{(R_1, \dots, R_n) \in \mathcal{R}^n \\ R_i \subset \bar{\Lambda}_M}} \left| \phi(R_1, \dots, R_n) \prod_{i=1}^n \zeta(R_i) \right| &\leq \sum_{n \geq 2} \frac{1}{n!} \sum_{\substack{R \in \mathcal{R} \\ R \subset \bar{\Lambda}_M}} \sum_{\substack{(R_1, \dots, R_n) \in \mathcal{R}^n \\ \exists R_i = R}} \left| \phi(R_1, \dots, R_n) \prod_{i=1}^n \zeta(R_i) \right| \\ &\leq \sum_{n \geq 2} \frac{n}{n!} \sum_{\substack{R \in \mathcal{R} \\ R \subset \bar{\Lambda}_M}} |\zeta(R)| \sum_{(R_2, \dots, R_n) \in \mathcal{R}^{n-1}} \left| \phi(R_1, \dots, R_n) \prod_{i=2}^n \zeta(R_i) \right| \\ &\leq \sum_{n \geq 2} \frac{n}{n!} \sum_{\substack{R \in \mathcal{R} \\ R \subset \bar{\Lambda}_M}} |\zeta(R)| \frac{1}{2} \left(2e \frac{5}{4} \frac{\delta}{1-\delta} \right)^{n-1} (n-2)! |R| e^{|R|}. \end{aligned}$$

Thus,

$$S \leq M(\delta + \sum_{r \geq 2} \delta^{r-1}) + M \frac{1}{2} \sum_{n \geq 2} \frac{1}{n-1} \left(2e \frac{5}{4} \frac{\delta}{1-\delta} \right)^{n-1} \left(\delta e + \sum_{r \geq 2} r(\delta e)^{r-1} \right), \quad (33)$$

which concludes the proof. \square

2.2. The small parameter of the cluster expansion

In order to identify a small parameter we set

$$E_{kl}(x-y) := J(x-y) - \bar{J}(k, l).$$

The expansion (26) suggests that the error depends on both β and

$$\sup_{k, l \in \bar{\Lambda}_M} \sup_{x \in C_k, y \in C_l} |E_{kl}(x-y)|.$$

This quantity can be evaluated more explicitly for the special choice of interaction potential defined in (4).

Lemma 2.2. *Assume that J satisfies (3)–(4) then the coarse-grained interaction potential \bar{J} at the coarse-graining level q approximates the potential J with the error*

$$|J(x-y) - \bar{J}(k, l)| \leq 2 \frac{q}{L^{d+1}} \sup_{\substack{x' \in C_k, \\ y' \in C_l}} \|\nabla V(x' - y')\| \quad (34)$$

for both $l \neq k$ and $l = k$.

Proof. Using the properties of the potential V , we have

$$V(z) = V(z') + (z - z') \cdot \nabla V(z') + o(\|z - z'\|).$$

Using the definition of J , (4) and setting $z = x - y$ and $z' = x' - y'$, where $x, x' \in C_k$ and $y, y' \in C_l$, we have

$$\begin{aligned} J(x-y) &= \frac{1}{q^{2d}} \sum_{x' \in C_k} \sum_{y' \in C_l} J(x' - y') \\ &+ \frac{1}{q^{2d}} \sum_{x' \in C_k} \sum_{y' \in C_l} \frac{n}{L^d L} ((x-y) - (x'-y')) \cdot \nabla V\left(\frac{n}{L}(x'-y')\right) \\ &+ \frac{1}{q^{2d}} \sum_{x' \in C_k} \sum_{y' \in C_l} o\left(\left\| \frac{n}{L}((x-y) - (x'-y')) \right\|\right), \end{aligned}$$

and using the estimate $\|(x - y) - (x' - y')\| \leq \|x - x'\| + \|y - y'\| \leq 2 \max\{\text{diam}(C_k)\} \sim \frac{2}{m}$ we obtain (34) in both cases $l \neq k$ and $l = k$. \square

Our goal is to construct higher-order corrections to the Hamiltonian $\bar{H}_M^{(0)}$ based on the expansion (30). First we check the condition (29) and identify the parameter δ . From the point of view of the cluster expansion theory we are in the high temperature, small activity regime because of the *a priori* estimate given by Lemma 2.2 on the coarse-grained approximation. Hence we expect the small parameter δ to depend on the characteristics of the coarse-graining, in particular on the parameters q and L .

Lemma 2.3 (Identification of the small parameter δ). *For ζ defined in (28), condition (29) holds with*

$$\delta \sim \beta \sup_{k \in \bar{\Lambda}_M} \sum_{l: l \neq k} \left| \sum_{x \in C_k, y \in C_l} E_{kl}(x - y) \right|.$$

In the particular case of Lemma 2.2 we obtain $\delta \sim \beta \frac{q^{d+1}}{L} \|\nabla V\|_\infty$, where q is the level of coarse-graining and L is the range of the interaction of the particles on the microscopic lattice.

Proof. To check the condition (29) we follow the analysis proposed in [22]. We start with the $r = 1$ case

$$\sup_{k \in \bar{\Lambda}_M} \sum_{\substack{R \in \mathcal{R}, R \supset \{k\} \\ |R|=1}} |\zeta(R)| = \sup_{k \in \bar{\Lambda}_M} \int |e^{-\beta \Delta_{kk} J(\sigma)} - 1| \tilde{\rho}_k(d\sigma) \sim \mathcal{O} \left(\left(\frac{q}{L} \right)^d \beta \frac{q^{d+1}}{L} \|\nabla V\|_\infty \right),$$

where we have used the fact that $|e^a - 1| \leq |a|e^{|a|}$ and that $e^{\beta|\Delta_{kk} J(\sigma)|} \sim \mathcal{O}(1)$. For the general case $r \geq 2$ we have

$$\begin{aligned} \sup_{k \in \bar{\Lambda}_M} \sum_{\substack{R \in \mathcal{R}, R \supset \{k\} \\ |R|=r}} \left| \int \sum_{g \in G_R} \prod_{\{l_1, l_2\} \in g} f_{l_1 l_2}(\sigma) \prod_{\{k\} \in R} \tilde{\rho}_k(d\sigma) \right| = \\ \sup_{k \in \bar{\Lambda}_M} \frac{1}{(r-1)!} \sum_{\substack{k_1 = k, k_2, \dots, k_r \\ k_i \neq k_j, i \neq j}} \left| \int \sum_{g \in G_r} \prod_{\{i, j\} \in g} (e^{-\beta \Delta_{k_i k_j} J(\sigma)} - 1) \prod_{i=1}^r \tilde{\rho}_{k_i}(d\sigma) \right|, \end{aligned} \tag{35}$$

where we have assumed $R = \{k_1, \dots, k_r\}$. We divide by $(r - 1)!$ because we can relabel k_2, \dots, k_r . Note that by G_r we denote the set of all connected graphs in $\{1, 2, \dots, r\}$ which may have self-loops $\{j, j\}$ for any combination of $j = 1, 2, \dots, r$ (from now on we simply call them loops instead of self-loops). In the sequel we need the following estimate: due to Lemma 2.2 we have that for a fixed $k \in \bar{\Lambda}_M$ which interacts with $(\frac{L}{q})^d$ many coarse cells,

$$\sum_{l: l \neq k} |\Delta_{kl} J(\sigma)| \sim \left(\frac{L}{q} \right)^d q^{2d} \frac{1}{L^d} \mathcal{O} \left(\frac{q}{L} \right) \sim \mathcal{O} \left(\frac{q^{d+1}}{L} \|\nabla V\|_\infty \right),$$

which implies that

$$\sup_{k \in \bar{\Lambda}_M} \sum_{l: l \neq k} |\Delta_{kl} J(\sigma)| \leq C, \tag{36}$$

for every σ and with $C \sim \sup_{k \in \bar{\Lambda}_M} \sum_{l: l \neq k} \left| \sum_{x \in C_k, y \in C_l} E_{kl}(x - y) \right|$ or in the special case of J given in terms of V $C \sim \frac{q^{d+1}}{L} \|\nabla V\|_\infty$.

Now we denote by G_r^0 the set of graphs g^0 without loops and we describe by $g \sim g^0$ the corresponding graph g with loops such that $g = g^0$ if we remove all loops $\{j, j\}$ from g . We have

$$\sum_{g \in G_r} \prod_{\{i,j\} \in g} (e^{-\beta \Delta_{k_i k_j} J(\sigma)} - 1) = \sum_{g^0 \in G_r^0} \prod_{\{i,j\} \in g^0} (e^{-\beta \Delta_{k_i k_j} J(\sigma)} - 1) \left(1 + \sum_{g \sim g^0} \prod_{\{j,j\} \in g/g^0} (e^{-\beta \Delta_{k_j k_j} J(\sigma)} - 1) \right).$$

We observe that for any $g^0 \in G_r^0$ with $|\text{supp}(g^0)| = r$ the terms in the parenthesis are bounded by 2^r , since for every $g \sim g^0$ we have that $\prod_{\{i,j\} \in g/g^0} \beta |\Delta_{k_i k_j} J(\sigma)| \leq 1$ (after the choice we will make at the end of the proof) and the summation $\sum_{g \sim g^0}$ has $\sum_{k=1}^r \binom{r}{k} = 2^r - 1$ many terms. Then, for the sum $\sum_{g^0 \in G_r^0} \prod_{\{i,j\} \in g^0} (e^{-\beta \Delta_{k_i k_j} J(\sigma)} - 1)$ we can use the tree-graph equality (e.g., [22], p. 5, Thm. 2) to estimate the sum over connected graphs with respect to the sum over the corresponding maximal spanning trees. From [22], Theorem 2, we have that

$$\begin{aligned} \sum_{g^0 \in G_r^0} \prod_{\{i,j\} \in g^0} (e^{-\beta \Delta_{k_i k_j} J(\sigma)} - 1) &= \sum_{\tau^0 \in \mathcal{T}_r^0} \prod_{\{i,j\} \in \tau^0} (-\beta \Delta_{k_i k_j} J(\sigma)) \int_0^1 dt_1 \dots \int_0^1 dt_{r-1} \\ &\times \sum_{\substack{X_1, \dots, X_{r-1} \\ \text{comp. with } \tau^0}} t_1^{b_1-1} \dots t_{r-1}^{b_{r-1}-1} e^{-W(X_1, \dots, X_{r-1}, t_1, \dots, t_{r-1})}, \end{aligned} \tag{37}$$

where g^0 is a connected graph in $\{1, 2, \dots, r\}$ without loops and τ^0 is the corresponding tree graph. The summation is over all sequences X_1, \dots, X_{r-1} of increasing subsets of $\{1, 2, \dots, r\}$ such that $X_1 = 1$, $X_n \subset X_{n+1}$ and $|X_n| = n$ that are compatible with τ^0 . A sequence X_1, \dots, X_{r-1} is compatible with a given tree τ^0 if, for all $n = 1, \dots, r-1$, X_n contains exactly $n-1$ links of τ^0 . We also say that a link $\{i, j\}$ crosses X_n if $i \in X_n$ and $j \notin X_n$ or *vice versa*; then b_n is the number of links of τ^0 which cross X_n , and

$$W(X_1, \dots, X_{r-1}, t_1, \dots, t_{r-1}) = \sum_{1 \leq i < j \leq r} t_1(\{i, j\}) \dots t_{r-1}(\{i, j\}) \beta \Delta_{k_i k_j} J(\sigma),$$

where

$$t_n(\{i, j\}) = \begin{cases} t_n \in [0, 1], & \text{if } \{i, j\} \text{ crosses } X_n \\ 1, & \text{otherwise} \end{cases}.$$

Then from (36) we obtain that

$$W(X_1, \dots, X_{r-1}, t_1, \dots, t_{r-1}) \leq \sum_{1 \leq k < l \leq r} |\beta \Delta_{kl} J(\sigma)| \leq r\beta \sup_{k \in \bar{A}_M} \sum_{l=1, l \neq k}^r |\Delta_{kl} J(\sigma)| \sim r\beta C,$$

where $C \sim \mathcal{O}(\frac{d+1}{L} \|\nabla V\|_\infty)$. Thus, uniformly in $X_1, \dots, X_{r-1}, t_1, \dots, t_{r-1}$ we have that

$$e^{-W(X_1, \dots, X_{r-1}, t_1, \dots, t_{r-1})} \leq e^{r\beta C}.$$

Furthermore, from [22], Lemma 4, we have that

$$\int_0^1 dt_1 \dots \int_0^1 dt_{r-1} \sum_{\substack{X_1, \dots, X_{r-1} \\ \text{comp. with } \tau^0}} t_1^{b_1-1} \dots t_{r-1}^{b_{r-1}-1} = 1.$$

This leads to the following tree-graph inequality for the case of graphs with loops

$$\begin{aligned} \left| \sum_{g \in G_r} \prod_{\{i,j\} \in g} (e^{-\beta \Delta_{k_i k_j} J(\sigma)} - 1) \right| &\leq 2^r \left| \sum_{g^0 \in G_r^0} \prod_{\{i,j\} \in g^0} (e^{-\beta \Delta_{k_i k_j} J(\sigma)} - 1) \right| \\ &\leq 2^r e^{r\beta C} \beta \sum_{\tau^0 \in \mathcal{T}_r^0} \prod_{\{i,j\} \in \tau^0} |\Delta_{k_i k_j} J(\sigma)|. \end{aligned}$$

From (36) we have that for any tree τ^0

$$\sup_{k \in \bar{\Lambda}_M} \sum_{\substack{k_1=k, k_2, \dots, k_r \\ k_i \neq k_j, i \neq j}} \prod_{\{i,j\} \in \tau^0} |\Delta_{k_i k_j} J(\sigma)| \leq C^{r-1}.$$

Using the Cayley formula $\sum_{\tau^0 \in \mathcal{T}_r^0} 1 = r^{r-2}$, we conclude that

$$\sup_{k \in \bar{\Lambda}_M} \sum_{\substack{R \in \mathcal{R}, R \supset \{k\} \\ |R|=r}} |\zeta(R)| \leq 2^r \frac{e^{r\beta C}}{(r-1)!} r^{r-2} (\beta C)^{r-1}.$$

Thus, we require

$$2\beta C e^{\beta C+1} \leq \delta,$$

so it suffices to consider $\delta \sim \beta \frac{q^{d+1}}{L} \|\nabla V\|_\infty$. □

2.3. Effective Hamiltonians and specific entropy estimates

In this section we use the results of Lemmas 2.1 and 2.3 and expand the Hamiltonian $\bar{H}_M(\eta)$ around $\bar{H}_M^{(0)}$ in powers of δ by truncating the series (30) at the levels of the number of interacting clusters and of the length of the clusters. We obtain the series

$$\bar{H}_M(\eta) = \bar{H}_M^{(0)}(\eta) + \bar{H}_M^{(1)}(\eta) + \dots + \bar{H}_M^{(p)}(\eta) + MO(\delta^{p+1}),$$

and define the corresponding Gibbs measures by

$$\bar{\mu}_{M,\beta}^{(p)}(d\eta) = \frac{1}{\bar{Z}_M^{(p)}} e^{-\beta(\bar{H}_M^{(0)}(\eta) + \dots + \bar{H}_M^{(p)}(\eta))} \bar{P}_M(d\eta), \tag{38}$$

where $p = 1, 2, \dots$ and $\bar{Z}_M^{(p)}$ is the corresponding partition function. In the following proposition we derive explicit formulas up to the order $\mathcal{O}(\delta^3)$ and state the corresponding error estimates for the relative entropy. Higher order corrections can be computed in a straightforward although tedious way. For the sake of simplicity we restrict ourselves to the case $d = 1$, however similar computations and results are derived in any dimension. To make the cluster summations more specific we also denote by $B_r(k)$ the ball centered at k of radius r for a given metric on the lattice. Furthermore, by $l \in B_r^+(k)$ we mean that $l \in B_r(k) \cap \{l > k\}$.

Proposition 2.4 (Corrections to $\bar{H}_M^{(0)}$). *For $\bar{H}_M^{(0)}$ we have the error estimate*

$$\bar{H}_M(\eta) = \bar{H}_M^{(0)}(\eta) + M\mathcal{O}(\delta). \tag{39}$$

Furthermore, the first and second order corrections to the Hamiltonian $\bar{H}_M^{(0)}$ are given by

$$\begin{aligned} \bar{H}_M(\eta) &= \bar{H}_M^{(0)}(\eta) + \bar{H}_M^{(1)}(\eta) + M\mathcal{O}(\delta^2) \\ \bar{H}_M(\eta) &= \bar{H}_M^{(0)}(\eta) + \bar{H}_M^{(1)}(\eta) + \bar{H}_M^{(2)}(\eta) + M\mathcal{O}(\delta^3), \end{aligned} \tag{40}$$

where

$$\begin{aligned} \bar{H}_M^{(1)}(\eta) &= -\frac{1}{\beta} \sum_{k \in \bar{\Lambda}_M} \int f_{kk}(\sigma) \tilde{\rho}_k(d\sigma) \\ &\quad - \frac{1}{\beta} \sum_k \sum_{l \in B_{\frac{L}{q}}^+(k)} \int (f_{kl} + f_{kl}f_{kk} + f_{kl}f_{ll} + f_{kl}f_{kk}f_{ll}) \tilde{\rho}_k(d\sigma) \tilde{\rho}_l(d\sigma) \end{aligned} \quad (41)$$

and

$$\begin{aligned} \bar{H}_M^{(2)}(\eta) &= \frac{1}{2\beta} \sum_{k \in \bar{\Lambda}_M} \left(\int f_{kk}(\sigma) \tilde{\rho}_k(d\sigma) \right)^2 \\ &\quad + \frac{1}{2\beta} \sum_k \sum_{l \in B_{\frac{L}{q}}^+(k)} \int f_{kk}(\sigma) \tilde{\rho}_k(d\sigma) \int f_{kl}(\sigma) \tilde{\rho}_k(d\sigma) \tilde{\rho}_l(d\sigma) \\ &\quad + \frac{1}{2\beta} \sum_k \sum_{l_1 \in B_{\frac{L}{q}}^+(k)} \sum_{l_2 \in B_{\frac{L}{q}}^+(k)} \int (f_{kl_1} + f_{kk}f_{kl_1} + f_{kl_1}f_{l_1l_1} + f_{kk}f_{kl_1}f_{l_1l_1}) \tilde{\rho}_k(d\sigma) \tilde{\rho}_{l_1}(d\sigma) \\ &\quad \times \int (f_{kl_2} + f_{kk}f_{kl_2} + f_{kl_2}f_{l_2l_2} + f_{kk}f_{kl_2}f_{l_2l_2}) \tilde{\rho}_k(d\sigma) \tilde{\rho}_{l_2}(d\sigma) \\ &\quad + \frac{1}{\beta} \sum_k \sum_{l_1 \in B_{\frac{L}{q}}^+(k)} \sum_{l_2 \in B_{\frac{L}{q}}(l_1) \cup B_{\frac{L}{q}}^+(k)} \int (f_{kl_1}f_{l_1l_2} + f_{kl_1}f_{kl_2} + f_{kl_2}f_{l_1l_2} + [\dots]) \tilde{\rho}_k(d\sigma) \tilde{\rho}_{l_1}(d\sigma) \tilde{\rho}_{l_2}(d\sigma) \\ &= I_1 + I_2 + I_3 + I_4, \end{aligned} \quad (42)$$

where $[\dots]$ means the previous three terms with all possible combinations of loops. The corresponding relative entropy error is

$$\frac{1}{N} \mathcal{R} \left(\bar{\mu}_{M,\beta}^{(p)} | \mu_{N,\beta} \circ \mathbf{F}^{-1} \right) \sim \mathcal{O} \left(\frac{1}{q} \delta^{p+1} \right).$$

Proof. The error in (39) is given by (33). To get an error $M\mathcal{O}(\delta^2)$ we need to include the terms $n = 1, r = 1, 2$, i.e.,

$$\bar{H}_M^{(1)}(\eta) = - \sum_{R: |R|=1,2} \zeta(R).$$

Similarly, for the $M\mathcal{O}(\delta^3)$ error we need to include (41) together with the terms $n = 2, r = 1, 2$ and $n = 1, r = 3$

$$\bar{H}_M^{(2)}(\eta) = -\frac{1}{2} \sum_{\substack{R_1, R_2 \\ |R_i|=1,2, i=1,2}} \phi(R_1, R_2) \zeta(R_1) \zeta(R_2) - \sum_{R: |R|=3} \zeta(R),$$

where $\phi(R_1, R_2) = 0$ if $R_1 \cap R_2 = \emptyset$ and -1 if $R_1 \cap R_2 \neq \emptyset$, which occurs when $R_1 = R_2 = \{k\}$ for $k \in \bar{\Lambda}_M$, in the case of $|R_i| = 1, i = 1, 2$. If $|R_i| = 2, i = 1, 2$ we have that $R_1 = \{k, l_1\}$ and $R_2 = \{k, l_2\}$. For the relative

entropy error for $\bar{H}_M^{(p)}$ with $p = 0, 1, \dots$, we have

$$\begin{aligned} \mathcal{R} \left(\bar{\mu}_{M,\beta}^{(p)} \mid \mu_{N,\beta} \circ \mathbf{F}^{-1} \right) &= \frac{1}{N} \int \log \frac{\frac{1}{\bar{Z}_M^{(p)}} e^{-\beta \bar{H}_M} \rho(\eta)}{\frac{1}{Z_N} \int_{\{\mathbf{F}(\sigma)=\eta\}} e^{-\beta H_N(\sigma)} P_N(d\sigma)} \bar{\mu}_{M,\beta}^{(p)}(d\eta) \\ &= \frac{1}{N} \int \log \frac{\frac{1}{\bar{Z}_M^{(p)}} e^{-\beta \bar{H}_M} \rho(\eta)}{\frac{1}{Z_M} e^{-\beta \bar{H}_M} \rho(\eta)} \bar{\mu}_{M,\beta}^{(p)}(d\eta) \\ &= \frac{1}{N} \log \frac{\bar{Z}_M}{\bar{Z}_M^{(p)}} + \frac{1}{N} \mathbb{E}_{\bar{\mu}_{M,\beta}^{(p)}} \left[\beta \left(\bar{H}_M - \bar{H}_M^{(p)} \right) \right], \end{aligned} \tag{43}$$

where for the partition functions we have

$$\bar{Z}_M = \sum_{\eta} e^{-\beta \bar{H}_M} \rho(\eta) = \bar{Z}_M^{(p)} \sum_{\eta} \frac{1}{\bar{Z}_M^{(p)}} e^{-\beta \bar{H}_M^{(p)}} e^{-\beta(\bar{H}_M - \bar{H}_M^{(p)})} \rho(\eta) = \bar{Z}_M^{(p)} \mathbb{E}_{\bar{\mu}_{M,\beta}^{(p)}} \left[e^{-\beta(\bar{H}_M - \bar{H}_M^{(p)})} \right]. \tag{44}$$

Thus, for the different choices of p in $\bar{H}_M^{(p)}$, given the fact that the estimates (39) and (40) are uniform in η we get that

$$\log \frac{\bar{Z}_M}{\bar{Z}_M^{(p)}} \sim \log \mathbb{E}_{\bar{\mu}_{M,\beta}^{(p)}} \left[e^{M\mathcal{O}(\delta^{p+1})} \right] \sim M\mathcal{O}(\delta^{p+1}), \quad \text{and} \quad \mathbb{E}_{\bar{\mu}_{M,\beta}^{(p)}} \left[\bar{H}_M - \bar{H}_M^{(p)} \right] \sim M\mathcal{O}(\delta^{p+1}),$$

where $p = 0, 1, 2$. □

3. NUMERICAL SCHEMES FOR COARSE-GRAINING AND A *POSTERIORI* ESTIMATES

We conclude the proof of Theorem 1.4 by giving explicit formulas for the correction terms (41) and (42) and estimating them with respect to the small parameter ϵ . The corrections consist of combinations of $f_{kl} = e^{-\beta \Delta_{kl} J(\sigma)} - 1$ for all k and l . Since the exponent $-\beta \Delta_{kl} J(\sigma)$ in (26) is small we have that

$$e^{-\beta \Delta_{kl} J(\sigma)} - 1 = \sum_{p=1}^{\infty} \frac{1}{p!} (-\beta \Delta_{kl} J(\sigma))^p \quad \text{with} \quad \Delta_{kl} J(\sigma) \sim \mathcal{O} \left(q^{2d} \frac{q}{L^{d+1}} \|\nabla V\|_{\infty} \right). \tag{45}$$

The key point of our calculations relies on the cancellation

$$\int \Delta_{kl} J(\sigma) \tilde{\rho}_k(d\sigma) \tilde{\rho}_l(d\sigma) = 0, \tag{46}$$

following from the definition of $\bar{H}_M^{(0)}$ in (17).

In order to keep expressions simple we express our results in terms of the variables $\alpha(k)$, the number of spins $\sigma(x) = 1$ in the coarse cell C_k , and $\omega(k) = q - \alpha(k)$. Next we define the following quantities for distinct x, y, z

and w

$$E_1(\alpha) := \mathbb{E}[\sigma(x)|\alpha] = \frac{2\alpha - q}{q}, \tag{47}$$

$$E_2(\alpha) := \mathbb{E}[\sigma(x)\sigma(y)|\alpha] = \frac{\alpha(\alpha - 1) - 2\alpha\omega + \omega(\omega - 1)}{q(q - 1)}, \tag{48}$$

$$\begin{aligned} E_3(\alpha) &:= \mathbb{E}[\sigma(x)\sigma(y)\sigma(z)|\alpha] \\ &= \frac{\alpha(\alpha - 1)(\alpha - 2) - 3\alpha(\alpha - 1)\omega + 3\alpha(\omega - 1)\omega - (\omega - 2)(\omega - 1)\omega}{q(q - 1)(q - 2)}, \end{aligned} \tag{49}$$

$$\begin{aligned} E_4(\alpha) &:= \mathbb{E}[\sigma(x)\sigma(y)\sigma(z)\sigma(w)|\alpha] \\ &= \frac{\alpha(\alpha - 1)(\alpha - 2)(\alpha - 3) - 4\alpha(\alpha - 1)(\alpha - 2)\omega}{q(q - 1)(q - 2)(q - 3)} \\ &\quad + \frac{6\alpha(\alpha - 1)(\omega - 1)\omega - 4\alpha(\omega - 2)(\omega - 1)\omega + \omega(\omega - 1)(\omega - 2)(\omega - 3)}{q(q - 1)(q - 2)(q - 3)}, \end{aligned} \tag{50}$$

where all spin sites in the expressions are different. Note that all terms are of order $\mathcal{O}(1)$. Furthermore, we introduce the notation

$$j_{kl}^1 := \sum_{\substack{x \in C_k \\ y \in C_l}} (J(x - y) - \bar{J}(k, l))^2 \tag{51}$$

$$j_{kl}^2 := \sum_{\substack{x \in C_k \\ y, z \in C_l}} (J(x - y) - \bar{J}(k, l))(J(x - z) - \bar{J}(k, l)) \tag{52}$$

$$j_{k_1 k_2 k_3}^2 := \sum_{\substack{x \in C_{k_1} \\ y \in C_{k_2}, z \in C_{k_3}}} (J(x - y) - \bar{J}(k_1, k_2))(J(y - z) - \bar{J}(k_2, k_3)). \tag{53}$$

If $k_1 = k_2$ then we also impose that for $x, y \in C_{k_1}$ we have $y \neq x$.

Note that these quantities have various symmetries following from translational invariance of $J(x - y)$; for example, $j_{lk}^2 = j_{kl}^2$ or $j_{kl}^1 = \tilde{j}_{k-l}^1 = \tilde{j}_{l-k}^1$ for some function \tilde{j}^1 and similarly j_{kl}^2 depends also only on $k - l$, moreover $j_{k_1 k_2 k_3}^2 = \tilde{j}_{k_1 - k_2, k_3 - k_2}^2$.

Based on (18) we get first estimates in terms of q and L

$$j_{kl}^1 \sim \left(\frac{q}{L}\right)^{2d+2}, \quad j_{kl}^2 \sim q^{3d} \left(\frac{q}{L^{d+1}}\right)^2 \sim q^d \left(\frac{q}{L}\right)^{2d+2}, \quad j_{k_1 k_2 k_3}^2 \sim q^d \left(\frac{q}{L}\right)^{2d+2}.$$

Proof of Theorem 1.4. Starting from the formulas for $\bar{H}_M^{(1)}$ and $\bar{H}_M^{(2)}$ in Proposition 2.4 we expand the factors f_{kl} and re-estimate the corrections using the cancellations in (46). Recalling that $\epsilon \sim \frac{q}{L}\beta\|\nabla V\|_\infty$, we have

$$\begin{aligned} \text{(i)} \quad & \sum_{k \in \bar{\Lambda}_M} \int f_{kk}(\sigma) \tilde{\rho}_k(d\sigma) \sim M \left(q^{2d} \frac{q}{L^{d+1}} \beta \|\nabla V\|_\infty \right)^2 \sim M q^{2d} \left(\frac{q}{L}\right)^{2d} \epsilon^2 \\ \text{(ii)} \quad & \sum_k \sum_{l \in B_{\frac{L}{q}}(k)} \int (f_{kl} + f_{kl} f_{kk} + f_{kl} f_{ll}) \tilde{\rho}_k(d\sigma) \tilde{\rho}_l(d\sigma) \sim M \left(\frac{L}{q}\right)^d \left(q^{2d} \frac{q}{L^{d+1}} \beta \|\nabla V\|_\infty \right)^2 \sim M q^{2d} \left(\frac{q}{L}\right)^d \epsilon^2 \end{aligned}$$

where in the estimate we included $\|\nabla V\|_\infty$ and β into the constant since here we want to focus on the L and q dependence. Note also that the term involving $f_{kl} f_{kk} f_{ll}$ is of a higher order.

Therefore it suffices to include only the first two contributions which we also compute explicitly. With a slight abuse of notation we still denote the new terms by $\bar{H}_M^{(1)}$. Collecting all the terms gives us

$$\begin{aligned}
 -\bar{H}_M^{(1)} &= \beta \sum_k \int \frac{1}{8} (\Delta_{kk} J(\sigma))^2 \tilde{\rho}_k(d\sigma) + \beta \sum_{k < l} \int \frac{1}{2} (\Delta_{kl} J(\sigma))^2 \tilde{\rho}_k(d\sigma) \tilde{\rho}_l(d\sigma) \\
 &\quad + \beta \sum_{k < l} \int \frac{1}{2} \Delta_{kk} J(\sigma) \Delta_{kl} J(\sigma) \tilde{\rho}_k(d\sigma) \tilde{\rho}_l(d\sigma) + \beta \sum_{k < l} \int \Delta_{kl} J(\sigma) \frac{1}{2} \Delta_{ll} J(\sigma) \tilde{\rho}_k(d\sigma) \tilde{\rho}_l(d\sigma).
 \end{aligned}$$

Explicit calculations of these integrals are detailed in Appendix A, yielding

$$\begin{aligned}
 -\bar{H}_M^{(1)} &= \frac{\beta}{8} \sum_k 4j_{kk}^2 [-E_4(\alpha(k)) + E_2(\alpha(k))] + 2j_{kk}^1 [E_4(\alpha(k)) + 1 - 2E_2(\alpha(k))] \\
 &\quad + \frac{\beta}{2} \sum_{k < l} j_{kl}^1 [E_2(\alpha(k))E_2(\alpha(l)) - E_2(\alpha(l)) - E_2(\alpha(k)) + 1] \\
 &\quad + j_{kl}^2 [-2E_2(\alpha(k))E_2(\alpha(l)) + E_2(\alpha(k)) + E_2(\alpha(l))] \\
 &\quad + \frac{\beta}{2} \sum_{k < l} 2j_{kkl}^2 [-E_3(\alpha(k))E_1(\alpha(l)) + E_1(\alpha(k))E_1(\alpha(l))] \\
 &\quad + \frac{\beta}{2} \sum_{k < l} 2j_{llk}^2 [-E_3(\alpha(l))E_1(\alpha(k)) + E_1(\alpha(l))E_1(\alpha(k))].
 \end{aligned}$$

Since $j_{kkl}^2 = j_{llk}^2$ this expression further simplifies to

$$\begin{aligned}
 -\bar{H}_M^{(1)} &= \frac{\beta}{8} \sum_k 4j_{kk}^2 [-E_4(\alpha(k)) + E_2(\alpha(k))] + 2j_{kk}^1 [E_4(\alpha(k)) + 1 - 2E_2(\alpha(k))] \\
 &\quad + \frac{\beta}{2} \sum_{k < l} j_{kl}^1 [E_2(\alpha(k))E_2(\alpha(l)) - E_2(\alpha(l)) - E_2(\alpha(k)) + 1] \\
 &\quad + j_{kl}^2 [-2E_2(\alpha(k))E_2(\alpha(l)) + E_2(\alpha(k)) + E_2(\alpha(l))] \\
 &\quad + \frac{\beta}{2} \sum_{k, l \neq k} j_{kkl}^2 [-E_3(\alpha(k))E_1(\alpha(l)) + 2E_1(\alpha(k))E_1(\alpha(l)) - E_3(\alpha(l))E_1(\alpha(k))].
 \end{aligned}$$

By the estimates on the terms j_{kl}^2 and the fact that the terms $E_i(\alpha)$ are of order one, and by counting the summations we obtain

$$\begin{aligned}
 \bar{H}_M^{(1)} &\sim M \frac{1}{\beta} q^{3d} \left(\frac{q}{L^{d+1}} \beta \|\nabla V\|_\infty \right)^2 + M \frac{1}{\beta} \left(\frac{L}{q} \right)^d q^{3d} \left(\frac{q}{L^{d+1}} \beta \|\nabla V\|_\infty \right)^2 \\
 &\sim M q^d \left(\frac{q}{L} \right)^{2d} \epsilon^2 + M q^d \left(\frac{L}{q} \right)^d \left(\frac{q}{L} \right)^{2d} \epsilon^2 \sim N \left(\frac{q}{L} \right)^d \epsilon^2,
 \end{aligned}$$

hence we have gained an extra q^d as compared with the previous estimate. In the same spirit we estimate the terms of the correction $\bar{H}_M^{(2)}$. From (42) we have

$$\begin{aligned}
 I_1 &\sim M \frac{1}{\beta} \left(q^{2d} \frac{q}{L^{d+1}} \beta \|\nabla V\|_\infty \right)^4, & I_2 &\sim M \frac{1}{\beta} \left(\frac{L}{q} \right)^d \left(q^{2d} \frac{q}{L^{d+1}} \beta \|\nabla V\|_\infty \right)^4, \\
 I_3 &\sim M \frac{1}{\beta} \left(\frac{L}{q} \right)^{2d} \left(q^{2d} \frac{q}{L^{d+1}} \beta \|\nabla V\|_\infty \right)^4, & I_4 &\sim M \frac{1}{\beta} \left(\frac{L}{q} \right)^{2d} \left(q^{2d} \frac{q}{L^{d+1}} \beta \|\nabla V\|_\infty \right)^2,
 \end{aligned}$$

where in the term I_4 we have neglected the higher order contribution of the terms [...] in (42). Obviously, for $\bar{H}_M^{(2)}$ it suffices to consider only the term I_4 . Abusing slightly the notation we call this term $\bar{H}_M^{(2)}$ and we calculate it explicitly

$$\begin{aligned} \bar{H}_M^{(2)} = & \beta \sum_{k_1} \sum_{k_2 > k_1} \sum_{k_3 > k_2} \int \Delta_{k_1 k_2} J(\sigma) \Delta_{k_2 k_3} J(\sigma) \tilde{\rho}_{k_1}(d\sigma) \tilde{\rho}_{k_2}(d\sigma) \tilde{\rho}_{k_3}(d\sigma) \\ & + \int \Delta_{k_2 k_3} J(\sigma) \Delta_{k_3 k_1} J(\sigma) \tilde{\rho}_{k_2}(d\sigma) \tilde{\rho}_{k_3}(d\sigma) \tilde{\rho}_{k_1}(d\sigma) + \int \Delta_{k_3 k_1} J(\sigma) \Delta_{k_1 k_2} J(\sigma) \tilde{\rho}_{k_3}(d\sigma) \tilde{\rho}_{k_1}(d\sigma) \tilde{\rho}_{k_2}(d\sigma). \end{aligned}$$

Similarly, as for the previous term, we evaluate the integrals in terms of E_k and obtain

$$\begin{aligned} \bar{H}_M^{(2)} = & \beta \sum_{k_1} \sum_{k_2 > k_1} \sum_{k_3 > k_2} j_{k_1 k_2 k_3}^2 (-E_1(\alpha(k_1)) E_2(\alpha(k_2)) E_1(\alpha(k_3)) + E_1(\alpha(k_1)) E_1(\alpha(k_3))) \\ & + j_{k_2 k_3 k_1}^2 (-E_1(\alpha(k_2)) E_2(\alpha(k_3)) E_1(\alpha(k_1)) + E_1(\alpha(k_2)) E_1(\alpha(k_1))) \\ & + j_{k_3 k_1 k_2}^2 (-E_1(\alpha(k_3)) E_2(\alpha(k_1)) E_1(\alpha(k_2)) + E_1(\alpha(k_3)) E_1(\alpha(k_2))). \end{aligned}$$

Using the estimates above we obtain the order of the correction

$$\bar{H}_M^{(2)} \sim M \frac{1}{\beta} \left(\frac{L}{q}\right)^{2d} q^{3d} \left(\frac{q}{L^{d+1}} \beta \|\nabla V\|_\infty\right)^2 \sim M q^d \epsilon^2.$$

Thus, from the above we deduce two results: the first one that due to the fact that the leading order terms that participate in the $\mathcal{O}(\delta)$ error vanish, the choice of the initial Hamiltonian $\bar{H}_M^{(0)}$ yields a second-order accurate approximation, in other words

$$\frac{\beta}{N} \left(\bar{H}_M - \bar{H}_M^{(0)}\right) \sim \mathcal{O}(\epsilon^2).$$

The second one is that to obtain a $\mathcal{O}(\epsilon^3)$ error, the corresponding correction term have to include both $\bar{H}_M^{(1)}$ and $\bar{H}_M^{(2)}$. In such a case we obtain

$$\frac{\beta}{N} \left(\bar{H}_M - \left(\bar{H}_M^{(0)} + \bar{H}_M^{(1)} + \bar{H}_M^{(2)}\right)\right) \sim \mathcal{O}(\epsilon^3).$$

In the same spirit we can derive error estimates for the additional terms in the expansion. □

In Section 4 we discuss the computational complexity of higher-order terms. Here we present only two schemes that are relevant for practical implementations.

Scheme 3.1 (2nd-order coarse-graining). *The 2nd-order coarse-graining algorithm has the following characteristics*

- (1) *Hamiltonian:* $\bar{H}_M^{(0)}$, given by (17).
- (2) *Gibbs measure:* $\bar{\mu}_{M,\beta}^{(0)}$, given by (38) for $p = 1$.
- (3) *Relative entropy error*

$$\mathcal{R} \left(\bar{\mu}_{M,\beta}^{(0)} \mid \mu_{N,\beta} \circ \mathbf{F}^{-1}\right) \sim \mathcal{O}(\epsilon^2).$$

Scheme 3.1 is the coarse-graining algorithm that has been extensively studied in [13–15, 17]. The novelty presented in this paper is the sharper error estimate that shows that the error is of the order $\mathcal{O}(\epsilon^2)$; this estimate readily follows from the cancellation due to (46) as is seen from the calculations above.

Scheme 3.2 (3rd-order coarse-graining). *We construct a higher-order Monte Carlo algorithm with the following characteristics*

(1) *Hamiltonian*: $\bar{H}_M^{(0)} + \bar{H}_M^{(1)} + \bar{H}_M^{(2)}$, where the corrections are

$$\begin{aligned}
 -\bar{H}_M^{(1)}(\eta) &= \frac{\beta}{8} \sum_k 4j_{kk}^2 [-E_4(\alpha(k)) + E_2(\alpha(k))] \\
 &\quad + 2j_{kk}^1 [E_4(\alpha(k)) + 1 - 2E_2(\alpha(k))] \\
 &\quad + \frac{\beta}{2} \sum_{k<l} j_{kl}^1 [E_2(\alpha(k))E_2(\alpha(l)) - E_2(\alpha(l)) - E_2(\alpha(k)) + 1] \\
 &\quad + j_{kl}^2 [-2E_2(\alpha(k))E_2(\alpha(l)) + E_2(\alpha(k)) + E_2(\alpha(l))] \\
 &\quad + \frac{\beta}{2} \sum_{k,l \neq k} j_{kl}^2 [-E_3(\alpha(k))E_1(\alpha(l)) + 2E_1(\alpha(k))E_1(\alpha(l)) - E_3(\alpha(l))E_1(\alpha(k))] \tag{54}
 \end{aligned}$$

and

$$\begin{aligned}
 \bar{H}_M^{(2)}(\eta) &= \beta \sum_{k_1} \sum_{k_2 > k_1} \sum_{k_3 > k_2} j_{k_1 k_2 k_3}^2 (-E_1(\alpha(k_1))E_2(\alpha(k_2))E_1(\alpha(k_3)) + E_1(\alpha(k_1))E_1(\alpha(k_3))) \\
 &\quad + j_{k_2 k_3 k_1}^2 (-E_1(\alpha(k_2))E_2(\alpha(k_3))E_1(\alpha(k_1)) + E_1(\alpha(k_2))E_1(\alpha(k_1))) \\
 &\quad + j_{k_3 k_1 k_2}^2 (-E_1(\alpha(k_3))E_2(\alpha(k_1))E_1(\alpha(k_2)) + E_1(\alpha(k_3))E_1(\alpha(k_2))). \tag{55}
 \end{aligned}$$

The terms E_i are defined in (47)–(50) and the quantities $j_{kl}^1, j_{kl}^2, j_{k_1 k_2 k_3}^2$ are defined in (51)–(53).

(2) *Gibbs measure* $\bar{\mu}_{M,\beta}^{(2)}(d\eta) = \frac{1}{\bar{Z}_M^{(2)}} e^{-(\bar{H}_M^{(0)} + \bar{H}_M^{(1)} + \bar{H}_M^{(2)})} \bar{P}_M(d\eta)$.

(3) *Relative entropy error*

$$\mathcal{R} \left(\bar{\mu}_{M,\beta}^{(2)} \mid \mu_{N,\beta} \circ \mathbf{F}^{-1} \right) \sim \mathcal{O}(\epsilon^3).$$

Remark 3.1. The estimates in Schemes 3.1 and 3.2 refer to the equilibrium Gibbs states of lattice systems. Non-equilibrium models such as the Arrhenius dynamics can be coarse-grained as shown in [14, 15]. In this case it is also possible to carry out a detailed error analysis between the exact microscopic and the approximating coarse-grained dynamics. In [13] an order $\mathcal{O}(\epsilon)$ estimate was proved for the specific relative entropy between microscopic and coarse-grained dynamics, while in [17] an improved $\mathcal{O}(\epsilon^2)$ estimate was shown in the weak topology. Both results are analogous to Scheme 3.1, as the invariant Gibbs measure corresponding to the coarse-grained Arrhenius dynamics studied there is (38).

We now turn our attention to the *a posteriori* error estimate in Theorem 1.6. The terms derived in Scheme 3.2 and the error estimate of Theorem 1.4 provide an explicit way of calculating the error made by the 2nd-order coarse-graining.

Proof of Theorem 1.6. As we have seen in Proposition 2.4, we can write the relative entropy error as the expectation with respect to the coarse-grained measure of some quantities, namely of the difference $\bar{H}_M - \bar{H}_M^{(p)}$ for any choice of K . On the other hand, in the Scheme 3.2 we have calculated explicitly the terms that contribute to the error of $\bar{H}_M^{(0)}$. We now conclude the proof by recalling the proof of Theorem 1.4 and relations (43) and (44). □

Using a Metropolis Monte Carlo sampler (see App. B), we can numerically compute the expectations in (43) and (44) and thus we can calculate the *a posteriori* error of $\mathcal{R}(\bar{\mu}_{M,\beta}^{(0)} \mid \mu_{N,\beta} \circ \mathbf{F}^{-1})$. Note also that as a consequence the quantity $\mathbb{E}_{\bar{\mu}_{M,\beta}^{(0)}} [R(\eta)]$ can be calculated on-the-fly using the coarse simulation only, at least up to the error $\mathcal{O}(\epsilon^3)$ according to Scheme 3.2.

3.1. Coarse-graining of external fields

To include an x -dependent external field in our analysis, we define an effective external field \bar{h} by

$$e^{-\beta\bar{h}(\eta(k))} = \int_{\{F(\sigma|_k)=\eta(k)\}} e^{-\beta\sum_{x\in C_k} h(x)\sigma(x)} \tilde{\rho}_k(d\sigma) \quad (56)$$

for every k . Then the effective full Hamiltonian (for both the interaction and the external field contribution) will be given by

$$e^{-\beta\bar{H}_M(\eta)} = \frac{1}{\bar{\rho}(\eta)} \int_{\{F(\sigma)=\eta\}} e^{-\beta H_N - \beta\sum_{x\in\Lambda} h(x)\sigma(x)} P_N(d\sigma)$$

and by perturbing around the coarse-grained Hamiltonian $\bar{H}_M^{(0)} + \sum_k \bar{h}(\eta(k))$ we obtain

$$\bar{H}_M(\eta) = \bar{H}_M(0) + \sum_k \bar{h}(\eta(k)) - \frac{1}{\beta} \log \frac{1}{\bar{\rho}(\eta)} \int_{\{F(\sigma)=\eta\}} e^{-\beta\Delta H} e^{\beta(\sum_k \bar{h}(\eta(k)) - \sum_{x\in\Lambda} h(x)\sigma(x))} P_N(d\sigma),$$

where $\Delta H = H_N - \bar{H}_M^{(0)}$. Since the part that involves the external field can be written in a product form, we obtain

$$\bar{H}_M(\eta) = \bar{H}_M^{(0)} + \sum_k \bar{h}(\eta(k)) - \frac{1}{\beta} \log \int e^{-\beta\Delta H} \prod_k \left(e^{\beta(\bar{h}(\eta(k)) - \sum_{x\in C_k} h(x)\sigma(x))} \tilde{\rho}_k(d\sigma) \right).$$

The new prior measure is normalized due to the definition (56) of the effective external field and therefore we can proceed with the cluster expansions as described before. Eventually, the only issue will be the analytic evaluation of the expected values $\mathbb{E}_h[\sigma(x)|\eta]$ with respect to the new prior measure due to the dependence on the external field. Furthermore, it is easy to see that in the case of a constant or slowly varying (at the same scale as the coarse cells) external field there is no extra contribution to the correction terms of the Hamiltonian.

4. COMPUTATIONAL ALGORITHMS AND NUMERICAL EXPERIMENTS

The 2nd-order approximation for the coarse-grained algorithm as described in Scheme 3.1 has been extensively studied in previous works, see, *e.g.*, [14, 15, 17], where it has been demonstrated that it performs well in certain regimes (*e.g.*, long-range interactions). In this section we present numerical experiments based on Scheme 3.2 and its comparison with the 2nd-order coarse-graining Scheme 3.1 for cases where the latter does not give satisfactory results.

Before we present numerical examples we briefly discuss the computational complexity of the approximations. As a simple measure of complexity we use the number of operations required for evaluating the Hamiltonian. Although the actual Monte Carlo step does not require evaluation of the full Hamiltonian the relative complexity with respect to the operation count of the full microscopic simulation $q = 1$ is properly reflected by this measure. Given a potential with the interaction radius L on a d -dimensional lattice with N sites, the number of operations for evaluation of the microscopic Hamiltonian H_N is $\mathcal{O}(NL^d)$. The count for the 2nd-order approximation $H_M^{(0)}$ on the coarse lattice with M sites and coarse-graining ratio q becomes $\mathcal{O}(ML^d/q^d)$. The 3rd-order approximation $H_M^{(1)}$ involves an additional summation over the interaction range and hence the operation count is $\mathcal{O}(ML^{2d}/q^{2d})$. In these estimates the ratio L/q is understood to be equal to one whenever $q \geq L$. In such a case the coarse interactions are reduced to the nearest-neighbor case. Thus the compression of the interaction kernel J in the corresponding approximations yields the speed-up of order $\mathcal{O}(q^{2d})$ for $H_M^{(0)}$ and $\mathcal{O}(q^{3d}/L^d)$ for $H_M^{(1)}$. We see that the 3rd-order approximation gives an improved error estimate at the same computational cost whenever $q = L$, in other words, whenever we can compress interactions to the nearest-neighbor potential.

We demonstrate the approximation properties in simulations of one-dimensional Ising-type spin systems. The one-dimensional system provides a suitable test bed since the exact (analytical) solutions are known for both the classical Ising system (*i.e.*, nearest-neighbor) and the mean-field model (Curie-Weiss model). We use the

exact solutions to ensure that the simulations are not influenced by finite-size effects. In all figures the exact solutions visually coincide with the fully resolved simulations, *i.e.*, $q = 1$. We computed error bars for statistical post-processing, however, they are not displayed in the figures due to their small relative size as compared to the scales of figures.

In the case of nearest-neighbor interactions the one-dimensional system does not exhibit phase transition. In fact, the exact solution is given by a well-known formula (see, *e.g.*, [18]), which we adopt to our choice of Hamiltonian with the constant nearest-neighbor ($L = 1$) interaction potential of strength J_0 . The equilibrium magnetization curve is then given by

$$m_\beta(h) = \frac{\sinh(\beta h)}{\sqrt{\sinh^2(\beta h) + e^{-2\beta J_0}}}. \quad (57)$$

On the other hand, for infinitely long attractive interactions there exists a 2nd-order phase transition and hysteresis behavior is observed according to the global mean-field theory for $\beta > \beta_c$ [9]. More explicitly, the mean-field (Curie-Weiss) model gives the magnetization curve as a solution of the non-linear equation

$$h = -J_0 m_\beta + \frac{1}{2\beta} \log \frac{1 + m_\beta}{1 - m_\beta}. \quad (58)$$

The Curie-Weiss model exhibits phase transition at the critical temperature given by $\beta_c J_0 = 1$ in the case of spins $\{-1, 1\}$ ($\beta_c J_0 = 4$ for spins $\{0, 1\}$).

The approximation of the hysteresis behavior in coarse-grained simulations provides a good test for derived coarse-graining schemes. It has been observed previously that hysteresis and critical behavior are not captured properly for short and intermediate range potentials [15]. Similar issues in predicting critical behavior were also observed in [21] for coarse-graining of complex fluids. There an artificial solidification effect was observed for higher levels of coarse-graining.

In the numerical tests presented here we demonstrate that the derived corrections improve this behavior even in the case of nearest-neighbor interactions or high coarse-graining ratio q . The sampling of the equilibrium measure is done by using microscopic and coarse-grained Metropolis dynamics discussed briefly in Appendix B. We compute isotherms similarly to natural parameter continuation, *i.e.*, we trace the magnetization m_β *vs.* external field h , first upon increasing the field h from low values and then decreasing it from high values. All simulations have been done with the fine lattice of the size $N = 512$. As derived in Section 2.2 the errors depend on the interplay of three parameters q , L and β . In examples presented here we investigate approximation properties for some regimes in this parameter space. In the computational examples we choose J to be constant on its support of the size $2L$, in particular $J(x) = \frac{J_0}{2L}$, for $|x| \leq L$ and 0 otherwise.

Test Case I: short-range interactions. We use the classical Ising model with nearest-neighbor interactions to show efficiency of the 3rd-order correction term. Figures 1 and 2 depicts simulations at two different temperatures $\beta = 2, 3$. The 2nd-order approximation exhibits hysteresis behavior for β above the critical value β_c for the Curie-Weiss model. It partly follows the magnetization curve predicted by the mean-field theory. Including the corrections into the effective Hamiltonian removes the hysteresis and in the case of smaller β gives even a reasonable approximation to the exact solution, which coincides with the case $q = 1$ at the scale of the figure.

Test Case II: intermediate-range interactions. In this case the system exhibits hysteresis behavior (and a 2nd-order phase transition) but the correct approximation of the transition between two states is not achieved by the 2nd-order effective Hamiltonian. In Figures 3 and 4 we present an example with extreme coarse-graining up to the interaction range, *i.e.*, $q = L = 8$, and beyond the interaction range $q = 32$. Note that this example is far from the $\epsilon \sim \frac{q}{L} \ll 1$ limit suggested by Lemma 2.1. However, as is the case in most asymptotics, computations perform well even for larger values of ϵ and especially in this case where the higher-order corrections are added.

Test Case III: long-range interactions. If the interaction range L is sufficiently large the behavior of the system is well-approximated by the mean-field solution and coarse-grained simulations will give good predictions already




FIGURE 1. Comparison of fully resolved $q = 1$ and coarse-grained $q = 8$ simulations. The interaction range is $L = 1$ and the inverse temperature is fixed at $\beta = 2$.




FIGURE 2. Comparison of fully resolved $q = 1$ and coarse-grained $q = 8$ simulations. The interaction range is $L = 1$ and the inverse temperature is fixed at $\beta = 3$.

in the 2nd-order approximation. This is observed in Figures 7 and 8 with coarse-graining to the range of interactions $q = L = 32$. However, coarse-graining beyond the interaction range ($q = 64$ in Fig. 8) shows that the 2nd and 3rd-order schemes give almost identical results. On the other hand the simulations in Figures 5




FIGURE 3. Comparison of fully resolved $q = 1$ and coarse-grained $q = 8$ simulations. The interaction range is $L = 8$ and the inverse temperature is fixed at $\beta = 2$.




FIGURE 4. Comparison of fully resolved $q = 1$ and coarse-grained $q = 32$ simulations. The interaction range is $L = 8$ and the inverse temperature is fixed at $\beta = 2$.




FIGURE 5. Comparison of fully resolved $q = 1$ and coarse-grained $q = 8$ simulations. The interaction range is $L = 8$ and the inverse temperature is fixed at $\beta = 2$.




FIGURE 6. Comparison of fully resolved $q = 1$ and coarse-grained $q = 8$ simulations in the “high temperature” regime. The interaction range is $L = 8$ and the inverse temperature is fixed at $\beta = 1$.

and 6 show two regimes where the 2nd-order approximation gives reasonable agreement with the microscopic magnetization curve.




FIGURE 7. Comparison of fully resolved $q = 1$ and coarse-grained $q = 32$ simulations. The interaction range is $L = 32$ and the inverse temperature is fixed at $\beta = 2$.




FIGURE 8. Comparison of fully resolved $q = 1$ and coarse-grained $q = 64$ simulations. The interaction range is $L = 32$ and the inverse temperature is fixed at $\beta = 2$.

APPENDIX A

In this section we present the detailed calculations involved in obtaining exact formulas for the corrections $\bar{H}_M^{(p)}$, $p = 1, 2, \dots$ to the coarse-grained Hamiltonian $\bar{H}_M^{(0)}$.

Computation of the term $\int (\Delta_{kk}J(\sigma))^2 \tilde{\rho}_k(d\sigma)$. Using the definition of $\Delta_{kk}J$ we write

$$\int (\Delta_{kk}J(\sigma))^2 \tilde{\rho}_k(d\sigma) = \sum_{\substack{x, y \in C_k \\ y \neq x}} \sum_{\substack{x', y' \in C_k \\ y' \neq x'}} (J(x - y) - \bar{J}(k, k))(J(x' - y') - \bar{J}(k, k)) \mathbb{E}[\sigma(x)\sigma(y)\sigma(x')\sigma(y')|\alpha].$$

In the above formula the expectation takes different values according to the following cases

- (1) Four particles: $x \neq x', y'$ and $y \neq x', y'$, in which case the expectation gives $E_4(\alpha)$.
- (2) Three particles: $x = x'$ and $y \neq y'$, or $y = y'$ and $x \neq x'$, or $x = y'$ and $x' \neq y$, or $x' = y$ and $x \neq y'$, in which case for some distinct x, y, z we obtain $\mathbb{E}[\sigma^2(x)\sigma(y)\sigma(z)|\alpha] = \mathbb{E}[\sigma(y)\sigma(z)|\alpha] = E_2(\alpha)$, since $\sigma^2(x) = 1$.
- (3) Two particles: $x = x'$ and $y = y'$, or $x = y'$ and $x' = y$, which for the distinct particle positions x, y gives $\mathbb{E}[\sigma^2(x)\sigma^2(y)|\alpha] = 1$.

We also keep in mind that we already have that $y \neq x$ and $y' \neq x'$. With the above cases we can substitute for $\mathbb{E}[\sigma(x)\sigma(y)\sigma(x')\sigma(y')|\alpha]$ the expression

$$\begin{aligned} & (1 - \delta_{x,x'})(1 - \delta_{x,y'})(1 - \delta_{y,x'}) (1 - \delta_{y,y'}) \times E_4(\alpha) \\ & + [\delta_{x,x'}(1 - \delta_{y,y'}) + \delta_{y,y'}(1 - \delta_{x,x'}) + \delta_{x,y'}(1 - \delta_{x',y}) + \delta_{x',y}(1 - \delta_{x,y'})] \times E_2(\alpha) \\ & + (\delta_{x,x'}\delta_{y,y'} + \delta_{x,y'}\delta_{x',y}) \times 1. \end{aligned}$$

We expand the above products and collect the terms with the factor $\delta_{x,x'}$ (note that the others with the factors $\delta_{x,y'}$, etc., are same by changing variables), and with the factor $\delta_{x,x'}\delta_{y,y'}$ and we obtain

$$4 \sum_{\substack{x, y \in C_k \\ y \neq x}} \sum_{\substack{x', y' \in C_k \\ y' \neq x'}} (J(x - y) - \bar{J}(k, k))(J(x' - y') - \bar{J}(k, k)) \delta_{x,x'} (-E_4(\alpha) + E_2(\alpha)) = 4j_{kk}^2 (-E_4(\alpha) + E_2(\alpha)),$$

and we also have

$$2 \sum_{\substack{x, y \in C_k \\ y \neq x}} \sum_{\substack{x', y' \in C_k \\ y' \neq x'}} (J(x - y) - \bar{J}(k, k))(J(x' - y') - \bar{J}(k, k)) \delta_{x,x'} \delta_{y,y'} (E_4 + 1 - 2E_2) = 2j_{kk}^1 (E_4(\alpha) + 1 - 2E_2(\alpha)).$$

Computation of the term $\int (\Delta_{kl}J(\sigma))^2 \tilde{\rho}_k(d\sigma) \tilde{\rho}_l(d\sigma)$. Similarly as in the calculation above we consider all possible cases for the position of the particles and we obtain

$$\begin{aligned} & \sum_{\substack{x \in C_k \\ y \in C_l}} \sum_{\substack{x' \in C_k \\ y' \in C_l}} (J(x-y) - \bar{J}(k,l))(J(x'-y') - \bar{J}(k,l)) \mathbb{E}[\sigma(x)\sigma(y)\sigma(x')\sigma(y')|\alpha] \\ &= \sum_{\substack{x \in C_k \\ y \in C_l}} \sum_{\substack{x' \in C_k \\ y' \in C_l}} (J(x-y) - \bar{J}(k,l))(J(x'-y') - \bar{J}(k,l)) \\ & \quad \times \left\{ (1 - \delta_{x,x'})(1 - \delta_{y,y'}) \mathbb{E}[\sigma(x)\sigma(x')|\alpha_k] \mathbb{E}[\sigma(y)\sigma(y')|\alpha_l] \right. \\ & \quad + \delta_{x,x'}(1 - \delta_{y,y'}) \mathbb{E}[\sigma^2(x)|\alpha_k] \mathbb{E}[\sigma(y)\sigma(y')|\alpha_l] \\ & \quad + \delta_{y,y'}(1 - \delta_{x,x'}) \mathbb{E}[\sigma(x)\sigma(x')|\alpha_k] \mathbb{E}[\sigma^2(y)|\alpha_l] \\ & \quad \left. + \delta_{x,x'}\delta_{y,y'} \mathbb{E}[\sigma^2(x)|\alpha_k] \mathbb{E}[\sigma^2(y)|\alpha_l] \right\} \\ &= \sum_{\substack{x \in C_k \\ y \in C_l}} \sum_{\substack{x' \in C_k \\ y' \in C_l}} (J(x-y) - \bar{J}(k,l))(J(x'-y') - \bar{J}(k,l)) \\ & \quad \times \left\{ \delta_{x,x'}(-E_2(\alpha_k)E_2(\alpha_l) + E_2(\alpha_l)) + \delta_{y,y'}(-E_2(\alpha_k)E_2(\alpha_l) + E_2(\alpha_k)) \right. \\ & \quad \left. + \delta_{x,x'}\delta_{y,y'}(E_2(\alpha_k)E_2(\alpha_l) - E_2(\alpha_l) - E_2(\alpha_k) + 1) \right\} \\ &= j_{kl}^2(-2E_2(\alpha_k)E_2(\alpha_l) + E_2(\alpha_k) + E_2(\alpha_l)) + j_{kl}^1(E_2(\alpha_k)E_2(\alpha_l) - E_2(\alpha_l) - E_2(\alpha_k) + 1). \end{aligned}$$

Computation of the term $\int \Delta_{kk}J(\sigma)\Delta_{kl}J(\sigma)\tilde{\rho}_k(d\sigma)\tilde{\rho}_l(d\sigma)$. In the same spirit we have

$$\begin{aligned} & \sum_{\substack{x,y \in C_k \\ y \neq x}} \sum_{\substack{x' \in C_k \\ y' \in C_l}} (J(x-y) - \bar{J}(k,k))(J(x'-y') - \bar{J}(k,l)) \mathbb{E}[\sigma(x)\sigma(y)\sigma(x')\sigma(y')|\alpha] \\ &= \sum_{\substack{x,y \in C_k \\ y \neq x}} \sum_{\substack{x' \in C_k \\ y' \in C_l}} (J(x-y) - \bar{J}(k,k))(J(x'-y') - \bar{J}(k,l)) \\ & \quad \times \left\{ (1 - \delta_{x,x'})(1 - \delta_{y,x'}) \mathbb{E}[\sigma(x)\sigma(y)\sigma(x')|\alpha_k] \mathbb{E}[\sigma(y')|\alpha_l] \right. \\ & \quad \left. + \delta_{x,x'} \mathbb{E}[\sigma^2(x)\sigma(y)|\alpha_k] \mathbb{E}[\sigma(y')|\alpha_l] + \delta_{y,x'} \mathbb{E}[\sigma(x)\sigma^2(y)|\alpha_k] \mathbb{E}[\sigma(y')|\alpha_l] \right\} \\ &= \sum_{\substack{x,y \in C_k \\ y \neq x}} \sum_{\substack{x' \in C_k \\ y' \in C_l}} (J(x-y) - \bar{J}(k,k))(J(x'-y') - \bar{J}(k,l)) \\ & \quad \times \left\{ -\delta_{x,x'}(E_3(\alpha_k)E_1(\alpha_l) - E_1(\alpha_k)E_1(\alpha_l)) - \delta_{y,x'}(E_3(\alpha_k)E_1(\alpha_l) - E_1(\alpha_k)E_1(\alpha_l)) \right\} \\ &= -2j_{kk}^2(E_3(\alpha_k)E_1(\alpha_l) - E_1(\alpha_k)E_1(\alpha_l)). \end{aligned}$$

Similar computation yields expressions $-2j_{ll}^2(E_3(\alpha_l)E_1(\alpha_k) - E_1(\alpha_l)E_1(\alpha_k))$ for the terms $f_l f_{kl}$.

Computation of the term $\int \Delta_{k_1 k_2} J(\sigma) \Delta_{k_2 k_3} J(\sigma) \tilde{\rho}_{k_1}(d\sigma) \tilde{\rho}_{k_2}(d\sigma) \tilde{\rho}_{k_3}(d\sigma)$. We repeat the same procedure and obtain

$$\begin{aligned} & \sum_{\substack{x \in C_{k_1} \\ y \in C_{k_2}}} \sum_{\substack{x' \in C_{k_2} \\ y' \in C_{k_3}}} (J(x-y) - \bar{J}(k_1, k_2))(J(x'-y') - \bar{J}(k_2, k_3)) \mathbb{E}[\sigma(x)\sigma(y)\sigma(x')\sigma(y')|\alpha] = \\ & \sum_{\substack{x \in C_{k_1} \\ y \in C_{k_2}}} \sum_{\substack{x' \in C_{k_2} \\ y' \in C_{k_3}}} (J(x-y) - \bar{J}(k_1, k_2))(J(x'-y') - \bar{J}(k_2, k_3)) \\ & \quad \times \left\{ (1 - \delta_{y,x'}) \mathbb{E}[\sigma(x)|\alpha_{k_1}] \mathbb{E}[\sigma(y)\sigma(x')|\alpha_{k_2}] \mathbb{E}[\sigma(y')|\alpha_{k_3}] \right. \\ & \quad \left. + \delta_{y,x'} \mathbb{E}[\sigma(x)|\alpha_{k_1}] \mathbb{E}[\sigma(y)^2|\alpha_{k_2}] \mathbb{E}[\sigma(y')|\alpha_{k_3}] \right\} \\ & \sum_{\substack{x \in C_{k_1} \\ y \in C_{k_2}}} \sum_{\substack{x' \in C_{k_2} \\ y' \in C_{k_3}}} (J(x-y) - \bar{J}(k_1, k_2))(J(x'-y') - \bar{J}(k_2, k_3)) \\ & \quad \times \delta_{y,x'} (-E_1(\alpha_{k_1})E_2(\alpha_{k_2})E_1(\alpha_{k_3}) + E_1(\alpha_{k_1})E_1(\alpha_{k_3})) \\ & = j_{k_1 k_2 k_3}^2 (-E_1(\alpha_{k_1})E_2(\alpha_{k_2})E_1(\alpha_{k_3}) + E_1(\alpha_{k_1})E_1(\alpha_{k_3})). \end{aligned}$$

APPENDIX B: SAMPLING OF THE EQUILIBRIUM MEASURE

In this section we briefly describe the background of Monte Carlo methods used in Section 4 for the sampling of the microscopic and coarse-grained equilibrium Gibbs measures. These algorithms rely on the construction of a suitable Markov Chain such that its unique invariant measure is the Gibbs measure we intend to sample.

Microscopic Monte Carlo algorithms. The sampling algorithm for the microscopic lattice system is given in terms a continuous-time jump Markov process that defines a change of the spin $\sigma(x)$ with the probability $c(x, \sigma)\Delta t$ over the time interval $[t, t + \Delta t]$. The function $c : \Lambda_N \times \mathcal{S}_N \rightarrow \mathbb{R}$ is called a rate of the process. The jump process $\{\sigma_t\}_{t \geq 0}$ is constructed in the following way: suppose that at the time t the configuration is σ_t , then the probability that over the time interval $[t, t + \Delta t]$ the spin at the site $x \in \Lambda_N$ spontaneously changes from $\sigma_t(x)$ to a new configuration $\sigma_{t+\Delta t}^x(x)$ is $c(x, \sigma)\Delta t + O(\Delta t^2)$. We denote the resulting configuration σ^x . We require that the dynamics is such that the invariant measure of this Markov process is the Gibbs measure (6). The sufficient condition is known as *detailed balance* and it imposes a condition on the form of the rate

$$c(x, \sigma)e^{-\beta H_N(\sigma)} = c(x, \sigma^x)e^{-\beta H_N(\sigma^x)}.$$

This condition has a simple interpretation $c(x, \sigma)$ is the rate of converting $\sigma(x)$ to the value $\sigma^x(x)$ while $c(x, \sigma^x)$ is the rate of changing the spin at the site x back to $\sigma(x)$. For the spin-flip dynamics we have $\sigma^x(y) = (1 - 2\delta_{xy})\sigma(y)$. The widely used class of Metropolis-type dynamics satisfies the detailed balance conditions and has the rate given by

$$c(x, \sigma) = G(\beta \Delta_x H_N(\sigma)), \text{ where } \Delta_x H_N(\sigma) = H_N(\sigma^x) - H_N(\sigma),$$

where G is a continuous function satisfying: $G(r) = G(-r)e^{-r}$ for all $r \in \mathbb{R}$. The most common choices in physics simulations are $G(r) = \frac{1}{1+e^r}$ (Glauber dynamics), $G(r) = e^{-[r]_+}$, (Metropolis dynamics), with $[r]_+ = r$ if $r \geq 0$ and $= 0$ otherwise, and $G(r) = e^{-r/2}$. Such dynamics are often used as samplers from the canonical equilibrium Gibbs measure.

Coarse-grained Monte Carlo algorithms. For the purpose of sampling the coarse-grained Gibbs measure given by $\bar{\mu}_{M,\beta}(\eta)$ we will consider a Markov jump process with the rates $\bar{c}_b(k, \eta), \bar{c}_d(k, \eta)$ correspond to the addition (“birth”) and removal of a particle (“death”) respectively, at the coarse cell k . The resulting process is a *birth-death* process $\{\eta_t\}_{t \geq 0}$ defined on the configuration space $(\Sigma^c)^{\bar{\Lambda}^M}$, where the state space is $\Sigma^c = \{-q, -q + 2, \dots, +q\}$ or $\Sigma^c = \{0, 1, \dots, q\}$.

As in the case of microscopic dynamics, we need to ensure the detailed balance condition for the coarse-grained rates expressed as

$$\bar{c}_b(k, \eta) \bar{\mu}_{M, \beta}(\eta) = \bar{c}_d(k, \eta + \delta_k) \bar{\mu}_{M, \beta}(\eta + \delta_k), \quad \text{and} \quad \bar{c}_d(k, \eta) \bar{\mu}_{M, \beta}(\eta) = \bar{c}_b(k, \eta - \delta_k) \bar{\mu}_{M, \beta}(\eta - \delta_k).$$

For the Metropolis-type dynamics the rates are given by

$$\bar{c}_b(k, \eta) = G(\bar{H}_M^{(0)}(\eta + \delta_k) - \bar{H}_M^{(0)}(\eta))(q - \eta(k)), \quad \text{and} \quad \bar{c}_d(k, \eta) = G(\bar{H}_M^{(0)}(\eta - \delta_k) - \bar{H}_M^{(0)}(\eta))\eta(k).$$

Notice that with these rates the detailed balance conditions hold since

$$\begin{aligned} G(\bar{H}_M^{(0)}(\eta + \delta_k) - \bar{H}_M^{(0)}(\eta))e^{-\bar{H}_M^{(0)}(\eta)} &= G(\bar{H}_M^{(0)}(\eta - \delta_k) - \bar{H}_M^{(0)}(\eta))e^{-\bar{H}_M^{(0)}(\eta + \delta_k)}, \\ G(\bar{H}_M^{(0)}(\eta - \delta_k) - \bar{H}_M^{(0)}(\eta))e^{-\bar{H}_M^{(0)}(\eta)} &= G(\bar{H}_M^{(0)}(\eta) - \bar{H}_M^{(0)}(\eta - \delta_k))e^{-\bar{H}_M^{(0)}(\eta - \delta_k)} \end{aligned}$$

and $G(r) = G(-r)e^{-r}$ for all $r \in \mathbb{R}$.

Higher-order coarse-grained Monte Carlo algorithms. In Theorem 1.4 we have suggested higher-order corrections ($\bar{H}_M^{(p)}$, for $p = 1, \dots$) to the coarse-grained Hamiltonian $\bar{H}_M^{(0)}$. Based on this equilibrium theory, and on the detailed balance condition, we define rates that are capable of sampling the corrected Gibbs measure corresponding to $\bar{H}_M^{(p)}$. In this case the Metropolis-type rates are given as

$$\bar{c}_a^{(p)}(k, \eta) = G(\bar{H}_M^{(p)}(\eta + \delta_k) - \bar{H}_M^{(p)}(\eta))(q - \eta(k)), \quad \text{and} \quad \bar{c}_d^{(p)}(k, \eta) = G(\bar{H}_M^{(p)}(\eta - \delta_k) - \bar{H}_M^{(p)}(\eta))\eta(k).$$

Acknowledgements. The research of M.K. was partially supported by DE-FG02-05ER25702, NSF-DMS-0413864 and NSF-ITR-0219211. The research of P.P. was partially supported by NSF-DMS-0303565. The research of L.R.-B. was partially supported by NSF-DMS-0306540. The research of D.T. was partially supported by NSF-DMS-0413864 and NSF-ITR-0219211.

REFERENCES

- [1] J. Bricmont, A. Kupiainen and R. Lefevre, Renormalization group pathologies and the definition of Gibbs states. *Comm. Math. Phys.* **194** (1998) 359–388.
- [2] C. Cammarota, Decay of correlations for infinite range interactions in unbounded spin systems. *Comm. Math. Phys.* **85** (1982) 517–528.
- [3] A. Chatterjee, M. Katsoulakis and D. Vlachos, Spatially adaptive lattice coarse-grained Monte Carlo simulations for diffusion of interacting molecules. *J. Chem. Phys.* **121** (2004) 11420–11431.
- [4] A. Chatterjee, M. Katsoulakis and D. Vlachos, Spatially adaptive grand canonical ensemble Monte Carlo simulations. *Phys. Rev. E* **71** (2005) 026702.
- [5] T.M. Cover and J.A. Thomas, *Elements of Information Theory*. John Wiley and Sons, Inc. (1991).
- [6] G.A. Gallavotti and S. Miracle-Sole, Correlation functions of a lattice system. *Comm. Math. Phys.* **7** (1968) 274–288.
- [7] N. Goldenfeld, *Lectures on Phase Transitions and the Renormalization Group*, Volume **85**. Addison-Wesley, New York (1992).
- [8] C. Gruber and H. Kunz, General properties of polymer systems. *Comm. Math. Phys.* **22** (1971) 133–161.
- [9] M. Hildebrand and A.S. Mikhailov, Mesoscopic modeling in the kinetic theory of adsorbates. *J. Chem. Phys.* **100** (1996) 19089.
- [10] A.E. Ismail, G.C. Rutledge and G. Stephanopoulos, Multiresolution analysis in statistical mechanics. I. Using wavelets to calculate thermodynamics properties. *J. Chem. Phys.* **118** (2003) 4414–4424.
- [11] A.E. Ismail, G.C. Rutledge and G. Stephanopoulos, Multiresolution analysis in statistical mechanics. II. Wavelet transform as a basis for Monte Carlo simulations on lattices. *J. Chem. Phys.* **118** (2003) 4424.
- [12] L. Kadanoff, Scaling laws for Ising models near t_c . *Physics* **2** (1966) 263.
- [13] M. Katsoulakis and J. Trashorras, Information loss in coarse-graining of stochastic particle dynamics. *J. Statist. Phys.* **122** (2006) 115–135.
- [14] M. Katsoulakis, A. Majda and D. Vlachos, Coarse-grained stochastic processes for microscopic lattice systems. *Proc. Natl. Acad. Sci.* **100** (2003) 782–782.
- [15] M.A. Katsoulakis, A.J. Majda and D.G. Vlachos, Coarse-grained stochastic processes and Monte Carlo simulations in lattice systems. *J. Comp. Phys.* **186** (2003) 250–278.

- [16] M.A. Katsoulakis, P. Plecháč, L. Rey-Bellet and D.K. Tsagkarogiannis, Coarse-graining schemes for lattice systems with short and long range interactions. (In preparation).
- [17] M.A. Katsoulakis, P. Plecháč and A. Sopasakis, Error analysis of coarse-graining for stochastic lattice dynamics. *SIAM J. Numer. Anal.* **44** (2006) 2270.
- [18] D.A. Lavis and G.M. Bell, *Statistical Mechanics of Lattice Systems, Volume I*. Springer Verlag (1999).
- [19] J.E. Mayer, Integral equations between distribution functions of molecules. *J. Chem. Phys.* **15** (1947) 187–201.
- [20] R. Peierls, On Ising's model of ferromagnetism. *Proc. Camb. Philos. Soc.* **32** (1936) 477–481.
- [21] I.V. Pivkin and G.E. Karniadakis, Coarse-graining limits in open and wall-bounded dissipative particle dynamics systems. *J. Chem. Phys.* **124** (2006) 184101.
- [22] A. Procacci, B.N.B. De Lima and B. Scoppola, A remark on high temperature polymer expansion for lattice systems with infinite range pair interactions. *Lett. Math. Phys.* **45** (1998) 303–322.
- [23] B. Simon, *The Statistical Mechanics of Lattice Gases, Vol. I*. Princeton series in Physics (1993).
- [24] A. Szepessy, R. Tempone and G.E. Zouraris, Adaptive weak approximation of stochastic differential equations. *Comm. Pure Appl. Math.* **54** (2001) 1169–1214.
- [25] A.C.D. van Enter, R. Fernández and A.D. Sokal, Regularity properties and pathologies of position-space renormalization-group transformations: scope and limitations of Gibbsian theory. *J. Statist. Phys.* **72** (1993) 879–1167.