

CONTINUUM LIMIT OF p -LAPLACIAN EVOLUTION PROBLEMS ON GRAPHS: L^q GRAPHONS AND SPARSE GRAPHS

IMAD EL BOUCHAIRI¹, JALAL M. FADILI^{1,*} AND ABDERRAHIM ELMOATAZ²

Abstract. In this paper we study continuum limits of the discretized p -Laplacian evolution problem on sparse graphs with homogeneous Neumann boundary conditions. This goes far beyond known results by handling much more general class of kernels, possibly singular, and graph sequences whose limit are the so-called L^q -graphons. More precisely, we derive a bound on the distance between two continuous-in-time trajectories defined by two different evolution systems (*i.e.*, with different kernels, second member and initial data). Similarly, we provide a bound in the case that one of the trajectories is discrete-in-time and the other is continuous. In turn, these results lead us to establish error estimates of the full discretization of the p -Laplacian problem on sparse random graphs. In particular, we provide rate of convergence of solutions for the discrete models to the solution of the continuous problem as the number of vertices grows.

Mathematics Subject Classification. 65N12, 34A12, 45G1, 05C90.

Received January 17, 2022. Accepted January 14, 2023.

1. INTRODUCTION

1.1. Problem formulation

Our main goal in this paper is to study the discretization of the following nonlinear diffusion problem on graphs, which we call the nonlocal p -Laplacian problem with homogeneous Neumann boundary conditions:

$$\begin{cases} \frac{\partial}{\partial t} u(\mathbf{x}, t) = \int_{\Omega} K(\mathbf{x}, \mathbf{y}) |u(\mathbf{y}, t) - u(\mathbf{x}, t)|^{p-2} (u(\mathbf{y}, t) - u(\mathbf{x}, t)) \, d\mathbf{y} + f(\mathbf{x}, t), & \mathbf{x} \in \Omega, t > 0, \\ u(\mathbf{x}, 0) = g(\mathbf{x}), & \mathbf{x} \in \Omega, \end{cases} \quad (\mathcal{P})$$

where $p \in [1, +\infty]$, $\Omega \subset \mathbb{R}^d$ is a bounded domain, $d \geq 1$, and $K : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ is the kernel function. Since our goal is to study nonlocal graph limits through the theory of L^q graphons, we assume in the sequel $\Omega = [0, 1]^d$. In the setting of graphons, one can even take $d = 1$ and it will be seen that K is the limit object for some convergent graph sequence $\{G_n\}_n$, $n \in \mathbb{N}$, whose meaning and form will be specified in the sequel. In terms of application, we are motivated by understanding collective dynamics in large ensembles of interacting dynamical systems

Keywords and phrases. Nonlocal diffusion, p -Laplacian, sparse graphs, L^q -graphons.

¹ Normandie Univ, ENSICAEN, UNICAEN, CNRS, GREYC, Caen, France.

² Normandie Univ, UNICAEN, CNRS, GREYC, Caen, France.

*Corresponding author: Jalal.Fadili@greyc.ensicaen.fr

with applications ranging from biology to physics and computer science (see later for a detailed discussion and review).

Throughout, we assume that

- (H.1) K is a nonnegative measurable function.
- (H.2) K is symmetric, *i.e.*, $K(\mathbf{x}, \mathbf{y}) = K(\mathbf{y}, \mathbf{x})$.
- (H.3) $\sup_{\mathbf{x} \in \Omega} \int_{\Omega} K(\mathbf{x}, \mathbf{y}) \, d\mathbf{y} < +\infty$.

By (H.2), it is straightforward to see that

$$\sup_{\mathbf{x} \in \Omega} \int_{\Omega} K(\mathbf{x}, \mathbf{y}) \, d\mathbf{y} = \sup_{\mathbf{y} \in \Omega} \int_{\Omega} K(\mathbf{x}, \mathbf{y}) \, d\mathbf{x},$$

and thus, (H.3) is equivalent to

$$\sup_{\mathbf{y} \in \Omega} \int_{\Omega} K(\mathbf{x}, \mathbf{y}) \, d\mathbf{x} < +\infty.$$

When the kernel is such that $K(\mathbf{x}, \mathbf{y}) = J(\mathbf{x} - \mathbf{y})$, where $J : \mathbb{R}^d \rightarrow \mathbb{R}$, then (H.1)–(H.3) read:

- (H[?].1) J is nonnegative and measurable.
- (H[?].2) J is symmetric, *i.e.*, $J(-\mathbf{x}) = J(\mathbf{x})$.
- (H[?].3) $\int_{\Omega - \Omega} J(\mathbf{x}) \, d\mathbf{x} < +\infty$.

Recall that $\Omega - \Omega$ is the Minkowski sum of Ω and $-\Omega$. In the case $\Omega = [0, 1]^d$, we obviously have $\Omega - \Omega = [-1, 1]^d$.

Partial differential equations (PDEs) involving the nonlocal p -Laplacian operator have become more and more popular both in the setting of Euclidean domains and on discrete graphs, as the p -Laplacian problem has been possessing many important features shared by many practical problems in mathematics, physics, engineering, biology, and economy, such as continuum mechanics, phase transition phenomena, population dynamics, see [2, 6, 7, 19, 27, 28, 45] and references therein. As explained in [27], when $K(\mathbf{x}, \mathbf{y}) = J(\mathbf{x} - \mathbf{y})$ with J radially symmetric with compact support verifying $J(0) > 0$ and $\int_{\mathbb{R}^d} J(\mathbf{x}) \, d\mathbf{x} = 1$, (P) has a nice interpretation in modeling the dispersal of organisms in space. In this case, $u(\mathbf{x}, t)$ can be viewed as their density at point \mathbf{x} and time t , $J(\mathbf{x} - \mathbf{y})$ is interpreted as the probability of jumping from position \mathbf{y} to position \mathbf{x} . In turn, the right hand side of (P) represents transport due to long-range dispersal mechanisms, that is the rate at which organisms are arriving to location \mathbf{x} from any other place. For $p = 2$ and $p = 1$, (P) correspond respectively to the nonlocal analogues of the heat equation and the total variation flow with homogeneous Neumann boundary conditions. The p -Laplacian, in its version on graphs, has found many applications in data processing, computer vision and machine learning [16, 18, 23–25, 35, 43, 44].

1.2. Contributions

In this work, our aim is to quantitatively analyze evolution problems of the form (P) driven by the p -Laplacian when discretized on *sparse* graphs, and to understand their continuum limit. The setting of sparse graphs affords great flexibility in applications where graph sequences can be of many types (biological, physical, social, meshes, images, etc.), whose details vary widely. More precisely:

- For $p \in [1, +\infty[$, we first provide a general error bound between the solutions of two evolution problems of the form (P) governed by two kernels and two initial data (see Thms. 5.1 and 5.2).
- This paves the way to studying consistency of numerical solutions regarded as a discretization of (P) with space discretization of the kernel K and initial data g , and forward and backward Euler discretization in time (see Thms. 6.3–6.5).

- Equipped with these results, and combining with probabilistic deviation arguments, we prove our main result, Theorem 7.1, which establishes an error bound that holds in probability for solution of a fully discretized problem on sparse random graphs (*i.e.*, those with unbounded average degree) whose limits are the so-called L^q -graphons [13, 14]. In a nutshell, this result states that error estimate between the discrete solution and the continuous one (that of (\mathcal{P})) decomposes in three terms. The first one corresponds to the error of spatial discretization of K , g and f . The second term comes from discretization in time (with appropriate exponents depending on p). The last term reflects the discretization of the problem on a sparse graph with n vertices. This term behaves as $(\rho_n n)^{-\beta/2}$ for any $\beta \in]0, 1[$, which vanishes as $n \rightarrow +\infty$ assuming that $\rho_n n \rightarrow +\infty$ (*i.e.*, the average graph degree tends to infinity). As ρ_n is the edge density parameter, and the graph is sparser as $\rho_n \rightarrow 0$, our result not only shows that discretization error degrades as the graphs becomes sparser, but also quantifies this precisely. Of course, our analysis recovers the result on dense graphs by just taking $\rho_n = 1$.

1.3. Relation to prior work

In the literature, numerous works, that will review shortly, have been carried out in the recent years to study continuum limits of evolution and variational problems on graphs. It is however important to stress the fact that our focus in this paper will be on nonlocal (in contrast to local ones) continuum limits of problems on graphs, and for which an appropriate limit object (L^q graphon introduced in the recent theory of graphs) of a sequence of graphs with an increasing number of vertices will be instrumental. This is more in line with a numerical analysis standpoint.

Kobayashi-type estimates

Motivated by either the existence, generation of nonlinear semigroups, or the algorithmic approximation of solutions to the abstract Cauchy problem

$$\begin{cases} \frac{\partial}{\partial t} v(t) + A(t)v(t) \ni f(t), \\ v(0) = g, \end{cases} \quad (\text{CP})$$

where $A(t)$ is a possibly time-dependent m -accretive operator with domain in a Banach space, several authors have considered some special implicit discretization schemes in different settings of (CP) (autonomous case, non-autonomous case, with or without f); see for instance [9, 10, 15, 20, 21, 34, 36, 37, 41]. The evolution problem (\mathcal{P}) we deal with is a particular case of (CP) in its autonomous case, *i.e.*, the operator $A(t) \equiv \Delta_p^K$ does not depend on time.

In particular, the work of [36] in the autonomous case with $f = 0$ constructed sequences of approximate solutions which converge in an appropriate sense to a solution to (CP). The key argument is an inequality (known as Kobayashi inequality) that estimates the distance between arbitrary points of two independent sequences generated by the so called proximal iterations, from which one can derive quantitative estimates to compare the continuous and discrete trajectories using the backward Euler scheme, as well as Lipschitz continuity properties of the solution trajectory. In this respect, all above cited of works will only capture the time discretization error, while our aim is also to establish space-discretization bounds. For this, one may think of using the results of [1] who generalized those of [36, 37] to the non-autonomous case as well as to the case where the trajectories are defined by two abstract Cauchy systems driven by two m -accretive operators A_1 and A_2 . The latter bounds, expressed in our notation, are provided only in terms of the difference between minimal norm elements of $A_1 g$ and $A_2 g$.

Our work goes much further by exploiting key properties of the p -Laplacian operator to get precise estimates in terms of the data. This is more meaningful in our context where we recall that the goal is to study the fully discretized nonlocal p -Laplacian problem on graphs. For the time discretization error, not only we tackle forward Euler discretization, which is much easier to implement in practice (and what many practitioners actually use), but even for backward Euler discretization our error estimates are strictly better than the standard results in [20, 36, 41] (see the detailed discussion after the proof of Thm. 6.5).

Non-local continuum limits

Our motivation comes from understanding collective dynamics in large ensembles of interacting dynamical systems which is a fundamental problem in nonlinear science with applications ranging from biology (*e.g.*, neural and genetic networks) to physics and chemistry (*e.g.*, interacting particle systems) and computer science (internet, data science, multiagent systems, opinion dynamics).

In particular, the work of [33, 40] focused on a nonlinear heat equation on sparse graphs, where Lipschitz-continuity of the operator played key role in their analysis, starting from well-posedness to consistency and error bounds. Our work differs from those works in many crucial aspects. The nonlocal p -Laplacian evolution problem considered here is much more general and cannot be covered by the approach of those previous papers because the lack of Lipschitzianity raises several challenges (including well-posedness and error estimates). Unlike those previous works, we also consider both the semi-discrete and fully-discrete versions with both forward and backward Euler approximations, that we fully characterize, and develop novel proof techniques. Moreover, while those previous results are asymptotic using central limit theorem-type arguments, our results on the other hand are non-asymptotic, rely on deviation inequalities with explicit convergence rates.

In [30], the authors derived the continuum limit of the p -Laplacian Cauchy problem, but was confined to the restrictive case of bounded kernels, bounded initial data and dense graphs. The kernels, initial data and graphs considered here are beyond reach of the approach developed in those works. Our setting is much more challenging and has not been considered in the literature to the best of our knowledge. We are then able to consider unbounded initial data and most importantly a far larger class of kernels (including singular ones). This in turn will allow to handle sparse graph sequences, and our error bounds reveal the key role played by the average degree in the convergence rate. Our framework also allows to consider the non-smooth case of $p = 1$ where the p -Laplacian is set-valued, while this case could not be addressed by Hafiene *et al.* [30]. Moreover, our error bounds are directly stated in $L^2(\Omega)$ and not in $L^p(\Omega)$ as done in this previous work. Our proof is also simpler, more elegant and the argument is made more transparent.

Motivated by applications in opinion dynamics, the authors in [5] used ideas from [39] to study a collective dynamics model with a time-varying kernel and Lipschitz continuous operator. They showed nonlocal continuum limits as the number of agents/vertices goes to infinity. Interestingly, those authors used such this continuum limit to derive the mean-field limit one. In fact, the graph limit approach is more general than the mean-field limit one and this subordination of the mean-field limit to the nonlocal continuum graph limit is natural as pointed out to in [11]. Mean-field limits of interacting particle systems using the theory of graphons is attracting increasing interest. Studying mean field limits for problems on graphs with the p -Laplacian is an open problem that we leave to a future work.

Local continuum limits

The literature on deriving local continuum limits of variational and evolution problems on graphs, including with the p -Laplacian operator, is also very active; see [16, 18, 23, 43, 44] and many others. The motivation comes essentially from data science and machine learning problems. The idea is to take $K(\mathbf{x}, \mathbf{y}) = \varepsilon^{-d} J((\mathbf{x} - \mathbf{y})/\varepsilon)$, $\varepsilon > 0$, for some radially symmetric kernel J , and to study the limit (in an appropriate topology) as the number of vertices n grows to infinity and for ε vanishing with n at an appropriate rate. This line of work is however fundamentally different from our perspective.

1.4. Paper organization

The rest of the paper is organized as follows. In Section 2 we start by reviewing some basic notations and recall some preliminary material necessary to our exposition. In Section 3, we provide some prerequisites on L^q -graphons and sparse K -random graph models that we are going to deal with. Section 4 is devoted to study the well-posedness of the problem (\mathcal{P}) . In Section 5, we study stability of the problem (\mathcal{P}) with respect to sequences of kernels K , initial data g and second member f . Error bounds for the semi-discrete (*i.e.*, space discretization of (K, g, f)) problem are established in Section 6.1, and those for the fully discrete (time and

space discretization) problem with forward and backward Euler time-discretization are provided in Section 6.2. Section 7 is devoted to applying these results to fully discretized problems on sparse random graph models.

2. PRELIMINARIES

2.1. Basic notations

For $q \in [1, +\infty]$, and $S \subset \mathbb{R}^d$, $L^q(S)$ is the standard Banach space of Lebesgue q -integrable functions on S . For a function $F : S \times S \rightarrow \mathbb{R}$, we define the $L^\infty, q(S^2)$ -norm as

$$\|F\|_{L^\infty, q(S^2)} \stackrel{\text{def}}{=} \sup_{\mathbf{x} \in S} \|F(\mathbf{x}, \cdot)\|_{L^q(S)}.$$

If F is symmetric, then

$$\|F\|_{L^\infty, q(S^2)} = \sup_{\mathbf{y} \in S} \|F(\cdot, \mathbf{y})\|_{L^q(S)}.$$

$L^\infty, q(S^2)$ is the space of functions on S^2 of bounded $L^\infty, q(S^2)$ -norm, which is of course a Banach space.

Throughout the paper, we will often use Fubini's theorem without explicitly referring to it.

$C([0, T]; L^q(S))$ denotes the space of functions $u : S \times [0, T] \rightarrow \mathbb{R}$ which are uniformly continuous in time with values $u(t, \cdot)$ in $L^q(S)$. $C([0, T]; L^q(S))$ is naturally endowed with the norm $\|u\|_{C([0, T]; L^q(S))} \stackrel{\text{def}}{=} \sup_{t \in [0, T]} \|u(\cdot, t)\|_{L^q(S)}$. Moreover, $L^1([0, T]; L^q(S))$ is the space of functions $u : S \times [0, T] \rightarrow \mathbb{R}$ such that $\|u\|_{L^1([0, T]; L^q(S))} \stackrel{\text{def}}{=} \int_0^T \|u(\cdot, t)\|_{L^q(S)} dt < +\infty$.

We define the q -norm on \mathbb{R}^m , $q \in [1, +\infty[$, as

$$\|\mathbf{v}\|_q = \left(\frac{1}{m} \sum_{i \in [m]} |\mathbf{v}_i|^q \right)^{\frac{1}{q}},$$

with the usual adaptation for $q = +\infty$. $(\cdot)_+$ is the positive part function on \mathbb{R} .

For a set-valued operator $A : X \rightarrow 2^X$ on a Banach space X , its domain and range are respectively

$$\text{dom}(A) = \{u \in X : Au \neq \emptyset\} \text{ and } \text{ran}(A) = A(X).$$

2.2. Projector and injector

Let $n \in \mathbb{N}^*$ and denote the multi-index $\mathbf{i} = (i_1, i_2, \dots, i_d) \in [n]^d$. Partition Ω into cells (hypercubes)

$$\Omega_{\mathbf{i}}^{(n)} \stackrel{\text{def}}{=} \left\{ \prod_{k=1}^d]\mathbf{x}_{i_k-1}, \mathbf{x}_{i_k}] : \mathbf{i} \in [n]^d \right\}$$

of size $h_{\mathbf{i}} \stackrel{\text{def}}{=} |\Omega_{\mathbf{i}}^{(n)}|$, and maximal mesh size

$$\delta_n \stackrel{\text{def}}{=} \max_{\mathbf{i} \in [n]^d} \max_{k \in [d]} (|\mathbf{x}_{i_k} - \mathbf{x}_{i_k-1}|).$$

When the cells are equi-spaced, then $h_{\mathbf{i}} = 1/n^d$.

We consider the operator $P_n : L^1(\Omega) \rightarrow \mathbb{R}^{n^d}$

$$(P_n u)_{\mathbf{i}} \stackrel{\text{def}}{=} \frac{1}{h_{\mathbf{i}}} \int_{\Omega_{\mathbf{i}}^{(n)}} u(\mathbf{x}) d\mathbf{x}. \tag{1}$$

This operator can be also seen as a piecewise constant projector of u on the space of discrete functions. For simplicity, and with a slight abuse of notation, we keep the same notation for the projector $P_n : L^1(\Omega^2) \rightarrow \mathbb{R}^{n^d \times n^d}$.

Our aim is to study the relationship between solutions of discrete approximations and the solution of the continuum model. It is then convenient to introduce an intermediate model which is the continuum extension of the discrete solution. Towards this goal, we consider the piecewise constant injector I_n of a vector $\mathbf{v} \in \mathbb{R}^{n^d}$ into $L^2(\Omega)$ defined as

$$I_n \mathbf{v}(\mathbf{x}) \stackrel{\text{def}}{=} \sum_{\mathbf{i} \in [n]^d} \mathbf{v}_i \chi_{\Omega_i^{(n)}}(\mathbf{x}), \tag{2}$$

where we recall that χ_C is the characteristic function of the set C , *i.e.*, takes 1 on C and 0 otherwise.

It is immediate to see that the operator $I_n P_n$ is the orthogonal projector on the subspace $\text{Span}\{\chi_{\Omega_i^{(n)}} : \mathbf{i} \in [n]^d\}$ of $L^1(\Omega)$. In turn, $I_n P_n u$ is the the piecewise constant approximation of u .

Lemma 2.1. *For a function $u \in L^q(\Omega)$, $q \in [1, +\infty]$, we have*

$$\|I_n P_n u\|_{L^q(\Omega)} \leq \|u\|_{L^q(\Omega)}. \tag{3}$$

For a function $K \in L^{\infty,q}(\Omega^2)$, $q \in [1, +\infty]$, we have

$$\|I_n P_n K\|_{L^{\infty,q}(\Omega^2)} \leq \|K\|_{L^{\infty,q}(\Omega^2)}. \tag{4}$$

Proof. We prove (4) as (3) is a consequence of it. Let $\mathbf{K} = P_n K$. We have, $\forall \mathbf{x} \in \Omega$,

$$\begin{aligned} \int_{\Omega} I_n P_n |K(\mathbf{x}, \mathbf{y})|^q d\mathbf{y} &= \int_{\Omega} \sum_{\mathbf{i}, \mathbf{j}} |\mathbf{K}_{\mathbf{i}, \mathbf{j}}|^q \chi_{\Omega_i^{(n)}}(\mathbf{x}) \chi_{\Omega_j^{(n)}}(\mathbf{y}) d\mathbf{y} \\ &= \sum_{\mathbf{i}} \left(\sum_{\mathbf{j}} \int_{\Omega_j^{(n)}} |\mathbf{K}_{\mathbf{i}, \mathbf{j}}|^q d\mathbf{y} \right) \chi_{\Omega_i^{(n)}}(\mathbf{x}) \\ &= \sum_{\mathbf{i}} \left(\sum_{\mathbf{j}} h_j \left| \frac{1}{h_i h_j} \int_{\Omega_i^{(n)} \times \Omega_j^{(n)}} K(\mathbf{x}', \mathbf{y}') d\mathbf{x}' d\mathbf{y}' \right|^q \right) \chi_{\Omega_i^{(n)}}(\mathbf{x}) \\ &\leq \sum_{\mathbf{i}} \left(\sum_{\mathbf{j}} \frac{1}{h_i} \int_{\Omega_i^{(n)} \times \Omega_j^{(n)}} |K(\mathbf{x}', \mathbf{y}')|^q d\mathbf{x}' d\mathbf{y}' \right) \chi_{\Omega_i^{(n)}}(\mathbf{x}) \\ &= \sum_{\mathbf{i}} \left(\frac{1}{h_i} \int_{\Omega_i^{(n)}} \left(\sum_{\mathbf{j}} \int_{\Omega_j^{(n)}} |K(\mathbf{x}', \mathbf{y}')|^q d\mathbf{y}' \right) d\mathbf{x}' \right) \chi_{\Omega_i^{(n)}}(\mathbf{x}) \\ &= \sum_{\mathbf{i}} \left(\frac{1}{h_i} \int_{\Omega_i^{(n)}} \left(\int_{\Omega} |K(\mathbf{x}', \mathbf{y}')|^q d\mathbf{y}' \right) d\mathbf{x}' \right) \chi_{\Omega_i^{(n)}}(\mathbf{x}) \\ &\leq \|K\|_{L^{\infty,q}(\Omega^2)}^q \sum_{\mathbf{i}} \chi_{\Omega_i^{(n)}}(\mathbf{x}) = \|K\|_{L^{\infty,q}(\Omega^2)}^q. \end{aligned}$$

Taking the supremum on the left-hand side yields the bound. □

2.3. Lipschitz spaces

For $N \in \mathbb{N}^*$, let S be a compact subset of \mathbb{R}^N . We introduce the Lipschitz spaces $\text{Lip}(s, L^q(S))$, $q \in [1, +\infty]$, which contain functions with, roughly speaking, s “derivatives” in $L^q(S)$ ([22], Chap. 2, Sect. 9).

Definition 2.1. For $F \in L^q(S)$, $q \in [1, +\infty]$, we define the (first-order) $L^q(S)$ modulus of smoothness by

$$\omega(F, h)_q \stackrel{\text{def}}{=} \sup_{z \in \mathbb{R}^d, |z| < h} \left(\int_{\mathbf{x}, \mathbf{x}+z \in S} |F(\mathbf{x}+z) - F(\mathbf{x})|^q d\mathbf{x} \right)^{1/q}. \quad (5)$$

The Lipschitz spaces $\text{Lip}(s, L^q(S))$ consist of all functions F for which

$$|F|_{\text{Lip}(s, L^q(S))} \stackrel{\text{def}}{=} \sup_{h > 0} h^{-s} \omega(F, h)_q < +\infty.$$

We restrict ourselves to values $s \in]0, 1]$ since for $s > 1$, only constant functions are in $\text{Lip}(s, L^q(S))$. It is easy to see that $|F|_{\text{Lip}(s, L^q(S))}$ is a semi-norm. $\text{Lip}(s, L^q(S))$ is endowed with the norm

$$\|F\|_{\text{Lip}(s, L^q(S))} \stackrel{\text{def}}{=} \|F\|_{L^q(S)} + |F|_{\text{Lip}(s, L^q(S))}.$$

The space $\text{Lip}(s, L^q(S))$ is the Besov space $B_{q, \infty}^s$ ([22], Chap. 2, Sect. 10) which are very popular in approximation theory. In particular, $\text{Lip}(s, L^{1/s}(S))$ contains the space $\text{BV}(S)$ of functions of bounded variation on S ; see Chapter 2, Lemma 9.2 of [22]. Thus Lipschitz spaces are rich enough to contain functions with both discontinuities and fractal structure.

We now state the following approximation error bounds whose proofs use standard arguments from approximation theory; see Section 6.2.1 of [30] for details.

Lemma 2.2. *There exists a positive constant C_s , depending only on s , such that for all $F \in \text{Lip}(s, L^q(S))$, $s \in]0, 1]$, $q \in [1, +\infty]$,*

$$\|F - \mathbf{I}_n \mathbf{P}_n F\|_{L^q(S)} \leq C_s \delta_n^s |F|_{\text{Lip}(s, L^q(S))}. \quad (6)$$

We denote by $\text{BV}([0, T]; L^q(S))$ the Banach space of functions $f : \Omega \times [0, T] \rightarrow \mathbb{R}$ such that

$$\text{Var}_q(f) \stackrel{\text{def}}{=} \sup_{0 \leq t_0 < t_1 < \dots < t_N \leq T} \sum_{i=1}^N \|f(\cdot, t_i) - f(\cdot, t_{i-1})\|_{L^q(S)} < +\infty,$$

endowed with the norm $\|f\|_{\text{BV}([0, T]; L^q(S))} \stackrel{\text{def}}{=} \|f(0)\|_{L^q(S)} + \text{Var}_q(f)$.

3. THE SPARSE GRAPH MODEL

One of our fundamental goals in this paper is to understand the behaviour of (\mathcal{P}) when discretized on a sequence of networks. Such networks can be of many types (biological, physical, social, data processing, etc.), whose details vary widely, but which bear similar structural phenomena. In turn, one may wonder whether discretization of (\mathcal{P}) is stable to the particular realization of the network and input data. It is then natural to consider a sequence of graphs with size tending to infinity and ask whether the solutions of discrete forms of (\mathcal{P}) on these graphs converge to any meaningful sort of limit. This will allow us in turn to establish a continuum limit of the solutions to these discrete evolution problems. For this, we need to be equipped with an appropriate theory of graph limits.

3.1. L^q graphons and graph limits

In [13, 14], the authors laid the foundations of L^q graphons, For $q \geq 1$, where the term L^q graphon refers to a symmetric function $K \in L^q([0, 1]^2)$. They developed a theory of limits for sequences of sparse graphs based on such graphons, which generalizes both the existing theory of bounded graphons that are tailored to dense graph limits [38], and its extension in [12] to sparse graphs under a no dense spots assumptions. The

latter graph model was studied in [31] in the context of continuum limits of p -Laplacian evolution problems on graphs. Nevertheless, the boundedness assumption of the graphon underlying these graph models is still highly restrictive. In particular, it does not allow to handle singular graphons and corresponding network models which have statistics governed by power laws. The theory of L^q graphons allows to analyze graphs with power law degree distributions, hence providing a broadly applicable limit theory for sparse graphs with unbounded average degree.

3.2. Sparse K -random graph models

We consider weighted graphs, which include as a special case simple unweighted graphs. Let $G = (V(G), E(G))$, be a weighted graph with vertex set $V(G)$ and edge set $E(G) \subseteq V(G)^2$, respectively. In G , every edge $(i, j) \in E(G)$ (allowing loops with $i = j$) is given a weight $\beta_{ij} \in \mathbb{R}_+^1$. We set $\beta_{ij} = 0$ whenever $(i, j) \notin E(G)$.

The theory of random graphs was founded in the 50’s–60’s by Erdős and Rényi [26], who started the systematic study of the space of graphs with n labeled vertices and $M = M(n)$ edges, with all graphs equiprobable. The aim is to turn the set of all graphs with n vertices into a probability space. Intuitively we should be able to generate a sequence of graphs $\{G_n\}_{n \in \mathbb{N}}$ randomly as follows: for each edge $(i, j) \in [n]^2$, we decide by some random experiment whether or not (i, j) shall be an edge of G_n , these experiments are performed independently.

The idea underlying the sparse K -random graph model proposed by Borgs *et al.* [14] is that each L^q graphon K gives rise to a natural random graph model, which produces a sequence of sparse graphs converging to K in an appropriate metric. Inspired by their work, we propose the following construction.

Definition 3.1. Fix $n \in \mathbb{N}^*$, let K be an L^1 graphon and $\rho_n > 0$. Take the equi-spaced partition of $[0, 1]$ in intervals $]x_{i-1}, x_i]$, $i \in [n]$, where $x_i = i/n$. Let $\mathbf{K} \in \mathbb{R}_+^{n \times n}$ be a weight matrix such that:

(H_w.1) $\|I_n \mathbf{K} - K\|_{L^1([0,1]^2)} \rightarrow 0$ as $n \rightarrow +\infty$.

(H_w.2) $\|I_n \mathbf{K}(x, \cdot) - K(x, \cdot)\|_{L^1([0,1])} \rightarrow 0$ uniformly in $x \in [0, 1]$.

Generate the random graph

$$G_n = (V(G_n), E(G_n)) \stackrel{\text{def}}{=} \mathbf{G}(n, K, \rho_n)$$

as follows: join each pair $(i, j) \in [n]^2$ of vertices independently, with probability

$$\mathbb{P}((i, j) \in E(G_n) | \mathbf{X}) = \rho_n \widehat{\mathbf{K}}_{ij}, \quad \text{where } \widehat{\mathbf{K}}_{ij} \stackrel{\text{def}}{=} \min(\mathbf{K}_{ij}, \rho_n^{-1}). \tag{7}$$

Remark 3.1. In the original sparse K -random graph model defined in [14], the x_i ’s are random iid samples drawn from the uniform distribution on $[0, 1]$. Moreover, $\mathbf{K}_{ij} = K(x_i, x_j)$. In this case, it follows from Theorem 2.14(a) of [14] (which relies on theorem from [32]) that assumptions (H_w.1) holds with probability 1.

Another interesting case is where $\mathbf{K} = P_n K$. Thanks to Lemma 2.1, $\|I_n P_n \mathbf{K}\|_{L^1(\Omega^2)} \leq \|K\|_{L^1(\Omega^2)}$ with probability 1. Thus, the Lebesgue differentiation theorem and the dominated convergence theorem allow to assert that $I_n P_n \mathbf{K}$ converges to K in $L^1(\Omega^2)$. In turn, assumption (H_w.1) holds.

For appropriate choices of ρ_n , the graph model constructed according to Definition 3.1 allows to sample both dense and sparse graphs from the graphon K . In particular, the sparsity assumption $\rho_n \rightarrow 0$ reflects the fact that ρ_n needs to be arbitrarily close to zero in order to see the unbounded/singular part of K . The assumption that $n\rho_n \rightarrow +\infty$ means the average degree tends to infinity. To check this, the average number of edges in this graph model is

$$\mathbb{E}(E(\mathbf{G}(n, K, \rho_n))) = \rho_n n^2 \left(n^{-2} \sum_{(i,j) \in [n]^2} \widehat{\mathbf{K}}_{ij} \right)$$

¹In [14], the weights are even allowed to be negative, but we will not consider this situation which is meaningless in our context.

$$= \rho_n n^2 \left(\|I_n \mathbf{K}\|_{L^1([0,1]^2)} - \left\| (I_n \mathbf{K} - \rho_n^{-1})_+ \right\|_{L^1([0,1]^2)} \right).$$

By assumption **(H_w.1)**, we have $\|I_n \mathbf{K}\|_{L^1([0,1]^2)} = \|K\|_{L^1([0,1]^2)} + o(1)$. Moreover, since $\rho_n \rightarrow 0$, we have from **(8)** that $\left\| (I_n \mathbf{K} - \rho_n^{-1})_+ \right\|_{L^1([0,1]^2)} = o(1)$. In turn,

$$\mathbb{E}(E(\mathbf{G}(n, K, \rho_n))) = \rho_n n^2 \left(\|K\|_{L^1([0,1]^2)} + o(1) \right).$$

As expected, this gives rise to a sparse graph whose edge density is $\rho_n \rightarrow 0$. For the average degree of this graph model, arguing similarly to above, and using **(H_w.2)**, we have

$$\begin{aligned} \mathbb{E}(\deg_{G_n}(i)) &= \rho_n n \left(n^{-1} \sum_{j \in [n]} \widehat{\mathbf{K}}_{ij} \right) \\ &= \rho_n n \left(\|I_n \mathbf{K}(x_i, \cdot)\|_{L^1([0,1])} - \left\| (I_n \mathbf{K}(x_i, \cdot) - \rho_n^{-1})_+ \right\|_{L^1([0,1])} \right) \\ &= \rho_n n \left(\int_0^1 K(x_i, y) \, dy + o(1) \right). \end{aligned}$$

As anticipated, the average degree is indeed unbounded since $\rho_n n \rightarrow +\infty$.

The above sequence of graphs generated also enjoys the following convergence result.

Proposition 3.1. *Let K be an L^1 graphon and \mathbf{K} be a weight matrix such that **(H_w.1)** holds. If $\rho_n > 0$ with $\rho_n \rightarrow 0$ and $n\rho_n \rightarrow +\infty$ as $n \rightarrow +\infty$, then $\rho_n^{-1} \mathbf{G}(n, K, \rho_n)$ converges almost surely to K in the cut distance metric (see [13, 14] for details about this metric).*

Proof. We essentially adapt the arguments of in the proof of Theorem 2.14(b) from [14]. More precisely, since **(H_w.1)** holds, one has to show ([14], (7.1)). For this, we invoke Lemma 7.3 of [14] by checking the condition (7.3) therein. We have by sub-linearity of $(\cdot)_+$ that

$$\begin{aligned} \frac{1}{n^2} \sum_{(i,j) \in [n]^2} (\mathbf{K}_{ij} - \rho_n^{-1})_+ &= \int_{[0,1]^2} (I_n \mathbf{K}(x, y) - \rho_n^{-1})_+ \, dx \, dy \\ &\leq \int_{[0,1]^2} (I_n \mathbf{K}(x, y) - K(x, y))_+ \, dx \, dy + \int_{[0,1]^2} (K(x, y) - \rho_n^{-1})_+ \, dx \, dy \quad (8) \\ &\leq \|I_n \mathbf{K} - K\|_{L^1([0,1]^2)} + \int_{[0,1]^2} (K(x, y) - \rho_n^{-1})_+ \, dx \, dy. \end{aligned}$$

The right-hand side in the above display goes to 0 as $n \rightarrow +\infty$ by **(H_w.1)** and since $\rho_n \rightarrow 0$. Indeed, for every $L > 0$, the limit superior of the last term is bounded by $\|(K - L)_+\|_{L^1([0,1]^2)}$, and this can be made arbitrarily small by choosing L large. \square

Example 3.1. For an example that cannot be handled using L^∞ graphons, and thus does not enter in the framework of [30, 31], consider a K -random graph model $\mathbf{G}(n, K, \rho_n)$ constructed according to Definition 3.1 with $\mathbf{K} = P_n K$, where $K(x, y) = J(x - y)$, $J : z \in [-1, 1] \mapsto 2^{-1}(1 - \beta)(2 - \beta)|z|^{-\beta}$, $\beta \in]0, 1[$. First, observe that the radially symmetric kernel J is singular but fulfills all assumptions **(H'.1)**–**(H'.3)**. In addition, by virtue of Remark 3.1, **(H_w.1)** and **(H_w.2)** also hold with

$$\|K\|_{L^1([0,1]^2)} = 1 \text{ and } \int_0^1 K(x, y) \, dy = 2^{-1}(2 - \beta)(x^{1-\beta} + (1 - x)^{1-\beta}) \in 2^{-1}(2 - \beta)[1, 2^\beta].$$

We also have the following convergence result in the $L^{\infty,1}$ norm that will be instrumental in Section 7. According to the construction in Definition 3.1, we let $\mathbf{\Lambda}_{ij}$, $(i, j) \in [n]^2, i \neq j$, be random variables such that $\rho_n \mathbf{\Lambda}_{ij}$ follows a Bernoulli distribution with parameter $\rho_n \widehat{\mathbf{K}}_{ij}$. For each row $i \in [n]$, $(\mathbf{\Lambda}_{ij})_{j \in [n]}$ are independent.

Lemma 3.1. *Let K be an $L^{\infty,1}$ graphon, i.e., it satisfies (H.1)–(H.3). Take the weight matrix $\mathbf{K} = P_n K$. Assume that $\rho_n \rightarrow 0$ and $n\rho_n = \omega((\log n)^\gamma)$ for some $\gamma > 1$. Then with probability 1,*

$$\|\mathbf{I}_n \mathbf{\Lambda}\|_{L^{\infty,1}([0,1]^2)} - \|\mathbf{I}_n \widehat{\mathbf{K}}\|_{L^{\infty,1}([0,1]^2)} \rightarrow 0.$$

If, moreover, (H_w.2) holds, then

$$\|\mathbf{I}_n \mathbf{\Lambda}\|_{L^{\infty,1}([0,1]^2)} \rightarrow \|K\|_{L^{\infty,1}([0,1]^2)}.$$

with probability 1.

Proof. For any $\varepsilon > 0$, we have by the union bound

$$\begin{aligned} \mathbb{P}\left(\left|\|\mathbf{I}_n \mathbf{\Lambda}\|_{L^{\infty,1}([0,1]^2)} - \|\mathbf{I}_n \widehat{\mathbf{K}}\|_{L^{\infty,1}([0,1]^2)}\right| \geq \varepsilon\right) &= \mathbb{P}\left(\left|\max_i \sum_j \mathbf{\Lambda}_{ij} - \max_i \sum_j \widehat{\mathbf{K}}_{ij}\right| \geq \varepsilon n\right) \\ &= \mathbb{P}\left(\left|\max_i \sum_j \rho_n \mathbf{\Lambda}_{ij} - \max_i \sum_j \rho_n \widehat{\mathbf{K}}_{ij}\right| \geq \varepsilon \rho_n n\right) \\ &\leq \mathbb{P}\left(\max_i \left|\sum_j \rho_n (\mathbf{\Lambda}_{ij} - \widehat{\mathbf{K}}_{ij})\right| \geq \varepsilon \rho_n n\right) \\ &\leq \sum_i \mathbb{P}\left(\left|\sum_j \rho_n (\mathbf{\Lambda}_{ij} - \widehat{\mathbf{K}}_{ij})\right| \geq \varepsilon \rho_n n\right). \end{aligned}$$

Since $(\rho_n \mathbf{\Lambda}_{ij})_j$ are independent Bernoulli variables with means $(\rho_n \widehat{\mathbf{K}}_{ij})_j$, it follows from the variant of the Chernoff bound in Lemma 7.1 of [14], that for every $\varepsilon > 0$,

$$\begin{aligned} \mathbb{P}\left(\left|\|\mathbf{I}_n \mathbf{\Lambda}\|_{L^{\infty,1}([0,1]^2)} - \|\mathbf{I}_n \widehat{\mathbf{K}}\|_{L^{\infty,1}([0,1]^2)}\right| \geq \varepsilon\right) &\leq 2 \sum_i \exp\left(-\frac{1}{3} \min\left(\frac{\varepsilon \rho_n n}{\rho_n \sum_j \widehat{\mathbf{K}}_{ij}}, 1\right) \varepsilon \rho_n n\right) \\ &\leq 2n \exp\left(-\frac{1}{3} \min\left(\frac{\varepsilon}{\|\mathbf{I}_n \widehat{\mathbf{K}}\|_{L^{\infty,1}([0,1]^2)}}, 1\right) \varepsilon \rho_n n\right) \\ &\leq 2n \exp\left(-\frac{1}{3} \min\left(\frac{\varepsilon}{\|K\|_{L^{\infty,1}([0,1]^2)}}, 1\right) \varepsilon \omega((\log n)^\gamma)\right) \\ &\leq 2n^{-\omega((\log n)^{\gamma-1})}, \end{aligned}$$

since $\gamma > 1$, and where we used (7) and Lemma 2.1 to show that

$$\|\mathbf{I}_n \widehat{\mathbf{K}}\|_{L^{\infty,1}([0,1]^2)} \leq \|\mathbf{I}_n \mathbf{K}\|_{L^{\infty,1}([0,1]^2)} = \|\mathbf{I}_n P_n K\|_{L^{\infty,1}([0,1]^2)} \leq \|K\|_{L^{\infty,1}([0,1]^2)}.$$

Invoking the (first) Borel–Cantelli lemma, we have the first claim. On the other hand,

$$\left|\|\mathbf{I}_n \widehat{\mathbf{K}}\|_{L^{\infty,1}([0,1]^2)} - \|K\|_{L^{\infty,1}([0,1]^2)}\right| \leq \|\mathbf{I}_n \widehat{\mathbf{K}} - K\|_{L^{\infty,1}([0,1]^2)}$$

$$\begin{aligned}
&\leq \left\| \mathbb{I}_n \widehat{\mathbf{K}} - \mathbb{I}_n \mathbb{P}_n K \right\|_{L^\infty,1([0,1]^2)} + \left\| \mathbb{I}_n \mathbb{P}_n K - K \right\|_{L^\infty,1([0,1]^2)} \\
&= \left\| (\mathbb{I}_n \mathbb{P}_n K - \rho_n^{-1})_+ \right\|_{L^\infty,1([0,1]^2)} + \left\| \mathbb{I}_n \mathbb{P}_n K - K \right\|_{L^\infty,1([0,1]^2)} \\
&\leq \left\| (K - \rho_n^{-1})_+ \right\|_{L^\infty,1([0,1]^2)} + \left\| (\mathbb{I}_n \mathbb{P}_n K - K)_+ \right\|_{L^\infty,1([0,1]^2)} \\
&\quad + \left\| \mathbb{I}_n \mathbb{P}_n K - K \right\|_{L^\infty,1([0,1]^2)} \\
&\leq \left\| (K - \rho_n^{-1})_+ \right\|_{L^\infty,1([0,1]^2)} + 2 \left\| \mathbb{I}_n \mathbb{P}_n K - K \right\|_{L^\infty,1([0,1]^2)}.
\end{aligned}$$

Since $\rho_n \rightarrow 0$ and in view of [\(H_w.2\)](#), the right-hand side in the above display goes to 0 as $n \rightarrow +\infty$. Combined with the first claim we obtain the desired conclusion. \square

4. WELL-POSEDNESS

4.1. The case $p \in]1, +\infty[$

To lighten notation, for $1 < p < +\infty$, we define the function

$$\Psi : x \in \mathbb{R} \mapsto |x|^{p-2} x = \text{sign}(x)|x|^{p-1},$$

where we take $\text{sign}(0) = 0$.

The next lemma summarizes key monotonicity and continuity properties of Ψ which will be instrumental to us.

Lemma 4.1. (i) *Monotonicity: assume that the constant β satisfies $\beta \in [\max(p, 2), +\infty[$. Then for all $x, y \in \mathbb{R}$,*

$$(\Psi(y) - \Psi(x))(y - x) \geq C_1 |y - x|^\beta (|y| + |x|)^{p-\beta}, \quad (9)$$

where the constant C_1 is sharp and given by

$$C_1 = 2^{2-p} \min(1, p - 1). \quad (10)$$

In particular,

$$(\Psi(y) - \Psi(x))(y - x) \geq C_1 \begin{cases} |y - x|^p & p \in [2, +\infty[, \\ |y - x|^2 (|y| + |x|)^{p-2} & p \in]1, 2]. \end{cases} \quad (11)$$

(ii) *Continuity: assume that the constant α satisfies $\alpha \in [0, \min(1, p - 1)]$. Then for all $x, y \in \mathbb{R}$,*

$$|\Psi(y) - \Psi(x)| \leq C_2 |y - x|^\alpha (|y| + |x|)^{p-1-\alpha}, \quad (12)$$

where the constant C_2 is sharp and given by

$$C_2 = \max(2^{2-p}, (p - 1)2^{2-p}, 1). \quad (13)$$

In particular,

$$|\Psi(y) - \Psi(x)| \leq C_2 \begin{cases} |y - x| (|y| + |x|)^{p-2} & p \in [2, +\infty[, \\ |y - x|^{p-1} & p \in]1, 2], \end{cases} \quad (14)$$

Proof. (i) For [\(9\)](#), see Theorem 2.2 of [\[17\]](#). For [\(11\)](#), set $\beta = p$ for $p \geq 2$ and $\beta = 2$ otherwise in [\(9\)](#); see also the seminal results of Lemmas 5.1 and 5.2 from [\[29\]](#).

(ii) For [\(12\)](#), see Theorem 2.1 of [\[17\]](#). For [\(14\)](#), set $\alpha = 1$ for $p \geq 2$ and $\alpha = p - 1$ otherwise in [\(12\)](#); see also the seminal results of Lemmas 5.3 and 5.4 from [\[29\]](#). \square

We now collect some preliminary properties of the nonlocal p -Laplacian, an operator on $L^1(\Omega)$ that we denote for short as

$$\Delta_p^K u(\mathbf{x}, t) = - \int_{\Omega} K(\mathbf{x}, \mathbf{y}) |u(\mathbf{y}, t) - u(\mathbf{x}, t)|^{p-2} (u(\mathbf{y}, t) - u(\mathbf{x}, t)) \, d\mathbf{y}.$$

Proposition 4.1. *Assume that K satisfies (H.1)–(H.3).*

- (i) Δ_p^K is positively homogeneous of degree $p - 1$.
- (ii) If $p > 2$, $L^{p-1}(\Omega) \subset \text{dom}(\Delta_p^K)$.
- (iii) If $1 < p \leq 2$, $\text{dom}(\Delta_p^K) = L^1(\Omega)$ and Δ_p^K is closed in $L^1(\Omega) \times L^1(\Omega)$.
- (iv) Let $h : \mathbb{R} \rightarrow \mathbb{R}$. Then for every $u, v \in L^p(\Omega)$,

$$\begin{aligned} 0 &\leq \int_{\Omega} (\Delta_p^K u(\mathbf{x}) - \Delta_p^K v(\mathbf{x})) h(u(\mathbf{x}) - v(\mathbf{x})) \, d\mathbf{x} \\ &= \frac{1}{2} \int_{\Omega^2} K(\mathbf{x}, \mathbf{y}) (\Psi(u(\mathbf{y}) - u(\mathbf{x})) - \Psi(v(\mathbf{y}) - v(\mathbf{x}))) (h(u(\mathbf{y}) - v(\mathbf{y})) - h(u(\mathbf{x}) - v(\mathbf{x}))) \, d\mathbf{y} \, d\mathbf{x}. \end{aligned} \tag{15}$$

If h is bounded, then this holds for any $u, v \in \text{dom}(\Delta_p^K)$.

- (v) For every $u, v \in L^p(\Omega)$,

$$\begin{aligned} &\int_{\Omega} (\Delta_p^K u(\mathbf{x}) - \Delta_p^K v(\mathbf{x})) (u(\mathbf{x}) - v(\mathbf{x})) \, d\mathbf{x} \\ &\geq \frac{C}{2} \left(\int_{\Omega^2} K(\mathbf{x}, \mathbf{y}) |u(\mathbf{y}) - u(\mathbf{x}) - (v(\mathbf{y}) - v(\mathbf{x}))|^p \, d\mathbf{y} \, d\mathbf{x} \right)^{\max(1, 2/p)} \end{aligned}$$

where

$$C = \begin{cases} C_1 & p \in [2, +\infty[, \\ 2^{2p-5} C_1 \|K\|_{L^{\infty,1}(\Omega^2)}^{1-2/p} (\|u\|_{L^p(\Omega)} + \|v\|_{L^p(\Omega)})^{p-2} & p \in]1, 2[. \end{cases}$$

and C_1 is the constant in (10). If $u, v \in L^\infty(\Omega)$, then

$$\begin{aligned} &\int_{\Omega} (\Delta_p^K u(\mathbf{x}) - \Delta_p^K v(\mathbf{x})) (u(\mathbf{x}) - v(\mathbf{x})) \, d\mathbf{x} \\ &\geq \frac{C}{2} \left(\int_{\Omega^2} K(\mathbf{x}, \mathbf{y}) |u(\mathbf{y}) - u(\mathbf{x}) - (v(\mathbf{y}) - v(\mathbf{x}))|^2 \, d\mathbf{y} \, d\mathbf{x} \right)^{\max(1, p/2)}, \end{aligned}$$

where

$$C = \begin{cases} C_1 & p \in [2, +\infty[, \\ 2^{p-2} C_1 (\|u\|_{L^\infty(\Omega)} + \|v\|_{L^\infty(\Omega)})^{p-2} & p \in]1, 2[. \end{cases}$$

- (vi) For $p \in]1, 2]$ and every $u, v \in L^2(\Omega)$,

$$\int_{\Omega} (\Delta_p^K u(\mathbf{x}) - \Delta_p^K v(\mathbf{x})) (u(\mathbf{x}) - v(\mathbf{x})) \, d\mathbf{x} \geq C \|\Delta_p^K u - \Delta_p^K v\|_{L^2(\Omega)}^{p/(p-1)}$$

where

$$C = 2^{\frac{p-2}{2(p-1)}} \left(C_2^{1/2} \|K\|_{L^{\infty,1}(\Omega^2)} \right)^{\frac{1}{1-p}} (1 - 1/p), \text{ and } C_2 \text{ is the constant in (13).}$$

- (vii) For $p \in]1, +\infty[$, Δ_p^K is completely accretive and satisfies the range condition

$$L^p(\Omega) \subset \text{ran}(\mathbf{I} + \Delta_p^K). \tag{16}$$

Consequently, the resolvent $J_{\lambda \Delta_p^K} \stackrel{\text{def}}{=} (\mathbf{I} + \lambda \Delta_p^K)^{-1}$, $\lambda > 0$, is single-valued on $L^p(\Omega)$ and non-expansive in $L^q(\Omega)$ for all $q \in [1, +\infty]$.

Proof. (i)–(iii) follow from Remark 2.2 of [3] which still holds for our larger class of kernels K .

For (iv), see Lemma A.2 of [30]. Monotonicity is immediate since h is non-decreasing.

The proof of (vii) is the same as that of Theorem 2.4 from [3], where we invoke the monotonicity claim (i).

We now show (v)². The case $p \in [2, +\infty[$ is immediate by inserting Lemma 4.1(i) into (15) with $h(x) = x$. For $p \in]1, 2]$, to lighten notation, denote the nonlocal gradient $\nabla^{\text{NL}}u(\mathbf{x}, \mathbf{y}) = u(\mathbf{y}) - u(\mathbf{x})$. We then have by Lemma 4.1(i) that

$$C_1 |\nabla^{\text{NL}}(u - v)(\mathbf{x}, \mathbf{y})|^2 \leq (\Psi(\nabla^{\text{NL}}u(\mathbf{x}, \mathbf{y})) - \Psi(\nabla^{\text{NL}}v(\mathbf{x}, \mathbf{y}))) \times (\nabla^{\text{NL}}u(\mathbf{x}, \mathbf{y}) - \nabla^{\text{NL}}v(\mathbf{x}, \mathbf{y})) (|\nabla^{\text{NL}}u(\mathbf{x}, \mathbf{y})| + |\nabla^{\text{NL}}v(\mathbf{x}, \mathbf{y})|)^{2-p}. \tag{17}$$

Taking the power $p/2$, multiplying by K and integrating, we get

$$\begin{aligned} & C_1^{p/2} \int_{\Omega^2} K(\mathbf{x}, \mathbf{y}) |\nabla^{\text{NL}}(u - v)(\mathbf{x}, \mathbf{y})|^p \, d\mathbf{x} \, d\mathbf{y} \\ & \leq \int_{\Omega^2} (K(\mathbf{x}, \mathbf{y}) (\Psi(\nabla^{\text{NL}}u(\mathbf{x}, \mathbf{y})) - \Psi(\nabla^{\text{NL}}v(\mathbf{x}, \mathbf{y}))) (\nabla^{\text{NL}}u(\mathbf{x}, \mathbf{y}) - \nabla^{\text{NL}}v(\mathbf{x}, \mathbf{y})))^{p/2} \\ & \quad \times \left(K(\mathbf{x}, \mathbf{y})^{1/p} (|\nabla^{\text{NL}}u(\mathbf{x}, \mathbf{y})| + |\nabla^{\text{NL}}v(\mathbf{x}, \mathbf{y})|) \right)^{(2-p)p/2} \, d\mathbf{x} \, d\mathbf{y}. \end{aligned}$$

It is easily seen that

$$\begin{aligned} & (K \cdot (\Psi(\nabla^{\text{NL}}u) - \Psi(\nabla^{\text{NL}}v)) (\nabla^{\text{NL}}u - \nabla^{\text{NL}}v))^{p/2} \in L^{2/p}(\Omega^2) \\ & \left(K^{1/p} \cdot (|\nabla^{\text{NL}}u| + |\nabla^{\text{NL}}v|) \right)^{(2-p)p/2} \in L^{2/(2-p)}(\Omega^2). \end{aligned}$$

It then follows from Hölder inequality and (15) that

$$\begin{aligned} C_1^{p/2} \int_{\Omega^2} K(\mathbf{x}, \mathbf{y}) |\nabla^{\text{NL}}(u - v)(\mathbf{x}, \mathbf{y})|^p \, d\mathbf{x} \, d\mathbf{y} & \leq 2 \left(\int_{\Omega} (\Delta_p^K u(\mathbf{x}) - \Delta_p^K v(\mathbf{x})) (u(\mathbf{x}) - v(\mathbf{x})) \, d\mathbf{x} \right)^{p/2} \\ & \quad \cdot \left(\int_{\Omega^2} K(\mathbf{x}, \mathbf{y}) (|\nabla^{\text{NL}}u(\mathbf{x}, \mathbf{y})| + |\nabla^{\text{NL}}v(\mathbf{x}, \mathbf{y})|)^p \, d\mathbf{x} \, d\mathbf{y} \right)^{(2-p)/2}. \end{aligned}$$

We have by Jensen’s inequality

$$\begin{aligned} & \int_{\Omega^2} K(\mathbf{x}, \mathbf{y}) (|\nabla^{\text{NL}}u(\mathbf{x}, \mathbf{y})| + |\nabla^{\text{NL}}v(\mathbf{x}, \mathbf{y})|)^p \, d\mathbf{x} \, d\mathbf{y} \\ & \leq 4^{p-1} \int_{\Omega^2} K(\mathbf{x}, \mathbf{y}) (|u(\mathbf{x})|^p + |u(\mathbf{y})|^p + |v(\mathbf{x})|^p + |v(\mathbf{y})|^p) \, d\mathbf{x} \, d\mathbf{y} \\ & \leq 2^{2p-1} \|K\|_{L^{\infty,1}(\Omega^2)} \left(\|u\|_{L^p(\Omega)}^p + \|v\|_{L^p(\Omega)}^p \right), \end{aligned}$$

whence we obtain

$$\begin{aligned} & C_1 \left(\int_{\Omega^2} K(\mathbf{x}, \mathbf{y}) |\nabla^{\text{NL}}(u - v)(\mathbf{x}, \mathbf{y})|^p \, d\mathbf{x} \, d\mathbf{y} \right)^{2/p} \\ & \leq 2^{5-2p} \left(\int_{\Omega} (\Delta_p^K u(\mathbf{x}) - \Delta_p^K v(\mathbf{x})) (u(\mathbf{x}) - v(\mathbf{x})) \, d\mathbf{x} \right) \\ & \quad \times \|K\|_{L^{\infty,1}(\Omega^2)}^{(2-p)/p} \left(\|u\|_{L^p(\Omega)} + \|v\|_{L^p(\Omega)} \right)^{2-p}. \end{aligned}$$

²This can be seen as a nonlocal analogue of Propositions 5.1 and 5.2 from [29].

Rearranging proves the bound. For $u, v \in L^\infty(\Omega)$ and $p \in [2, +\infty]$ we use that $L^p(\Omega) \subset L^2(\Omega)$. For $p \in]1, 2]$, we embark from (17) and use that for all $(\mathbf{x}, \mathbf{y}) \in \Omega^2$,

$$|\nabla^{\text{NL}}u(\mathbf{x}, \mathbf{y})| + |\nabla^{\text{NL}}v(\mathbf{x}, \mathbf{y})| \leq 2\left(\|u\|_{L^\infty(\Omega)} + \|v\|_{L^\infty(\Omega)}\right).$$

Multiplying (17) by K , integrating and using (15), we conclude.

To prove (vi), we start by showing that Δ_p^K is Hölder continuous with exponent $p - 1$ on $L^2(\Omega)$. We have by Jensen inequality (twice) and (14),

$$\begin{aligned} \|\Delta_p^K u - \Delta_p^K v\|_{L^2(\Omega)}^2 &= \int_{\Omega} \left| \int_{\Omega} K(\mathbf{x}, \mathbf{y}) (\Psi(\nabla^{\text{NL}}u(\mathbf{x}, \mathbf{y})) - \Psi(\nabla^{\text{NL}}v(\mathbf{x}, \mathbf{y}))) \, d\mathbf{y} \right|^2 \, d\mathbf{x} \\ &\leq \|K\|_{L^\infty,1(\Omega^2)} \int_{\Omega^2} K(\mathbf{x}, \mathbf{y}) (\Psi(\nabla^{\text{NL}}u(\mathbf{x}, \mathbf{y})) - \Psi(\nabla^{\text{NL}}v(\mathbf{x}, \mathbf{y})))^2 \, d\mathbf{x} \, d\mathbf{y} \\ &\leq C_2 \|K\|_{L^\infty,1(\Omega^2)} \int_{\Omega^2} K(\mathbf{x}, \mathbf{y}) (\nabla^{\text{NL}}(u - v)(\mathbf{x}, \mathbf{y}))^{2(p-1)} \, d\mathbf{x} \, d\mathbf{y} \\ &\leq 2^p C_2 \|K\|_{L^\infty,1(\Omega^2)} \int_{\Omega^2} K(\mathbf{x}, \mathbf{y}) (u(\mathbf{x}) - v(\mathbf{x}))^{2(p-1)} \, d\mathbf{x} \, d\mathbf{y} \\ &\leq 2^p C_2 \|K\|_{L^\infty,1(\Omega^2)}^2 \left(\int_{\Omega} (u(\mathbf{x}) - v(\mathbf{x}))^2 \, d\mathbf{x} \right)^{p-1} \\ &= 2^p C_2 \|K\|_{L^\infty,1(\Omega^2)}^2 \|u - v\|_{L^2(\Omega)}^{2(p-1)}. \end{aligned} \tag{18}$$

We are now in position to invoke ([8], Cor. 18.14(i) \Rightarrow (v)) to show that the claimed inequality holds. □

Solutions of (P) will be understood in the following sense:

Definition 4.1. Let $p \in]1, +\infty[$. A solution of (P) in $[0, T]$ is a function

$$u \in C([0, T]; L^1(\Omega)) \cap W^{1,1}([0, T]; L^1(\Omega)),$$

that satisfies $u(\mathbf{x}, 0) = g(\mathbf{x})$ a.e. $\mathbf{x} \in \Omega$ and

$$\frac{\partial}{\partial t} u(\mathbf{x}, t) = -\Delta_p^K u(\mathbf{x}, t) + f(\mathbf{x}, t) \quad \text{a.e. in } \Omega \times]0, T[.$$

Such a solution is also a strong solution (see [4], Def. A.3).

The main result of existence and uniqueness of a global solution, that is, a solution on $[0, T]$ for $T > 0$ is stated in the following theorem.

Theorem 4.1. Suppose that $p \in]1, +\infty[$ and assumptions (H.1)–(H.3) hold. Let $g \in L^p(\Omega)$ and $f \in L^1([0, T]; L^p(\Omega))$.

- (i) For any $T > 0$, there exists a unique strong solution in $[0, T]$ of (P).
- (ii) Moreover, for $q \in [1, +\infty]$, if $g_i \in L^q(\Omega)$ and $f_i \in L^1([0, T]; L^q(\Omega))$, $i = 1, 2$, and u_i is the solution of (P) with data (f_i, g_i) , then

$$\|u_1(\cdot, t) - u_2(\cdot, t)\|_{L^q(\Omega)} \leq \|g_1 - g_2\|_{L^q(\Omega)} + \|f_1 - f_2\|_{L^1([0, T]; L^q(\Omega))}, \quad \forall t \in [0, T]. \tag{19}$$

Proof. The proof follows the same lines as that of Theorem 1.2 from [3] extended to the case where $f \not\equiv 0$ thanks to the results of [9], where we invoke Proposition 4.1(ii), (iii) and (vii). □

Remark 4.1. In [3] (see also [4], Chap. 6), the authors impose the following stringent assumptions: $K(\mathbf{x}, \mathbf{y}) = J(\mathbf{x} - \mathbf{y})$, where J is nonnegative, continuous, radially symmetric, compactly supported, $J(0) > 0$ and $\int_{\mathbb{R}^d} J(\mathbf{x}) d\mathbf{x} < +\infty$. Actually, these assumptions are not needed for existence and uniqueness. The particular form $J(\mathbf{x} - \mathbf{y})$ of the kernel is not needed. Continuity with radial symmetry and support compactness play a pivotal role to study convergence to the local p -Laplacian problem in Theorem 1.5 from [3]. In addition, $J(0) > 0$ was mandatory to prove a Poincaré-type inequality in Proposition 4.1 from [3]. Even for the form $J(\mathbf{x} - \mathbf{y})$, our assumptions **(H'.1)**–**(H'.3)** are weaker than those of [3]. This discussion remains true also for the case $p = 1$.

4.2. The case $p = 1$

We will need to define subdifferential of the absolute value function on \mathbb{R} , which is the well-known set-valued mapping $\partial|\cdot| : \mathbb{R} \rightarrow 2^{\mathbb{R}}$,

$$\partial|\cdot|(x) = \begin{cases} 1 & x > 0 \\ [-1, 1] & x = 0 \\ -1 & x < 0. \end{cases}$$

It will be convenient to denote the 1-Laplacian Δ_1^K . This is a set-valued operator in $L^1(\Omega) \times L^1(\Omega)$ such that $\eta \in \Delta_1^K u$ if and only if

$$\eta(\mathbf{x}) = - \int_{\Omega} K(\mathbf{x}, \mathbf{y}) w(\mathbf{x}, \mathbf{y}) d\mathbf{y} \quad \text{a.e. in } \Omega,$$

for a subgradient function w satisfying $\|w\|_{L^\infty(\Omega^2)} \leq 1$, $w(\mathbf{x}, \mathbf{y}) = -w(\mathbf{y}, \mathbf{x})$, and

$$w(\mathbf{x}, \mathbf{y}) \in \partial|\cdot|(u(\mathbf{y}) - u(\mathbf{x})).$$

Solutions of **(P)** will be understood in the following sense.

Definition 4.2. A solution of **(P)** for $p = 1$ in $[0, T]$ is a function

$$u \in C([0, T]; L^1(\Omega)) \cap W^{1,1}([0, T[; L^1(\Omega)),$$

that satisfies $u(\mathbf{x}, 0) = g(\mathbf{x})$ for a.e. $\mathbf{x} \in \Omega$ and

$$\frac{\partial}{\partial t} u(\mathbf{x}, t) = -\eta(\mathbf{x}, t) + f(\mathbf{x}, t) \quad \text{a.e. in } \Omega \times]0, T[,$$

where $\eta(\cdot, t) \in \Delta_1^K u(\cdot, t)$.

Observe that for $p = 1$, the evolution problem **(P)** reads

$$\begin{cases} \frac{\partial}{\partial t} u(\mathbf{x}, t) = \int_{\Omega} K(\mathbf{x}, \mathbf{y}) \text{sign}(u(\mathbf{y}, t) - u(\mathbf{x}, t)) d\mathbf{y} + f(\mathbf{x}, t), & \mathbf{x} \in \Omega, t > 0, \\ u(\mathbf{x}, 0) = g(\mathbf{x}), & \mathbf{x} \in \Omega, \end{cases}$$

where

$$\text{sign}(x) = \begin{cases} \frac{x}{|x|} & x \neq 0 \\ 0 & x = 0. \end{cases}$$

Thus, it satisfies

$$\frac{\partial}{\partial t} u(\cdot, t) \in -\Delta_1^K u(\cdot, t).$$

In the same vein as Proposition 4.1, the 1-Laplacian enjoys the following properties.

Proposition 4.2. Assume that K satisfies **(H.1)**–**(H.3)**.

- (i) $\text{dom}(\Delta_1^K) = L^1(\Omega)$ and (the graph of) Δ_1^K is closed in $L^1(\Omega) \times L^1(\Omega)$.
(ii) Let $h \in C^1(\mathbb{R})$ be a nondecreasing function. Then for every $u_i \in L^1(\Omega)$ and any $\eta_i \in \Delta_1^K u_i$, $i = 1, 2$,

$$\begin{aligned} 0 &\leq \int_{\Omega} (\eta_1(\mathbf{x}) - \eta_2(\mathbf{x}))(h(u_1(\mathbf{x}) - u_2(\mathbf{x}))) \, d\mathbf{x} \\ &= \frac{1}{2} \int_{\Omega^2} K(\mathbf{x}, \mathbf{y})(w_1(\mathbf{x}, \mathbf{y}) - w_2(\mathbf{x}, \mathbf{y}))(h(u_1(\mathbf{y}) - u_2(\mathbf{y})) - h(u_1(\mathbf{x}) - u_2(\mathbf{x}))) \, d\mathbf{x} \, d\mathbf{y} \end{aligned} \quad (20)$$

where w_i are the corresponding subgradient functions defined above. In particular,

$$\int_{\Omega^2} K(\mathbf{x}, \mathbf{y})w_i(\mathbf{x}, \mathbf{y})u_i(\mathbf{x}) \, d\mathbf{x} \, d\mathbf{y} = -\frac{1}{2} \int_{\Omega^2} K(\mathbf{x}, \mathbf{y})|u_i(\mathbf{y}) - u_i(\mathbf{x})| \, d\mathbf{x} \, d\mathbf{y}.$$

- (iii) Δ_1^K is completely accretive and satisfies the range condition

$$L^\infty(\Omega) \subset \text{ran}(\mathbb{I} + \Delta_1^K). \quad (21)$$

Proof. For (i), see Remark 2.8 of [3] which still holds for our class of kernels K .

The proof of (iii) is again the same as that of Theorem 2.9 from [3], where we invoke the monotonicity claim (ii) to which we turn now.

For any $v \in L^1(\Omega)$, we have the integration by parts formula

$$\begin{aligned} \int_{\Omega^2} K(\mathbf{x}, \mathbf{y})w_i(\mathbf{x}, \mathbf{y})(v(\mathbf{y}) - v(\mathbf{x})) \, d\mathbf{x} \, d\mathbf{y} &= - \int_{\Omega^2} K(\mathbf{y}, \mathbf{x})w_i(\mathbf{y}, \mathbf{x})v(\mathbf{y}) \, d\mathbf{y} \, d\mathbf{x} \\ &\quad - \int_{\Omega^2} K(\mathbf{x}, \mathbf{y})w_i(\mathbf{x}, \mathbf{y})v(\mathbf{x}) \, d\mathbf{x} \, d\mathbf{y} \\ &= -2 \int_{\Omega^2} K(\mathbf{x}, \mathbf{y})w_i(\mathbf{x}, \mathbf{y})v(\mathbf{x}) \, d\mathbf{x} \, d\mathbf{y}. \end{aligned} \quad (22)$$

Taking $v(\mathbf{x}) = h(u_1(\mathbf{x}) - u_2(\mathbf{x}))$ in (22) with w_1 and w_2 , and then taking the difference, we arrive at

$$\begin{aligned} &-2 \int_{\Omega} \left(\int_{\Omega} K(\mathbf{x}, \mathbf{y})(w_1(\mathbf{x}, \mathbf{y}) - w_2(\mathbf{x}, \mathbf{y})) \, d\mathbf{y} \right) h(u_1(\mathbf{x}) - u_2(\mathbf{x})) \, d\mathbf{x} \\ &= 2 \int_{\Omega} (\eta_1(\mathbf{x}) - \eta_2(\mathbf{x}))(h(u_1(\mathbf{x}) - u_2(\mathbf{x}))) \, d\mathbf{x} \\ &= \int_{\Omega^2} K(\mathbf{x}, \mathbf{y})(w_1(\mathbf{x}, \mathbf{y}) - w_2(\mathbf{x}, \mathbf{y}))(h(u_1(\mathbf{y}) - u_2(\mathbf{y})) - h(u_1(\mathbf{x}) - u_2(\mathbf{x}))) \, d\mathbf{x} \, d\mathbf{y}. \end{aligned}$$

By the mean-value theorem applied to h , we get

$$\begin{aligned} &= 2 \int_{\Omega} (\eta_1(\mathbf{x}) - \eta_2(\mathbf{x}))(h(u_1(\mathbf{x}) - u_2(\mathbf{x}))) \, d\mathbf{x} \\ &= \int_{\Omega^2} K(\mathbf{x}, \mathbf{y})(w_1(\mathbf{x}, \mathbf{y}) - w_2(\mathbf{x}, \mathbf{y}))h'(\zeta(\mathbf{x}, \mathbf{y}))((u_1(\mathbf{y}) - u_2(\mathbf{y})) - (u_1(\mathbf{x}) - u_2(\mathbf{x}))) \, d\mathbf{x} \, d\mathbf{y} \\ &= \int_{\Omega^2} K(\mathbf{x}, \mathbf{y})h'(\zeta(\mathbf{x}, \mathbf{y}))(w_1(\mathbf{x}, \mathbf{y}) - w_2(\mathbf{x}, \mathbf{y}))((u_1(\mathbf{y}) - u_1(\mathbf{x})) - (u_2(\mathbf{y}) - u_2(\mathbf{x}))) \, d\mathbf{x} \, d\mathbf{y}, \end{aligned}$$

where $\zeta(\mathbf{x}, \mathbf{y})$ is an intermediate value between $u_1(\mathbf{y}) - u_2(\mathbf{y})$ and $u_1(\mathbf{x}) - u_2(\mathbf{x})$. Since h is increasing, that $w_i(\mathbf{x}, \mathbf{y}) \in \partial|\cdot|(u_i(\mathbf{y}) - u_i(\mathbf{x}))$, and $\partial|\cdot|$ is a monotone operator, we get the claimed monotonicity.

To get the particular identity, we specialize (22) by taking $v = u_i$, which entails

$$- \int_{\Omega^2} K(\mathbf{x}, \mathbf{y})w_i(\mathbf{x}, \mathbf{y})(u_i(\mathbf{y}) - u_i(\mathbf{x})) \, d\mathbf{x} \, d\mathbf{y} = 2 \int_{\Omega^2} K(\mathbf{x}, \mathbf{y})w_i(\mathbf{x}, \mathbf{y})u_i(\mathbf{x}) \, d\mathbf{x} \, d\mathbf{y}.$$

We finally use the equivalent characterization of $\partial|\cdot|$, which originates from the Fenchel's identity since $|\cdot|$ is positively homogeneous,

$$\partial|\cdot|(x) = \{\xi \in \mathbb{R} : |\xi| \leq 1 \text{ and } \xi x = |x|\}.$$

Applying this identity with $x = u_i(\mathbf{y}) - u_i(\mathbf{x})$ and $\xi = w_i(\mathbf{x}, \mathbf{y})$ gives the claim. \square

Theorem 4.2. *Suppose that $p = 1$, and assumptions (H.1)–(H.3) hold. Let $g \in L^1(\Omega)$ and $f \in L^1([0, T]; L^1(\Omega))$. For any $T > 0$, there exists a unique solution in $[0, T]$ of (P) in the sense of Definition 4.2.*

Proof. The proof is an adaptation of Theorem 1.4 from [3] to the case where $f \not\equiv 0$ thanks to the results of [9], where we invoke Proposition 4.2(i) and (iii). \square

5. CONTINUOUS-CONTINUOUS ESTIMATES

In this section, we provide an estimate that compares solutions of two p -Laplacian evolution problems of the form (P) with two different kernels and initial data. This estimate will be instrumental to derive error bounds in the totally discrete case.

5.1. The case $p \in]1, +\infty[$

We have the following error bounds and convergence result.

Theorem 5.1. *Suppose that $p \in]1, +\infty[$. Let u be a solution of (P) with kernel K and data (f, g) . Let u_n be a sequence of solutions to (P) with kernels K_n and data (f_n, g_n) . Assume that K and K_n satisfy (H.1), (H.2) and $K, K_n \in L^{\infty, 2}(\Omega^2)$, and that either one of the following holds:*

- (a) $p \in]1, 2[$, $g, g_n \in L^2(\Omega)$, and $f, f_n \in L^1([0, T]; L^2(\Omega))$;
- (b) $p \geq 2$, $g, g_n \in L^{2(p-1)}(\Omega)$ and $f, f_n \in L^1([0, T]; L^{2(p-1)}(\Omega))$;
- (c) $g, g_n \in L^\infty(\Omega)$ and $f, f_n \in L^1([0, T]; L^\infty(\Omega))$.

Then, the following hold.

- (i) u and u_n are the unique solutions (P) with respectively data (f, g) and (f_n, g_n) .
- (ii) We have the error estimate

$$\begin{aligned} \|u_n - u\|_{C([0, T]; L^2(\Omega))} &\leq \|g_n - g\|_{L^2(\Omega)} + \|f_n - f\|_{L^1([0, T]; L^2(\Omega))} \\ &\quad + CT \begin{cases} \|K_n - K\|_{L^{\infty, 2}(\Omega^2)}, & \text{under (a) or (b)} \\ \|K_n - K\|_{L^2(\Omega^2)}, & \text{under (c)} \end{cases} \end{aligned} \quad (23)$$

where C is positive constant that may depend only on p, g and f .

- (iii) Moreover, if (c) holds, $\sup_{n \in \mathbb{N}} |g_n(\mathbf{x})| < +\infty$ a.e. on Ω and $g_n \rightarrow g$ pointwise a.e. on Ω , $\sup_{n \in \mathbb{N}} |f_n(\mathbf{x}, t)| < +\infty$ a.e. on $\Omega \times [0, T]$ and $f_n \rightarrow f$ pointwise a.e. on $\Omega \times [0, T]$, and the sequence $\{|K_n|^2\}_{n \in \mathbb{N}}$ is uniformly integrable over Ω^2 and $K_n \rightarrow K$ pointwise a.e. on Ω^2 . Then

$$\lim_{n \rightarrow +\infty} \|u_n - u\|_{C([0, T]; L^2(\Omega))} = 0.$$

Remark 5.1. Observe that since $L^\infty(\Omega) \subset L^2(\Omega)$ and $L^{2(p-1)}(\Omega) \subset L^2(\Omega)$ for $p \geq 2$, then the first two terms involved in (23) provide a non-trivial bound. Similarly, since $L^{\infty, 2}(\Omega^2) \subset L^2(\Omega^2)$, the last term in the bound for case (c) is also non-trivial. In fact, both bounds in (23) can be summarized in one bound; the first one. However, the second bound for case (c) is obviously sharper.

Proof. In the proof, C is any positive constant that may depend solely on p and g .

- (i) Since $L^{\infty,2}(\Omega^2) \subset L^{\infty,1}(\Omega^2)$, assumption **(H.3)** holds for both K and K_n . We also have the embeddings
 - $L^2(\Omega) \subset L^p(\Omega)$ under **(a)**,
 - $L^{2(p-1)}(\Omega) \subset L^p(\Omega)$ under **(b)**, and
 - $L^\infty(\Omega) \subset L^p(\Omega)$ under **(c)**.

Thus $g, g_n \in L^p(\Omega)$ and $f, f_n \in L^1([0, T]; L^p(\Omega))$. Existence and uniqueness of the solutions u and u_n in the sense of Definition 4.1 is a consequence of Theorem 4.1.

- (ii) Denote the error function $\xi_n(\mathbf{x}, t) = u_n(\mathbf{x}, t) - u(\mathbf{x}, t)$, then from **(P)**, we have a.e.

$$\begin{aligned} \frac{\partial \xi_n(\mathbf{x}, t)}{\partial t} &= -(\Delta_p^{K_n}(u_n(\mathbf{x}, t)) - \Delta_p^K(u(\mathbf{x}, t))) + f_n(\mathbf{x}, t) - f(\mathbf{x}, t) \\ &= -(\Delta_p^{K_n}(u_n(\mathbf{x}, t)) - \Delta_p^{K_n}(u(\mathbf{x}, t))) - (\Delta_p^{K_n}(u(\mathbf{x}, t)) - \Delta_p^K(u(\mathbf{x}, t))) \\ &\quad + f_n(\mathbf{x}, t) - f(\mathbf{x}, t). \end{aligned} \tag{24}$$

Multiplying both sides of (24) by $\xi_n(\mathbf{x}, t)$ and integrating, we get

$$\begin{aligned} \frac{1}{2} \frac{\partial}{\partial t} \|\xi_n(\cdot, t)\|_{L^2(\Omega)}^2 &= - \int_{\Omega} (\Delta_p^{K_n} u_n(\mathbf{x}, t) - \Delta_p^{K_n} u(\mathbf{x}, t))(u_n(\mathbf{x}, t) - u(\mathbf{x}, t)) \, d\mathbf{x} \\ &\quad + \int_{\Omega^2} (K_n(\mathbf{x}, \mathbf{y}) - K(\mathbf{x}, \mathbf{y}))\Psi(u(\mathbf{y}, t) - u(\mathbf{x}, t))\xi_n(\mathbf{x}, t) \, d\mathbf{x} \, d\mathbf{y} \\ &\quad + \int_{\Omega} (f_n(\mathbf{x}, t) - f(\mathbf{x}, t))\xi_n(\mathbf{x}, t) \, d\mathbf{x}. \end{aligned} \tag{25}$$

Since $g, g_n \in L^p(\Omega)$ and $f, f_n \in L^1([0, T]; L^p(\Omega))$, $u_n(\cdot, t), u(\cdot, t) \in L^p(\Omega)$ for any $t \in [0, T]$ thanks to (19). We can then apply Proposition 4.1(iv) with $h(x) = x$ to assert that the first term on the right-hand side of (25) is nonpositive. Let us now bound the second term.

- Case **(c)**: in this case $\|u\|_{C([0,T];L^\infty(\Omega))} \leq \|g\|_{L^\infty(\Omega)} + \|f\|_{L^1([0,T];L^\infty(\Omega))}$ thanks to (19), and we get from Cauchy–Schwartz inequality that

$$\begin{aligned} &\left| \int_{\Omega^2} (K_n(\mathbf{x}, \mathbf{y}) - K(\mathbf{x}, \mathbf{y}))\Psi(u(\mathbf{y}, t) - u(\mathbf{x}, t))\xi_n(\mathbf{x}, t) \, d\mathbf{x} \, d\mathbf{y} \right| \\ &\leq 2^{p-1} \|u(\cdot, t)\|_{L^\infty(\Omega)}^{p-1} \int_{\Omega^2} |K_n(\mathbf{x}, \mathbf{y}) - K(\mathbf{x}, \mathbf{y})| |\xi_n(\mathbf{x}, t)| \, d\mathbf{x} \, d\mathbf{y} \\ &\leq 2^{p-1} \left(\|g\|_{L^\infty(\Omega)} + \|f\|_{L^1([0,T];L^\infty(\Omega))} \right)^{p-1} \|K_n - K\|_{L^2(\Omega^2)} \|\xi_n(\cdot, t)\|_{L^2(\Omega)} \\ &= C \|K_n - K\|_{L^2(\Omega^2)} \|\xi_n(\cdot, t)\|_{L^2(\Omega)}. \end{aligned} \tag{26}$$

- Case **(a)** or **(b)**: applying again Cauchy–Schwartz inequality we obtain

$$\begin{aligned} &\left| \int_{\Omega^2} (K_n(\mathbf{x}, \mathbf{y}) - K(\mathbf{x}, \mathbf{y}))\Psi(u(\mathbf{y}, t) - u(\mathbf{x}, t))\xi_n(\mathbf{x}, t) \, d\mathbf{x} \, d\mathbf{y} \right| \\ &\leq \left(\int_{\Omega^2} |u(\mathbf{y}, t) - u(\mathbf{x}, t)|^{2(p-1)} \, d\mathbf{x} \, d\mathbf{y} \right)^{1/2} \left(\int_{\Omega^2} |K_n(\mathbf{x}, \mathbf{y}) - K(\mathbf{x}, \mathbf{y})|^2 |\xi_n(\mathbf{x}, t)|^2 \, d\mathbf{x} \, d\mathbf{y} \right)^{1/2} \\ &= \left(\int_{\Omega^2} |u(\mathbf{y}, t) - u(\mathbf{x}, t)|^{2(p-1)} \, d\mathbf{x} \, d\mathbf{y} \right)^{1/2} \left(\int_{\Omega} \left(\int_{\Omega} |K_n(\mathbf{x}, \mathbf{y}) - K(\mathbf{x}, \mathbf{y})|^2 \, d\mathbf{y} \right) |\xi_n(\mathbf{x}, t)|^2 \, d\mathbf{x} \right)^{1/2} \\ &= \left(\int_{\Omega^2} |u(\mathbf{y}, t) - u(\mathbf{x}, t)|^{2(p-1)} \, d\mathbf{x} \, d\mathbf{y} \right)^{1/2} \|K_n - K\|_{L^{\infty,2}(\Omega^2)} \|\xi_n(\cdot, t)\|_{L^2(\Omega)}. \end{aligned}$$

On the one hand, under (a), Jensen’s inequality applied to the concave function $x \in \mathbb{R}^+ \mapsto x^{p-1}$ entails

$$\begin{aligned} \left(\int_{\Omega^2} |u(\mathbf{y}, t) - u(\mathbf{x}, t)|^{2(p-1)} \, d\mathbf{x} \, d\mathbf{y} \right)^{1/2} &\leq \left(\int_{\Omega^2} |u(\mathbf{y}, t) - u(\mathbf{x}, t)|^2 \, d\mathbf{x} \, d\mathbf{y} \right)^{(p-1)/2} \\ &\leq 2^{p-1} \|u(\cdot, t)\|_{L^2(\Omega)}^{p-1} \leq 2^{p-1} \left(\|g\|_{L^2(\Omega)} + \|f\|_{L^1([0,T];L^2(\Omega))} \right)^{p-1}, \end{aligned}$$

where we used (19) in the last inequality. On the other hand, under (b), we have

$$\begin{aligned} \left(\int_{\Omega^2} |u(\mathbf{y}, t) - u(\mathbf{x}, t)|^{2(p-1)} \, d\mathbf{x} \, d\mathbf{y} \right)^{1/2} &\leq 2^{p-1} \|u(\cdot, t)\|_{L^{2(p-1)}(\Omega)}^{p-1} \\ &\leq 2^{p-1} \left(\|g\|_{L^{2(p-1)}(\Omega)} + \|f\|_{L^1([0,T];L^{2(p-1)}(\Omega))} \right)^{p-1}. \end{aligned}$$

In turn, under either (a) or (b), we have the bound

$$\left| \int_{\Omega^2} (K_n(\mathbf{x}, \mathbf{y}) - K(\mathbf{x}, \mathbf{y})) \Psi(u(\mathbf{y}, t) - u(\mathbf{x}, t)) \xi_n(\mathbf{x}, t) \, d\mathbf{x} \, d\mathbf{y} \right| \leq C \|K_n - K\|_{L^\infty,2(\Omega^2)} \|\xi_n(\cdot, t)\|_{L^2(\Omega)}. \tag{27}$$

Inserting (26) and (27) into (25), ignoring the first term which is non-positive as argued above, and using Cauchy–Schwartz inequality on the last term, we obtain

$$\frac{\partial}{\partial t} \|\xi_n(\cdot, t)\|_{L^2(\Omega)} \leq \|f_n(\cdot, t) - f(\cdot, t)\|_{L^2(\Omega)} + \begin{cases} C \|K_n - K\|_{L^\infty,2(\Omega^2)}, & \text{under (a) or (b)} \\ C \|K_n - K\|_{L^2(\Omega^2)}, & \text{under (c)}. \end{cases}$$

Integrating this inequality on $[0, t]$ and taking the supremum over $t \in [0, T]$, we get (23).

- (iii) By assumptions on $\{K_n\}_{n \in \mathbb{N}}$, we are in position to apply the Vitali convergence theorem ([42], p. 133) in $L^2(\Omega^2)$ to get that $\|K_n - K\|_{L^2(\Omega^2)} \rightarrow 0$ as $n \rightarrow +\infty$. We have by assumption that the sequence $\{g_n\}_{n \in \mathbb{N}}$ is dominated by a constant function. The latter is obviously in $L^2(\Omega)$ since $|\Omega| < +\infty$. It then follows from the dominated convergence theorem that $\|g_n - g\|_{L^2(\Omega)} \rightarrow 0$ as $n \rightarrow +\infty$. We now turn to the sequence f_n . We have

$$\|f_n - f\|_{L^1([0,T];L^2(\Omega))} \leq T^{1/2} \|f_n - f\|_{L^2([0,T];L^2(\Omega))} = T^{1/2} \|f_n - f\|_{L^2(\Omega \times [0,T])}.$$

Arguing as for g_n , using our assumptions, entails again that $\|f_n - f\|_{L^1([0,T];L^2(\Omega))} \rightarrow 0$ as $n \rightarrow +\infty$. Passing to the limit in the second inequality of (23), the claim follows. □

In the case where the kernel takes the form $K(\mathbf{x}, \mathbf{y}) = J(\mathbf{x} - \mathbf{y})$, we have the following consequence of Theorem 5.1.

Corollary 5.1. *Suppose that $p \in]1, +\infty[$. Let u be a solution of (P) with kernel $K(\mathbf{x}, \mathbf{y}) = J(\mathbf{x} - \mathbf{y})$ and data (f, g) . Let u_n be a sequence of solutions to (P) with kernels $K_n(\mathbf{x}, \mathbf{y}) = J_n(\mathbf{x} - \mathbf{y})$ and data (f_n, g_n) . Assume that J and J_n satisfy (H’1), (H’2) and $J, J_n \in L^2(\Omega - \Omega)$, and that either one of (a), (b) or (c) in Theorem 5.1 holds. Then, the following hold.*

- (i) u and u_n are the unique solutions of the corresponding evolution problems.
- (ii) We have the error estimate

$$\|u_n - u\|_{C([0,T];L^2(\Omega))} \leq \|g_n - g\|_{L^2(\Omega)} + \|f_n - f\|_{L^1([0,T];L^2(\Omega))} + CT \|J_n - J\|_{L^2(\Omega - \Omega)}, \tag{28}$$

where C is positive constant that may depend only on p, g and f .

- (iii) Moreover, if the sequence $\{|J_n|^2\}_{n \in \mathbb{N}}$ is uniformly integrable over $\Omega - \Omega$, $J_n \rightarrow J$ pointwise a.e. on $\Omega - \Omega$, $g_n \rightarrow g$ pointwise a.e. on Ω , $f_n \rightarrow f$ pointwise a.e. on $\Omega \times [0, T]$, and either one of the following holds:
 - (a') $p \in]1, 2[$, $\{|g_n|^2\}_{n \in \mathbb{N}}$ (resp. $\{|f_n|^2\}_{n \in \mathbb{N}}$) is uniformly integrable over Ω (resp. $\Omega \times [0, T]$);
 - (b') $p \geq 2$, $\{|g_n|^{2(p-1)}\}_{n \in \mathbb{N}}$ (resp. $\{|f_n|^{2(p-1)}\}_{n \in \mathbb{N}}$) is uniformly integrable over Ω (resp. $\Omega \times [0, T]$);
 - (c') $\sup_{n \in \mathbb{N}} |g_n(\mathbf{x})| < +\infty$ a.e. on Ω and $\sup_{n \in \mathbb{N}} |f_n(\mathbf{x}, t)| < +\infty$ a.e. on $\Omega \times [0, T]$.

Then

$$\lim_{n \rightarrow +\infty} \|u_n - u\|_{C([0, T]; L^2(\Omega))} = 0.$$

Proof. (i) We argue in the same way as in the proof of Theorem 5.1 since $L^2(\Omega - \Omega) \subset L^1(\Omega - \Omega)$ implies that assumption (H' 3) holds for both J and J_n .

(ii) The error bound (28) is a specialization of (23) since

$$\int_{\Omega} |K_n(\mathbf{x}, \mathbf{y}) - K(\mathbf{x}, \mathbf{y})|^2 d\mathbf{y} = \int_{\Omega - \mathbf{x}} |J_n(\mathbf{z}) - J(\mathbf{z})|^2 d\mathbf{z} \leq \|J_n - J\|_{L^2(\Omega - \Omega)}^2.$$

Thus

$$\|K_n - K\|_{L^2(\Omega^2)} \leq \|K_n - K\|_{L^{\infty, 2}(\Omega^2)} \leq \|J_n - J\|_{L^2(\Omega - \Omega)}.$$

- (iii) Case (a') follows from the Vitali convergence theorem applied to J_n , g_n and f_n . The latter argument also applies to case (b') since $L^{2(p-1)}(\Omega - \Omega) \subset L^2(\Omega - \Omega)$, $L^{2(p-1)}(\Omega) \subset L^2(\Omega)$ and $L^{2(p-1)}(\Omega \times [0, T]) \subset L^1([0, T]; L^2(\Omega))$. Case (c') uses the Vitali convergence theorem on J_n and the dominated convergence theorem on g_n and f_n as argued in the proof of Theorem 5.1(iii). □

Remark 5.2. At this stage, we only relied on the monotonicity property of Δ_p^K in Proposition 4.1(iv) to get our bounds. One may then wonder if the stronger notions of monotonicity established in Proposition 4.1(v) can yield bounds better than (28). We answer this question positively by (slightly) improving the dependence on T for $p \in]1, 2]$ but at the price of more stringent assumptions on J . For this, we embark from (25), bound all terms as in the proof of Theorem 5.1, use Proposition 4.1(v) and that $L^2(\Omega) \subset L^p(\Omega)$ in this case to get

$$\begin{aligned} & \frac{1}{2} \frac{\partial}{\partial t} \|\xi_n(\cdot, t)\|_{L^2(\Omega)}^2 + C_1 \int_{\Omega^2} J(\mathbf{x} - \mathbf{y}) |\nabla^{\text{NL}} \xi_n(\mathbf{x}, \mathbf{y})|^2 d\mathbf{y} d\mathbf{x} \\ & \leq \left(C \|J_n - J\|_{L^2(\Omega - \Omega)} + \|f_n(\cdot, t) - f(\cdot, t)\|_{L^2(\Omega)} \right) \|\xi_n(\cdot, t)\|_{L^2(\Omega)}, \end{aligned}$$

for two positive constants C, C_1 (in the following C_i is a positive constant). Assume in addition that J is compactly supported and $J(0) > 0$. One can then invoke the Poincaré inequality ([3], Prop. 4.1) to show that

$$C_2 \left\| \xi_n(\cdot, t) - \int_{\Omega} \xi_n(\mathbf{x}, t) d\mathbf{x} \right\|_{L^2(\Omega)}^2 \leq \int_{\Omega^2} J(\mathbf{x} - \mathbf{y}) |\nabla^{\text{NL}} \xi_n(\mathbf{x}, \mathbf{y})|^2 d\mathbf{y} d\mathbf{x}.$$

Thus

$$\frac{1}{2} \|\xi_n(\cdot, t)\|_{L^2(\Omega)}^2 \leq \left\| \xi_n(\cdot, t) - \int_{\Omega} \xi_n(\mathbf{x}, t) d\mathbf{x} \right\|_{L^2(\Omega)}^2 + \left(\int_{\Omega} \xi_n(\mathbf{x}, t) d\mathbf{x} \right)^2.$$

Altogether, we arrive at

$$\begin{aligned} & \frac{1}{2} \frac{\partial}{\partial t} \|\xi_n(\cdot, t)\|_{L^2(\Omega)}^2 + \frac{C_1 C_2}{2} \|\xi_n(\cdot, t)\|_{L^2(\Omega)}^2 \\ & \leq \left(C \|J_n - J\|_{L^2(\Omega - \Omega)} + \|f_n(\cdot, t) - f(\cdot, t)\|_{L^2(\Omega)} \right) \|\xi_n(\cdot, t)\|_{L^2(\Omega)} + C_1 C_2 \left(\int_{\Omega} \xi_n(\mathbf{x}, t) d\mathbf{x} \right)^2. \end{aligned}$$

By integrating (\mathcal{P}) , it is easy to see by applying Proposition 4.1(v) and (iv) with $h(x) = 1$ that the solution of (\mathcal{P}) preserves the total mass in Ω , whence we deduce

$$\int_{\Omega} \xi_n(\mathbf{x}, t) \, d\mathbf{x} = \int_{\Omega} (g_n(\mathbf{x}) - g(\mathbf{x})) + \int_0^t \int_{\Omega} (f_n(\mathbf{x}, s) - f(\mathbf{x}, s)) \, d\mathbf{x} \, ds.$$

If (f, g) and (f_n, g_n) have the same mass, we get

$$\begin{aligned} \frac{1}{2} \frac{\partial}{\partial t} \|\xi_n(\cdot, t)\|_{L^2(\Omega)}^2 + \frac{C_1 C_2}{2} \|\xi_n(\cdot, t)\|_{L^2(\Omega)}^2 \\ \leq \left(C \|J_n - J\|_{L^2(\Omega-\Omega)} + \|f_n(\cdot, t) - f(\cdot, t)\|_{L^2(\Omega)} \right) \|\xi_n(\cdot, t)\|_{L^2(\Omega)}, \end{aligned}$$

and therefore

$$\frac{\partial}{\partial t} \|\xi_n(\cdot, t)\|_{L^2(\Omega)} + \frac{C_1 C_2}{2} \|\xi_n(\cdot, t)\|_{L^2(\Omega)} \leq \left(C \|J_n - J\|_{L^2(\Omega-\Omega)} + \|f_n(\cdot, t) - f(\cdot, t)\|_{L^2(\Omega)} \right).$$

Applying Gronwall's lemma yields the estimate

$$\begin{aligned} \|u_n(\mathbf{x}, t) - u(\mathbf{x}, t)\|_{L^2(\Omega)} &\leq \|f_n - f\|_{L^1([0, T]; L^2(\Omega))} + \exp(-C_1 C_2 t/2) \|g_n - g\|_{L^2(\Omega)} \\ &\quad + \frac{2C}{C_1 C_2} (1 - \exp(-C_1 C_2 t/2)) \|J_n - J\|_{L^2(\Omega-\Omega)}. \end{aligned}$$

This bound is clearly better than (28). In turn,

$$\|u_n - u\|_{C([0, T]; L^2(\Omega))} \leq \|f_n - f\|_{L^1([0, T]; L^2(\Omega))} + \max \left(\|g_n - g\|_{L^2(\Omega)}, \frac{2C}{C_1 C_2} \|J_n - J\|_{L^2(\Omega-\Omega)} \right).$$

The same reasoning as above can be carried out to sharpen the error bounds for the discrete problems in Section 6. Nevertheless, this will not be detailed further in this work.

5.2. The case $p = 1$

We now turn to the case $p = 1$.

Theorem 5.2. *Let u be a solution of (\mathcal{P}) for $p = 1$ with kernel K and data (f, g) . Let u_n be a sequence of solutions to (\mathcal{P}) for $p = 1$ with kernels K_n and data (f_n, g_n) . Assume that K and K_n satisfy (H.1) and (H.2), that $K, K_n \in L^{\infty, 2}(\Omega^2)$, $g, g_n \in L^2(\Omega)$ and $f, f_n \in L^1([0, T]; L^2(\Omega))$. Then, the following hold.*

- (i) u and u_n are the unique solutions in the sense of Definition 4.2 of the corresponding evolution problems.
- (ii) We have the error estimate

$$\|u_n - u\|_{C([0, T]; L^2(\Omega))} \leq \|g_n - g\|_{L^2(\Omega)} + \|f_n - f\|_{L^1([0, T]; L^2(\Omega))} + T \|K_n - K\|_{L^2(\Omega^2)}. \tag{29}$$

- (iii) Moreover, if $K_n \rightarrow K$ pointwise a.e. on Ω^2 , $g_n \rightarrow g$ pointwise a.e. on Ω , $f_n \rightarrow f$ pointwise a.e. on $\Omega \times [0, T]$, and $\{|K_n|^2\}_{n \in \mathbb{N}}$ is uniformly integrable over Ω^2 , $\{|g_n|^2\}_{n \in \mathbb{N}}$ is uniformly integrable on Ω , and $\{|f_n|^2\}_{n \in \mathbb{N}}$ is uniformly integrable on $\Omega \times [0, T]$. Then

$$\lim_{n \rightarrow +\infty} \|u_n - u\|_{C([0, T]; L^2(\Omega))} = 0.$$

Proof. (i) Existence and uniqueness of u and u_n follow from Theorem 4.2 where we argue as in Theorem 5.1(i) since $g, g_n \in L^2(\Omega) \subset L^1(\Omega)$ and $K, K_n \in L^{\infty, 2}(\Omega^2) \subset L^{\infty, 1}(\Omega^2)$.

(ii) Denote the error function $\xi_n(\mathbf{x}, t) = u_n(\mathbf{x}, t) - u(\mathbf{x}, t)$, then from Definition 4.2, we have a.e.

$$\begin{aligned} \frac{\partial \xi_n(\mathbf{x}, t)}{\partial t} &= \int_{\Omega} (K_n(\mathbf{x}, \mathbf{y})w_n(\mathbf{x}, \mathbf{y}, t) \, d\mathbf{y} - K(\mathbf{x}, \mathbf{y})w(\mathbf{x}, \mathbf{y}, t)) \, d\mathbf{y} + f_n(\mathbf{x}, t) - f(\mathbf{x}, t) \\ &= \int_{\Omega} K_n(\mathbf{x}, \mathbf{y})(w_n(\mathbf{x}, \mathbf{y}, t) - w(\mathbf{x}, \mathbf{y}, t)) \, d\mathbf{y} + \int_{\Omega} (K_n(\mathbf{x}, \mathbf{y}) - K(\mathbf{x}, \mathbf{y}))w(\mathbf{x}, \mathbf{y}, t) \, d\mathbf{y} \\ &\quad + f_n(\mathbf{x}, t) - f(\mathbf{x}, t), \end{aligned} \tag{30}$$

where w (resp. w_n) is the subgradient function associated to u (resp. u_n) as in Definition 4.2. Multiplying both sides of (30) by $\xi_n(\mathbf{x}, t)$ and integrating, we get

$$\begin{aligned} \frac{1}{2} \frac{\partial}{\partial t} \|\xi_n(\cdot, t)\|_{L^2(\Omega)}^2 &= \int_{\Omega^2} K_n(\mathbf{x}, \mathbf{y})(w_n(\mathbf{x}, \mathbf{y}, t) - w(\mathbf{x}, \mathbf{y}, t))\xi_n(\mathbf{x}, t) \, d\mathbf{x} \, d\mathbf{y} \\ &\quad + \int_{\Omega^2} (K_n(\mathbf{x}, \mathbf{y}) - K(\mathbf{x}, \mathbf{y}))w(\mathbf{x}, \mathbf{y}, t)\xi_n(\mathbf{x}, t) \, d\mathbf{x} \, d\mathbf{y} \\ &\quad + \int_{\Omega} (f_n(\mathbf{x}, t) - f(\mathbf{x}, t))\xi_n(\mathbf{x}, t) \, d\mathbf{x}. \end{aligned} \tag{31}$$

In view of the monotonicity claim in Proposition 4.2(ii), we have

$$\int_{\Omega^2} K_n(\mathbf{x}, \mathbf{y})(w_n(\mathbf{x}, \mathbf{y}, t) - w(\mathbf{x}, \mathbf{y}, t))\xi_n(\mathbf{x}, t) \, d\mathbf{x} \, d\mathbf{y} \leq 0.$$

Let us turn to bounding the second term. We have by the Cauchy–Schwartz inequality and that $\|w\|_{L^\infty(\Omega^2 \times]0, T])} \leq 1$,

$$\begin{aligned} \left| \int_{\Omega^2} (K_n(\mathbf{x}, \mathbf{y}) - K(\mathbf{x}, \mathbf{y}))w(\mathbf{x}, \mathbf{y}, t)\xi_n(\mathbf{x}, t) \, d\mathbf{x} \, d\mathbf{y} \right| &\leq \int_{\Omega^2} |K_n(\mathbf{x}, \mathbf{y}) - K(\mathbf{x}, \mathbf{y})|\xi_n(\mathbf{x}, t) \, d\mathbf{x} \, d\mathbf{y} \\ &\leq \|K_n - K\|_{L^2(\Omega^2)} \|\xi_n(\cdot, t)\|_{L^2(\Omega)}. \end{aligned} \tag{32}$$

Inserting (32) into (31), ignoring the first term which is non-positive as argued above, and using Cauchy–Schwartz inequality on the last term, we obtain

$$\frac{\partial}{\partial t} \|\xi_n(\cdot, t)\|_{L^2(\Omega)} \leq \|f_n(\cdot, t) - f(\cdot, t)\|_{L^2(\Omega)} + \|K_n - K\|_{L^2(\Omega^2)}.$$

Integrating this inequality on $[0, t]$ and taking the supremum over $t \in [0, T]$, we get (29).

(iii) We argue again using the Vitali convergence theorem since $K, K_n \in L^\infty \cdot 2(\Omega^2) \subset L^2(\Omega^2)$ and $L^1([0, T]; L^2(\Omega)) \subset L^2(\Omega \times [0, T])$. □

The following corollary is immediate in the same vein as Corollary 5.1.

Corollary 5.2. *Let u be a solution of (\mathcal{P}) for $p = 1$ with kernel $K(\mathbf{x}, \mathbf{y}) = J(\mathbf{x} - \mathbf{y})$ and data (f, g) . Let u_n be a sequence of solutions to (\mathcal{P}) for $p = 1$ with kernels $K_n(\mathbf{x}, \mathbf{y}) = J_n(\mathbf{x} - \mathbf{y})$ and data (f_n, g_n) . Assume that J and J_n satisfy $(H'.1)$, $(H'.2)$ and $J, J_n \in L^2(\Omega - \Omega)$, that $g, g_n \in L^2(\Omega)$ and $f, f_n \in L^1([0, T]; L^2(\Omega))$. Then, the following hold.*

- (i) u and u_n are the unique solutions in the sense of Definition 4.2 of the corresponding evolution problems.
- (ii) We have the error estimate

$$\|u_n - u\|_{C([0, T]; L^2(\Omega))} \leq \|g_n - g\|_{L^2(\Omega)} + \|f_n - f\|_{L^1([0, T]; L^2(\Omega))} + T\|J_n - J\|_{L^2(\Omega - \Omega)}. \tag{33}$$

(iii) Moreover, if $J_n \rightarrow J$ pointwise a.e. on $\Omega - \Omega$, $g_n \rightarrow g$ pointwise a.e. on Ω , $f_n \rightarrow f$ pointwise a.e. on $\Omega \times [0, T]$, and $\{|J_n|^2\}_{n \in \mathbb{N}}$ is uniformly integrable over $\Omega - \Omega$, $\{|g_n|^2\}_{n \in \mathbb{N}}$ is uniformly integrable on Ω , and $\{|f_n|^2\}_{n \in \mathbb{N}}$ is uniformly integrable on $\Omega \times [0, T]$. Then

$$\lim_{n \rightarrow +\infty} \|u_n - u\|_{C([0, T]; L^2(\Omega))} = 0.$$

6. ERROR BOUNDS FOR THE DISCRETE PROBLEM

Let $\mathbf{K} \in \mathbb{R}^{n^d \times n^d}$ and $\mathbf{g} \in \mathbb{R}^{n^d}$ be discrete approximations of, respectively, the kernel K and initial data g in (\mathcal{P}) , on a regular mesh of size δ_n . Typically, one can take $\mathbf{K} = P_n K$ and $\mathbf{g} = P_n g$. For $1 < p < \infty$, the discrete p -Laplacian operator with kernel \mathbf{K} is

$$\widehat{\Delta}_p^{\mathbf{K}} : \mathbf{u} \in \mathbb{R}^{n^d} \mapsto - \sum_{j \in [n]^d} h_j \mathbf{K}_{ij} |\mathbf{u}_j - \mathbf{u}_i|^{p-2} (\mathbf{u}_j - \mathbf{u}_i) = - \sum_{j \in [n]^d} h_j \mathbf{K}_{ij} \Psi(\mathbf{u}_j - \mathbf{u}_i).$$

In the same way, we define the discrete 1-Laplacian operator as the set-valued operator $\widehat{\Delta}_1^{\mathbf{K}} : \mathbb{R}^{n^d} \rightarrow 2^{\mathbb{R}^{n^d}}$ such that $\boldsymbol{\eta} \in \widehat{\Delta}_1^{\mathbf{K}} \mathbf{u}$ if and only if

$$\boldsymbol{\eta}_i = - \sum_{j \in [n]^d} h_j \mathbf{K}_{ij} \mathbf{w}_{ij},$$

where $\|\mathbf{w}\|_\infty \leq 1$, $\mathbf{w}_{ij} = -\mathbf{w}_{ji}$, and

$$\mathbf{w}_{ij} \in \partial |\cdot| (\mathbf{u}_j - \mathbf{u}_i).$$

By construction, we have the following simple lemma whose proof is immediate.

Lemma 6.1. For any $\mathbf{K} \in \mathbb{R}^{n^d \times n^d}$ and $\mathbf{u} \in \mathbb{R}^{n^d}$, the following holds:

(i) If $1 < p < +\infty$,

$$I_n \widehat{\Delta}_p^{\mathbf{K}}(\mathbf{u}) = \Delta_p^{I_n \mathbf{K}}(I_n \mathbf{u}).$$

(ii) If $p = 1$,

$$I_n \boldsymbol{\eta}(\mathbf{x}) = - \int_{\Omega} I_n K(\mathbf{x}, \mathbf{y}) I_n \mathbf{w}(\mathbf{x}, \mathbf{y}) \, d\mathbf{y}, \quad \text{where } I_n \mathbf{w}(\mathbf{x}, \mathbf{y}) \in \partial |\cdot| (I_n \mathbf{u}(\mathbf{y}) - I_n \mathbf{u}(\mathbf{x})).$$

Moreover, $\|I_n \mathbf{w}\|_{L^\infty(\Omega^2)} \leq 1$ and $I_n \mathbf{w}(\mathbf{x}, \mathbf{y}) = -I_n \mathbf{w}(\mathbf{y}, \mathbf{x})$.

6.1. The semi-discrete problem

Case $p \in]1, +\infty[$: We start with the case $1 < p < +\infty$ and consider the space semi-discretization of (\mathcal{P}) ,

$$\begin{cases} \frac{\partial}{\partial t} \mathbf{u}(t) = -\widehat{\Delta}_p^{\mathbf{K}} \mathbf{u}(t) + \mathbf{f}(t), & t > 0, \\ \mathbf{u}(0) = \mathbf{g}. \end{cases} \tag{\mathcal{P}_p^{\text{SD}}}$$

where $\mathbf{u} : t \in \mathbb{R}^+ \mapsto \mathbf{u}(t) \in \mathbb{R}^{n^d}$ and similarly for \mathbf{f} .

Our aim is to compare the solutions of problems (\mathcal{P}) and $(\mathcal{P}_p^{\text{SD}})$. The solution of $(\mathcal{P}_p^{\text{SD}})$ being discrete in space, we consider its continuum space extensions of \mathbf{u} and \mathbf{f} on Ω for any $t > 0$ as

$$u_n(\mathbf{x}, t) = (I_n \mathbf{u}(t))(\mathbf{x}) \quad \text{and} \quad f_n(\mathbf{x}, t) = (I_n \mathbf{f}(t))(\mathbf{x}). \tag{34}$$

Theorem 6.1. *Suppose that $p \in]1, +\infty[$. Let u be a solution of (\mathcal{P}) with kernel K and data (f, g) , and \mathbf{u} that of $(\mathcal{P}_p^{\text{SD}})$ with $\mathbf{K} = \mathbf{P}_n K$, $\mathbf{g} = \mathbf{P}_n g$ and $\mathbf{f}(t) = \mathbf{P}_n f(\cdot, t)$ for $t \in [0, T]$. Let u_n and f_n as defined in (34). Assume that K satisfies (H.1), (H.2) and $K \in L^{\infty,2}(\Omega^2)$, and that g and f satisfy either one of the conditions (a), (b) or (c) in Theorem 5.1. Then, the following hold.*

- (i) u and u_n are the unique solutions of (\mathcal{P}) with data respectively (f, g) and $(f_n, \mathbf{I}_n \mathbf{P}_n g)$.
- (ii) We have the error estimate

$$\begin{aligned} \|u_n - u\|_{C([0,T];L^2(\Omega))} &\leq \|\mathbf{I}_n \mathbf{P}_n g - g\|_{L^2(\Omega)} + \|f_n - f\|_{L^1([0,T];L^2(\Omega))} \\ &\quad + CT \begin{cases} \|\mathbf{I}_n \mathbf{P}_n K - K\|_{L^{\infty,2}(\Omega^2)}, & \text{under (a), (b)} \\ \|\mathbf{I}_n \mathbf{P}_n K - K\|_{L^2(\Omega^2)}, & \text{under (c)} \end{cases} \end{aligned} \quad (35)$$

where C is positive constant that depends only on p, g and f .

- (iii) If, moreover, $g \in L^\infty(\Omega) \cap \text{Lip}(s, L^2(\Omega))$, $K \in \text{Lip}(s, L^2(\Omega^2))$ and $f(\cdot, t) \in L^\infty(\Omega) \cap \text{Lip}(s, L^2(\Omega))$ for every $t \in [0, T]$, then

$$\|u_n - u\|_{C([0,T];L^2(\Omega))} \leq C(1+T)\delta_n^s, \quad (36)$$

where C is positive constant that depends only on p, g, f, K, s .

Proof. (i) Existence and uniqueness of u were proved in Theorem 5.1(i). We also see that $\mathbf{I}_n \mathbf{K}$ verifies (H.1) and (H.2). Using Lemma 2.1, we have $\mathbf{I}_n \mathbf{g} \in L^p(\Omega)$, $f_n \in L^1([0, T], L^p(\Omega))$ and $\mathbf{I}_n \mathbf{K} \in L^{\infty,2}(\Omega^2) \subset L^{\infty,1}(\Omega^2)$, and thus $\mathbf{I}_n \mathbf{K}$ fulfills (H.3). In view of Lemma 6.1(i), it follows from $(\mathcal{P}_p^{\text{SD}})$ that the function u_n satisfies (\mathcal{P}) with kernel $\mathbf{I}_n \mathbf{K}$ and data $(f_n, \mathbf{I}_n \mathbf{g})$. Existence and uniqueness of u_n then follow from Theorem 4.1.

- (ii) The claim is a specialization of (23) in Theorem 5.1(ii).
- (iii) As $K \in L^{\infty,2}(\Omega^2) \subset L^2(\Omega^2)$, we insert the estimate (6) (see Lem. 2.2) in the second bound of (35). □

Case $p = 1$: we now turn to the case $p = 1$, and consider the evolution problem

$$\begin{cases} \frac{\partial}{\partial t} \mathbf{u}(t) = -\boldsymbol{\eta}(t) + \mathbf{f}(t), & t > 0, \\ \mathbf{u}(0) = \mathbf{g}, \end{cases} \quad (\mathcal{P}_1^{\text{SD}})$$

where

$$\boldsymbol{\eta}_i(t) = - \sum_{j \in [n]^d} h_j \mathbf{K}_{ij} \text{sign}(\mathbf{u}_j - \mathbf{u}_i), \quad \text{and thus} \quad \boldsymbol{\eta}(t) \in \widehat{\Delta}_1^{\mathbf{K}} \mathbf{u}(t).$$

Theorem 6.2. *Let u be a solution of (\mathcal{P}) for $p = 1$ with kernel K and data (f, g) , and \mathbf{u} is that of $(\mathcal{P}_1^{\text{SD}})$ with $\mathbf{K} = \mathbf{P}_n K$, $\mathbf{g} = \mathbf{P}_n g$ and $\mathbf{f}(t) = \mathbf{P}_n f(\cdot, t)$ for $t \in [0, T]$. Let u_n and f_n as defined in (34). Assume that K satisfies (H.1), (H.2) and $K \in L^{\infty,2}(\Omega^2)$, and that $g \in L^2(\Omega)$ and $f \in L^1([0, T]; L^2(\Omega))$. Then, the following hold.*

- (i) u and u_n are the unique solutions in the sense of Definition 4.2 of the corresponding evolution problems.
- (ii) We have the error estimate

$$\|u_n - u\|_{C([0,T];L^2(\Omega))} \leq \|\mathbf{I}_n \mathbf{P}_n g - g\|_{L^2(\Omega)} + \|f_n - f\|_{L^1([0,T];L^2(\Omega))} + T \|\mathbf{I}_n \mathbf{P}_n K - K\|_{L^2(\Omega^2)}. \quad (37)$$

- (iii) If, moreover, $g \in \text{Lip}(s, L^2(\Omega))$, $K \in \text{Lip}(s, L^2(\Omega^2))$ and $f(\cdot, t) \in \text{Lip}(s, L^2(\Omega))$ for every $t \in [0, T]$, then

$$\|u_n - u\|_{C([0,T];L^2(\Omega))} \leq C(1+T)\delta_n^s, \quad (38)$$

where C is positive constant that depends only on p, g, f, K and s .

Proof. (i) Existence and uniqueness of u were proved in Theorem 5.2(i). In addition, $I_n \mathbf{K}$ verifies (H.1) and (H.2). Using Lemma 2.1, $I_n \mathbf{g} \in L^2(\Omega) \subset L^1(\Omega)$, $f_n \in L^1([0, T]; L^2(\Omega)) \subset L^1([0, T]; L^1(\Omega))$ and $I_n \mathbf{K} \in L^{\infty, 2}(\Omega^2) \subset L^{\infty, 1}(\Omega^2)$, and thus $I_n \mathbf{K}$ fulfills (H.3). By virtue of Lemma 6.1(ii), u_n , the space continuum extension of \mathbf{u} , will satisfy (P) with kernel $I_n \mathbf{K}$ and data $(f_n, I_n \mathbf{g})$. Existence and uniqueness of u_n in the sense of Definition 4.2 follow from Theorem 4.2.

(ii) This claim is a specialization of (29) in Theorem 5.2(ii).

(iii) Insert the estimate (6) in (37). □

6.2. The totally discrete problem

We establish in this section error bounds for fully discrete (in time and space) approximations of (P). For that, let $0 < t_1 < t_2 < \dots < t_{N-1} < t_N = T$ be a partition (not necessarily equispaced) of $[0, T]$. Let $\tau_{k-1} \stackrel{\text{def}}{=} |t_k - t_{k-1}|$ and denote $\tau = \max_{k \in [N]} \tau_k$.

6.2.1. Forward/Explicit Euler discretization

Case $p \in]1, 2]$: we start with $p \in]1, 2]$ and consider a totally discrete problem with forward/explicit Euler scheme in time,

$$\begin{cases} \frac{\mathbf{u}^k - \mathbf{u}^{k-1}}{\tau_{k-1}} = -\widehat{\Delta}_p^{\mathbf{K}} \mathbf{u}^{k-1} + \mathbf{f}, & k \in [N], \\ \mathbf{u}^0 = \mathbf{g}, \end{cases} \tag{P}_p^{\text{TDF}}$$

where $\mathbf{u}^k, \mathbf{f} \in \mathbb{R}^{n^d}$. We have implicitly assumed that \mathbf{f} does not depend on time, which is a standard assumption in the context of explicit discretization.

Since our aim is to compare the solutions of problems (P) and $(P)_p^{\text{TDF}}$, we introduce the following continuum extensions in space and/or time of $\{\mathbf{u}^k\}_{k \in [N]}$ as

$$\begin{aligned} u_n^k &= I_n \mathbf{u}^k, k \in [N], \text{ and } f_n = I_n \mathbf{f}, \\ \check{u}_n(\mathbf{x}, t) &= \frac{t_k - t}{\tau_{k-1}} u_n^{k-1}(\mathbf{x}) + \frac{t - t_{k-1}}{\tau_{k-1}} u_n^k(\mathbf{x}), & (\mathbf{x}, t) \in \Omega \times]t_{k-1}, t_k], k \in [N], \\ \bar{u}_n(\mathbf{x}, t) &= \sum_{k=1}^N u_n^{k-1}(\mathbf{x}) \chi_{]t_{k-1}, t_k]}(t), & (\mathbf{x}, t) \in \Omega \times]0, T]. \end{aligned}$$

Then, in the same vein as Lemma 6.1, it is easy to see that $(P)_p^{\text{TDF}}$ is equivalent to the following evolution problem

$$\begin{cases} \frac{\partial}{\partial t} \check{u}_n(\mathbf{x}, t) = -\Delta_p^{I_n \mathbf{K}} \bar{u}_n(\mathbf{x}, t) + f_n(\mathbf{x}), & (\mathbf{x}, t) \in \Omega \times]0, T], \\ \check{u}_n(\mathbf{x}, 0) = I_n \mathbf{g}(\mathbf{x}), & \mathbf{x} \in \Omega. \end{cases} \tag{39}$$

Before turning to the consistency result, we collect some useful estimates.

Lemma 6.2. Consider problem $(P)_p^{\text{TDF}}$ with kernel \mathbf{K} , data (\mathbf{f}, \mathbf{g}) and variable step-size $\tau_k \leq 2C \|\Delta_p^{I_n \mathbf{K}} u_n^k - f_n\|_{L^2(\Omega)}^{\frac{2-p}{p-1}}$, where C is the constant in Proposition 4.1(vi). Assume that $I_n \mathbf{g} \in L^2(\Omega)$ and $I_n \mathbf{K}$ satisfies (H.1)–(H.3). Suppose also that for each $n \in \mathbb{N}$, \mathbf{f} is such that $(P)_p^{\text{TDF}}$ has a stationary solution \mathbf{u}^* and that $\sup_{n \in \mathbb{N}} \|I_n \mathbf{g} - I_n \mathbf{u}^*\|_{L^2(\Omega)} < +\infty$. Then

$$\bar{u}_n(\cdot, t) \in L^2(\Omega), \forall t \in [0, T], \text{ and } \sup_{t \in [0, T], n \in \mathbb{N}} \|\bar{u}_n(\cdot, t) - I_n \mathbf{u}^*\|_{L^2(\Omega)} < +\infty.$$

Remark 6.1.

- (i) Condition on the time-step τ_k can be seen as an abstract non-linear CFL condition. It is better than the one in [30] since we here exploit the Hölder continuity of $\Delta_p^{I_n \mathbf{K}}$ on $L^2(\Omega)$ for $p \in]1, 2]$, see Proposition 4.1(vi). For $p = 2$, where $\Delta_2^{I_n \mathbf{K}}$ is linear Lipschitz continuous operator on $L^2(\Omega)$, the condition reads $\tau_k \leq 2C$. Such condition for explicit time-discretization of evolution problems with accretive and Lipschitz-continuous operators is known, see *e.g.*, [41]. It is also consistent with known convergence results for finding zeros of co-called co-coercive operators on Hilbert spaces [8].
- (ii) The assumption on \mathbf{f} and \mathbf{K} imply that $f_n \in L^2(\Omega)$. Indeed, (18) entails

$$\|f_n\|_{L^2(\Omega)} = \|\Delta_p^{I_n \mathbf{K}}(\mathbf{I}_n \mathbf{u}^*)\|_{L^2(\Omega)} \leq 2^p C_2 \|K\|_{L^\infty, 1(\Omega^2)}^2 \|\mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^{p-1}.$$

- (iii) The assumption made on \mathbf{f} is trivially true when $\mathbf{f} = \mathbf{0}$ since $\mathbf{0}$ is a stationary solution in this case. In turn, using Lemma 2.1, one can see that the uniform boundedness conditions on \mathbf{g} and \mathbf{K} are fulfilled if $\mathbf{g} = P_n g$ and $\mathbf{K} = P_n K$, where $g \in L^2(\Omega)$ and K satisfies (H.1)–(H.3).

Proof. We show the claim by an induction argument. Since $\Delta_p^{I_n \mathbf{K}}(\mathbf{I}_n \mathbf{u}^*) = f_n$, we have

$$\begin{aligned} \|u_n^1 - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 &= \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 - 2\tau_0 \int_{\Omega} (\Delta_p^{I_n \mathbf{K}}(\mathbf{I}_n \mathbf{g})(\mathbf{x}) - f_n(\mathbf{x})) (\mathbf{I}_n \mathbf{g}(\mathbf{x}) - \mathbf{I}_n \mathbf{u}^*) \, d\mathbf{x} \\ &\quad + \tau_0^2 \|\Delta_p^{I_n \mathbf{K}}(\mathbf{I}_n \mathbf{g}) - f_n\|_{L^2(\Omega)}^2 \\ &= \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 - 2\tau_0 \int_{\Omega} (\Delta_p^{I_n \mathbf{K}}(\mathbf{I}_n \mathbf{g})(\mathbf{x}) - \Delta_p^{I_n \mathbf{K}}(\mathbf{I}_n \mathbf{u}^*)(\mathbf{x})) (\mathbf{I}_n \mathbf{g}(\mathbf{x}) - \mathbf{I}_n \mathbf{u}^*) \, d\mathbf{x} \\ &\quad + \tau_0^2 \|\Delta_p^{I_n \mathbf{K}}(\mathbf{I}_n \mathbf{g}) - f_n\|_{L^2(\Omega)}^2. \end{aligned}$$

By assumption on \mathbf{g} , \mathbf{u}^* and τ_k , we can invoke Proposition 4.1(vi) to get

$$\begin{aligned} \|u_n^1 - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 &\leq \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 - 2C\tau_0 \|\Delta_p^{I_n \mathbf{K}}(\mathbf{I}_n \mathbf{g}) - f_n\|_{L^2(\Omega)}^{p/(p-1)} + \tau_0^2 \|\Delta_p^{I_n \mathbf{K}}(\mathbf{I}_n \mathbf{g}) - f_n\|_{L^2(\Omega)}^2 \\ &\leq \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 + \tau_0 \|\Delta_p^{I_n \mathbf{K}}(\mathbf{I}_n \mathbf{g}) - f_n\|_{L^2(\Omega)}^2 \left(2C \|\Delta_p^{I_n \mathbf{K}}(\mathbf{I}_n \mathbf{g}) - f_n\|_{L^2(\Omega)}^{(2-p)/(p-1)} - \tau_0 \right) \\ &\leq \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2. \end{aligned}$$

Suppose now that, for any $k > 1$,

$$\|u_n^k - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 \leq \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2,$$

and thus $u_n^k \in L^2(\Omega)$. We can then use Proposition 4.1(vi) as above to see that

$$\begin{aligned} \|u_n^{k+1} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 &\leq \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 - \tau_k \|\Delta_p^{I_n \mathbf{K}}(u_n^k) - f_n\|_{L^2(\Omega)}^2 \left(2C \|\Delta_p^{I_n \mathbf{K}}(u_n^k) - f_n\|_{L^2(\Omega)}^{(2-p)/(p-1)} - \tau_k \right) \\ &\leq \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2. \end{aligned}$$

Thus the sequence $\left\{ \|u_n^k\|_{L^2(\Omega)} \right\}_{k \in \mathbb{N}}$ is bounded, and so is $\|\bar{u}_n(\cdot, t)\|_{L^2(\Omega)}$ for $t \in [0, T]$ by its definition. We also have

$$\sup_{t \in [0, T], n \in \mathbb{N}} \|\bar{u}_n(\cdot, t) - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)} = \sup_{(n, N) \in \mathbb{N}^2, k \in [N]} \|u_n^k - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)} \leq \sup_{n \in \mathbb{N}} \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)} < +\infty.$$

□

Lemma 6.3. *In addition to the assumptions of Lemma 6.2, suppose that $\sup_{n \in \mathbb{N}} \|\mathbf{I}_n \mathbf{K}\|_{L^\infty,1(\Omega^2)} < +\infty$. Then*

$$\sup_{t \in [0,T], n \in \mathbb{N}} \|\check{u}_n(\cdot, t) - \bar{u}_n(\cdot, t)\|_{L^2(\Omega)} \leq C\tau,$$

where C is a positive constant that does not depend on (n, N, T) .

Proof. It is easy to see that for $t \in]t_{k-1}, t_k]$, $k \in \mathbb{N}$,

$$\begin{aligned} \|\check{u}_n(\cdot, t) - \bar{u}_n(\cdot, t)\|_{L^2(\Omega)} &= (t - t_{k-1}) \left\| \frac{u_n^k - u_n^{k-1}}{\tau_{k-1}} \right\|_{L^2(\Omega)} \\ &= (t - t_{k-1}) \|\Delta_p^{\mathbf{I}_n \mathbf{K}} u_n^{k-1} - f_n\|_{L^2(\Omega)} \\ &= (t - t_{k-1}) \|\Delta_p^{\mathbf{I}_n \mathbf{K}} u_n^{k-1} - \Delta_p^{\mathbf{I}_n \mathbf{K}} \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)} \\ &\leq \tau \|\Delta_p^{\mathbf{I}_n \mathbf{K}} u_n^{k-1} - \Delta_p^{\mathbf{I}_n \mathbf{K}} \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)} = \tau \|\Delta_p^{\mathbf{I}_n \mathbf{K}} \bar{u}_n(\cdot, t) - \Delta_p^{\mathbf{I}_n \mathbf{K}} \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}. \end{aligned}$$

As $\Delta_p^{\mathbf{I}_n \mathbf{K}}$ is Hölder continuous on $L^2(\Omega)$ with exponent $p - 1$, see (18), we get

$$\|\check{u}_n(\cdot, t) - \bar{u}_n(\cdot, t)\|_{L^2(\Omega)} \leq \tau 2^{p/2} C_2^{1/2} \|K\|_{L^\infty,1(\Omega^2)} \|\bar{u}_n(\cdot, t) - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^{p-1}.$$

We then take the supremum over t and n , and use Lemma 6.2 to conclude. □

We are now in position to state the error bound for the totally discrete problem $(\mathcal{P}_p^{\text{TDF}})$.

Theorem 6.3. *Suppose that $p \in]1, 2]$. Let u be a solution of (\mathcal{P}) with kernel K and data (f, g) where f is time-independent, and $\{\mathbf{u}^k\}_{k \in [N]}$ is the sequence generated by $(\mathcal{P}_p^{\text{TDF}})$ with $\mathbf{K} = \mathbf{P}_n K$, $\mathbf{g} = \mathbf{P}_n g$, $\mathbf{f} = \mathbf{P}_n f$ and τ_k as prescribed in Lemma 6.2. Assume that K satisfies (H.1), (H.2) and $K \in L^\infty,2(\Omega^2)$, and that f, g belong either to $L^2(\Omega)$ or $L^\infty(\Omega)$. Then, the following hold.*

- (i) u is the unique solution of (\mathcal{P}) , $\{\mathbf{u}^k\}_{k \in [N]}$ is uniquely defined and $\left\{ \|\mathbf{I}_n \mathbf{u}^k\|_{L^2(\Omega)} \right\}_{k \in [N]}$ is bounded (uniformly in n when $\mathbf{f} = \mathbf{0}$).
- (ii) We have the error estimate

$$\begin{aligned} \sup_{k \in [N], t \in]t_{k-1}, t_k]} \|\mathbf{I}_n \mathbf{u}^{k-1} - u(\cdot, t)\|_{L^2(\Omega)} &\leq \exp(T/2) \left(\|\mathbf{I}_n \mathbf{P}_n g - g\|_{L^2(\Omega)} \right. \\ &\quad \left. + CT^{1/2} \left(\tau^{1/(3-p)} + \|f_n - f\|_{L^2(\Omega)} + \begin{cases} \|\mathbf{I}_n \mathbf{P}_n K - K\|_{L^\infty,2(\Omega^2)} & g \in L^2(\Omega) \\ \|\mathbf{I}_n \mathbf{P}_n K - K\|_{L^2(\Omega^2)} & g \in L^\infty(\Omega) \end{cases} \right) \right) \end{aligned} \quad (40)$$

for τ sufficiently small, where C is positive constant that depends only on p, g, f and K .

- (iii) If, moreover, $f, g \in L^\infty(\Omega) \cap \text{Lip}(s, L^2(\Omega))$ and $K \in \text{Lip}(s, L^2(\Omega^2))$, then

$$\sup_{k \in [N], t \in]t_{k-1}, t_k]} \|\mathbf{I}_n \mathbf{u}^{k-1} - u(\cdot, t)\|_{L^2(\Omega)} \leq C \exp(T/2) \left((1 + T^{1/2}) \delta_n^s + T^{1/2} \tau^{1/(3-p)} \right), \quad (41)$$

for τ sufficiently small, where C is positive constant that depends only on p, g, f, K and s .

Proof. In the proof, C is any positive constant that may depend only on p, g, f, K and/or s , and that may be different at each line.

- (i) Existence and uniqueness of u were proved in Theorem 5.1(i). The claimed well-posedness of the sequence $\{\mathbf{u}^k\}_{k \in [N]}$ is a consequence of Lemma 6.2 and Remark 6.1(iii).

(ii) Denote $\check{\xi}_n(\mathbf{x}, t) = \check{u}_n(\mathbf{x}, t) - u(\mathbf{x}, t)$, $\bar{\xi}_n(\mathbf{x}, t) = \bar{u}_n(\mathbf{x}, t) - u(\mathbf{x}, t)$, $g_n = \mathbf{I}_n \mathbf{P}_n g$ and $K_n = \mathbf{I}_n \mathbf{P}_n K$. We thus have a.e.

$$\begin{aligned} \frac{\partial \check{\xi}_n(\mathbf{x}, t)}{\partial t} &= -(\Delta_p^{K_n}(\bar{u}_n(\mathbf{x}, t)) - \Delta_p^K(u(\mathbf{x}, t))) + (f_n(\mathbf{x}) - f(\mathbf{x})) \\ &= -(\Delta_p^{K_n}(\bar{u}_n(\mathbf{x}, t)) - \Delta_p^{K_n}(u(\mathbf{x}, t))) - (\Delta_p^{K_n}(u(\mathbf{x}, t)) - \Delta_p^K(u(\mathbf{x}, t))) + (f_n(\mathbf{x}) - f(\mathbf{x})). \end{aligned}$$

Multiplying both sides by $\check{\xi}_n(\mathbf{x}, t)$, integrating and rearranging the terms, we get

$$\begin{aligned} \frac{1}{2} \frac{\partial}{\partial t} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2 &= - \int_{\Omega} (\Delta_p^{K_n} \bar{u}_n(\mathbf{x}, t) - \Delta_p^{K_n} u(\mathbf{x}, t)) (\bar{u}_n(\mathbf{x}, t) - u(\mathbf{x}, t)) \, d\mathbf{x} \\ &\quad - \int_{\Omega} (\Delta_p^{K_n} u(\mathbf{x}, t) - \Delta_p^K u(\mathbf{x}, t)) \check{\xi}_n(\mathbf{x}, t) \, d\mathbf{x} \\ &\quad - \int_{\Omega} (\Delta_p^{K_n} \bar{u}_n(\mathbf{x}, t) - \Delta_p^{K_n} u(\mathbf{x}, t)) (\check{u}_n(\mathbf{x}, t) - \bar{u}_n(\mathbf{x}, t)) \, d\mathbf{x} \\ &\quad + \int_{\Omega} (f_n(\mathbf{x}) - f(\mathbf{x})) \check{\xi}_n(\mathbf{x}, t) \, d\mathbf{x}. \end{aligned} \tag{42}$$

Since $f, g \in L^p(\Omega)$ in both cases, so is $u(\cdot, t)$ thanks to (19). We also have $\bar{u}_n(\cdot, t) \in L^2(\Omega) \subset L^p(\Omega)$ by Lemma 6.2. We are then in position to use Proposition 4.1(iv) with $h(x) = x$ to assert that the first term on the right-hand side of (42) is nonpositive. Let us now bound the second term.

Similarly to the estimates (27) and (26) in the proof of Theorem 5.1, and using Young inequality, we have

$$\begin{aligned} \left| \int_{\Omega} (\Delta_p^{K_n} u(\mathbf{x}, t) - \Delta_p^K u(\mathbf{x}, t)) \check{\xi}_n(\mathbf{x}, t) \, d\mathbf{x} \right| &\leq \begin{cases} C \|K_n - K\|_{L^\infty, 2(\Omega^2)} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}, & g \in L^2(\Omega) \\ C \|K_n - K\|_{L^2(\Omega^2)} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}, & g \in L^\infty(\Omega), \end{cases} \\ &\leq \frac{1}{6} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2 + \begin{cases} C \|K_n - K\|_{L^\infty, 2(\Omega^2)}^2, & g \in L^2(\Omega) \\ C \|K_n - K\|_{L^2(\Omega^2)}^2, & g \in L^\infty(\Omega). \end{cases} \end{aligned}$$

For the third term in (42), we invoke Lemma 6.3 to get

$$\begin{aligned} &\left| \int_{\Omega} (\Delta_p^{K_n} \bar{u}_n(\mathbf{x}, t) - \Delta_p^{K_n} u(\mathbf{x}, t)) (\check{u}_n(\mathbf{x}, t) - \bar{u}_n(\mathbf{x}, t)) \, d\mathbf{x} \right| \\ &\leq \|\Delta_p^{K_n} \bar{u}_n(\cdot, t) - \Delta_p^{K_n} u(\cdot, t)\|_{L^2(\Omega)} \|\check{u}_n(\cdot, t) - \bar{u}_n(\cdot, t)\|_{L^2(\Omega)} \\ &\leq C \|\Delta_p^{K_n} \bar{u}_n(\cdot, t) - \Delta_p^{K_n} u(\cdot, t)\|_{L^2(\Omega)} \tau. \end{aligned}$$

We then use the fact that $\Delta_p^{\mathbf{I}_n \mathbf{K}}$ is Hölder continuous on $L^2(\Omega)$ with exponent $p - 1$, see (18), to obtain

$$\|\Delta_p^{K_n} \bar{u}_n(\cdot, t) - \Delta_p^{K_n} u(\cdot, t)\|_{L^2(\Omega)} \leq C \|\bar{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^{p-1} \leq C \left(\|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^{p-1} + \tau^{p-1} \right),$$

where we used Lemma 6.3 to go from $\bar{\xi}_n$ to $\check{\xi}_n$, and that $p \in]1, 2]$. It then follows by Cauchy–Schwartz inequality that

$$\begin{aligned} \left| \int_{\Omega} (\Delta_p^{K_n} \bar{u}_n(\mathbf{x}, t) - \Delta_p^{K_n} u(\mathbf{x}, t)) (\check{u}_n(\mathbf{x}, t) - \bar{u}_n(\mathbf{x}, t)) \, d\mathbf{x} \right| &\leq C \left(\|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^{p-1} \tau + \tau^p \right) \\ &\leq \frac{1}{6} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2 + C(\tau^{2/(3-p)} + \tau^p). \end{aligned}$$

Using Young inequality to bound the last term in (42), and combining the bounds on the three other terms, we have shown that

$$\begin{aligned} \frac{\partial}{\partial t} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2 &\leq \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2 + C \left(\tau^{2/(3-p)} + \tau^p + \|f_n - f\|_{L^2(\Omega)}^2 \right. \\ &\quad \left. + \begin{cases} \|K_n - K\|_{L^\infty,2(\Omega^2)}^2, & g \in L^2(\Omega) \\ \|K_n - K\|_{L^2(\Omega^2)}^2, & g \in L^\infty(\Omega) \end{cases} \right). \end{aligned}$$

Using the Gronwall's lemma and taking the square-root, we get

$$\begin{aligned} \|\check{u}_n - u\|_{C([0,T];L^2(\Omega))} &\leq \exp(T/2) \left(\|\mathbf{I}_n \mathbf{P}_n g - g\|_{L^2(\Omega)} \right. \\ &\quad \left. + CT^{1/2} \left(\tau^{1/(3-p)} + \tau^{p/2} + \|f_n - f\|_{L^2(\Omega)} + \begin{cases} \|\mathbf{I}_n \mathbf{P}_n K - K\|_{L^\infty,2(\Omega^2)} & g \in L^2(\Omega) \\ \|\mathbf{I}_n \mathbf{P}_n K - K\|_{L^2(\Omega^2)} & g \in L^\infty(\Omega) \end{cases} \right) \right). \end{aligned} \quad (43)$$

Since $1/2 < 1/(3-p) \leq p/2$ for $p \in]1, 2]$ the dependence on τ scales as $O(\tau^{1/(3-p)})$ for τ sufficiently small (or N large enough). Inserting (43) into

$$\sup_{k \in [N], t \in]t_{k-1}, t_k]} \|u_n^{k-1} - u(\cdot, t)\|_{L^2(\Omega)} = \|\bar{u}_n - u\|_{C([0,T];L^2(\Omega))} \leq \|\check{u}_n - u\|_{C([0,T];L^2(\Omega))} + C\tau, \quad (44)$$

completes the proof of the error bound.

(iii) Plug (6) into (40).

□

Remark 6.2. Error bounds in $L^p(\Omega)$ were derived in [30] for forward Euler discretization with bounded data (kernel and initial condition) and $f = 0$. Their rate is provided for the range $p \in]1, +\infty[$. However, we believe that their proof contains an invalid argument for $p \geq 2$.

Case $p = 1$: we now turn to the case $p = 1$, and consider the discrete system

$$\begin{cases} \frac{\mathbf{u}^k - \mathbf{u}^{k-1}}{\tau_{k-1}} = -\boldsymbol{\eta}^{k-1} + \mathbf{f}, & k \in [N], \\ \mathbf{u}^0 = \mathbf{g} \end{cases} \quad (\mathcal{P}_1^{\text{TDF}})$$

where

$$\boldsymbol{\eta}^k = - \sum_{j \in [n]^d} h_j \mathbf{K}_{ij} \text{sign}(\mathbf{u}_j^k - \mathbf{u}_i^k), \quad \text{and thus } \boldsymbol{\eta}^k \in \widehat{\Delta}_1^{\mathbf{K}} \mathbf{u}^k.$$

We consider the continuum extensions in space and/or time of $\{\mathbf{u}^k\}_{k \in [N]}$ as before, namely u_n^k , \check{u}_n and \bar{u}_n , $f_n = \mathbf{I}_n \mathbf{f}$, and the space-time continuum extension of $\{\boldsymbol{\eta}^k\}_{k \in [N]}$

$$\bar{\eta}_n(\mathbf{x}, t) = \sum_{k=1}^N (\mathbf{I}_n \boldsymbol{\eta}^{k-1})(\mathbf{x}) \chi_{]t_{k-1}, t_k]}(t) = - \int_{\Omega} \mathbf{I}_n \mathbf{K}(\mathbf{x}, \mathbf{y}) \text{sign}(\bar{u}_n(\mathbf{y}, t) - \bar{u}_n(\mathbf{x}, t)), \quad (\mathbf{x}, t) \in \Omega \times]0, T].$$

In view of Lemma 6.1, these extensions satisfy the evolution problem

$$\begin{cases} \frac{\partial}{\partial t} \check{u}_n(\mathbf{x}, t) = -\bar{\eta}_n(\mathbf{x}, t) + f_n(\mathbf{x}), & (\mathbf{x}, t) \in \Omega \times]0, T], \\ \check{u}_n(\mathbf{x}, 0) = \mathbf{I}_n \mathbf{g}(\mathbf{x}), & \mathbf{x} \in \Omega, \end{cases} \quad (45)$$

and

$$\bar{\eta}_n(\mathbf{x}, t) \in \Delta_1^{\mathbf{I}_n \mathbf{K}} \bar{u}_n(\mathbf{x}, t).$$

We have the following counterpart estimates of Lemma 6.2.

Lemma 6.4. Consider problem $(\mathcal{P}_1^{\text{TDF}})$ with kernel \mathbf{K} , data (\mathbf{f}, \mathbf{g}) and variable step-size

$$\tau_k = \frac{\alpha_k}{\max\left(\|\mathbf{I}_n \boldsymbol{\eta}^k - f_n\|_{L^2(\Omega)}, 1\right)}, \quad \text{where} \quad \sum_{k \in \mathbb{N}} \alpha_k^2 < +\infty.$$

Assume that $\mathbf{I}_n \mathbf{g} \in L^2(\Omega)$ and $\mathbf{I}_n \mathbf{K}$ satisfies **(H.1)–(H.3)**. Suppose also that for each $n \in \mathbb{N}$, \mathbf{f} is such that $(\mathcal{P}_p^{\text{TDF}})$ has a stationary solution \mathbf{u}^* and that $\sup_{n \in \mathbb{N}} \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)} < +\infty$. Then

$$\bar{u}_n(\cdot, t) \in L^2(\Omega), \forall t \in [0, T], \quad \text{and} \quad \sup_{t \in [0, T], n \in \mathbb{N}} \|\bar{u}_n(\cdot, t) - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)} < +\infty.$$

Remark 6.3. The condition on the time-step τ_k is reminiscent of subgradient descent and has been used in [30]. The assumptions on $(\mathbf{f}, \mathbf{g}, \mathbf{K})$ are again verified when $\mathbf{f} = 0$, $\mathbf{g} = P_n g$ and $\mathbf{K} = P_n K$, where $g \in L^2(\Omega)$ and K satisfies **(H.1)–(H.3)**.

Proof. Define the series $s_k \stackrel{\text{def}}{=} \sum_{i=0}^k \alpha_i^2$. As in Lemma 6.2, we proceed by induction using the monotonicity of the 1-Laplacian (Prop. 4.2(ii)). Indeed, since $f_n \in \Delta_p^{\mathbf{K}}(\mathbf{I}_n \mathbf{u}^*)$, we have

$$\|u_n^1 - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 = \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 - 2\tau_0 \int_{\Omega} (\Delta_p^{\mathbf{I}_n \mathbf{K}}(\mathbf{I}_n \mathbf{g})(\mathbf{x}) - \Delta_p^{\mathbf{I}_n \mathbf{K}}(\mathbf{I}_n \mathbf{u}^*)(\mathbf{x})) (\mathbf{I}_n \mathbf{g}(\mathbf{x}) - \mathbf{I}_n \mathbf{u}^*) \, d\mathbf{x} + \alpha_0^2.$$

By assumption on \mathbf{g}, \mathbf{u}^* , we can invoke Proposition 4.2(ii) to get

$$\|u_n^1 - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 \leq \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 + s_0.$$

Suppose now that, for any $k > 1$,

$$\|u_n^k - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 \leq \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 + s_{k-1},$$

and thus $u_n^k \in L^2(\Omega)$. We can then invoke again Proposition 4.2(ii) to see that

$$\begin{aligned} \|u_n^{k+1} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 &= \|u_n^k - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 - 2\tau_k \int_{\Omega} (\Delta_p^{\mathbf{I}_n \mathbf{K}}(u_n^k)(\mathbf{x}) - \Delta_p^{\mathbf{I}_n \mathbf{K}}(\mathbf{I}_n \mathbf{u}^*)(\mathbf{x})) (u_n^k(\mathbf{x}) - \mathbf{I}_n \mathbf{u}^*) \, d\mathbf{x} + \alpha_k^2 \\ &\leq \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 + s_k. \end{aligned}$$

This shows that for all $k \in \mathbb{N}$,

$$\|u_n^k - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 \leq \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)}^2 + s_{\infty},$$

and thus $\left\{ \|\mathbf{I}_n \mathbf{u}^k\|_{L^2(\Omega)} \right\}_{k \in [N]}$ is bounded. In turn, so is $\|\bar{u}_n(\cdot, t)\|_{L^2(\Omega)}$ for $t \in [0, T]$ by its definition. Moreover,

$$\begin{aligned} \sup_{t \in [0, T], n \in \mathbb{N}} \|\bar{u}_n(\cdot, t) - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)} &= \sup_{(n, N) \in \mathbb{N}^2, k \in [N]} \|u_n^k - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)} \\ &\leq \sup_{n \in \mathbb{N}} \|\mathbf{I}_n \mathbf{g} - \mathbf{I}_n \mathbf{u}^*\|_{L^2(\Omega)} + s_{\infty}^{1/2} < +\infty. \end{aligned}$$

□

We also have the following analogue of Lemma 6.3.

Lemma 6.5. *In addition to the assumptions of Lemma 6.4, suppose that $\sup_{n \in \mathbb{N}} \|\mathbf{I}_n \mathbf{K}\|_{L^\infty,1(\Omega^2)} < +\infty$. Then*

$$\sup_{t \in [0, T], n \in \mathbb{N}} \|\check{u}_n(\cdot, t) - \bar{u}_n(\cdot, t)\|_{L^2(\Omega)} \leq C\tau,$$

where C is a positive constant that does not depend on (n, N, T) .

Proof. Arguing as the beginning of Lemma 6.3, we get for any $t \in]t_{k-1}, t_k]$, $k \in \mathbb{N}$,

$$\|\check{u}_n(\cdot, t) - \bar{u}_n(\cdot, t)\|_{L^2(\Omega)} \leq \tau \|\bar{\eta}_n(\mathbf{x}, t) - f_n\|_{L^2(\Omega)}.$$

By Hölder inequality, we have

$$\begin{aligned} \|\bar{\eta}_n(\mathbf{x}, t)\|_{L^2(\Omega)}^2 &= \int_{\Omega} \left| \int_{\Omega} \mathbf{I}_n \mathbf{K}(\mathbf{x}, \mathbf{y}) \operatorname{sign}(\bar{u}_n(\mathbf{y}, t) - \bar{u}_n(\mathbf{x}, t)) \, d\mathbf{y} \right|^2 \, d\mathbf{x} \\ &\leq \int_{\Omega} \left(\int_{\Omega} \mathbf{I}_n \mathbf{K}(\mathbf{x}, \mathbf{y}) \, d\mathbf{y} \right)^2 \, d\mathbf{x} \leq \|\mathbf{I}_n \mathbf{K}\|_{L^\infty,1(\Omega^2)}^2. \end{aligned}$$

The same bound also holds on $\|f_n\|_{L^2(\Omega)}$. We then take the supremum over t and n to conclude. \square

Theorem 6.4. *Let u be a solution of (\mathcal{P}) with kernel K and data (f, g) where f is time-independent, and $\{\mathbf{u}^k\}_{k \in [N]}$ is the sequence generated by $(\mathcal{P}_1^{\text{TDF}})$ with $\mathbf{K} = P_n K$, $\mathbf{g} = P_n g$, $\mathbf{f} = P_n f$ and τ_k as prescribed in Lemma 6.4. Assume that K satisfies $(\mathbf{H.1})$, $(\mathbf{H.2})$ and $K \in L^{\infty,2}(\Omega^2)$, and that $f, g \in L^2(\Omega)$. Then, the following hold.*

- (i) u is the unique solution of (\mathcal{P}) , $\{\mathbf{u}^k\}_{k \in [N]}$ is uniquely defined and $\left\{ \|\mathbf{I}_n \mathbf{u}^k\|_{L^2(\Omega)} \right\}_{k \in [N]}$ is bounded (uniformly in n when $\mathbf{f} = \mathbf{0}$).
- (ii) We have the error estimate

$$\begin{aligned} \sup_{k \in [N], t \in]t_{k-1}, t_k]} \|\mathbf{I}_n \mathbf{u}^{k-1} - u(\cdot, t)\|_{L^2(\Omega)} &\leq \exp(T/2) \left(\|\mathbf{I}_n P_n g - g\|_{L^2(\Omega)} \right. \\ &\quad \left. + CT^{1/2} \left(\tau^{1/2} + \|f_n - f\|_{L^2(\Omega)} + \|\mathbf{I}_n P_n K - K\|_{L^2(\Omega^2)} \right) \right) \end{aligned} \tag{46}$$

where C is positive constant that depends only on K .

- (iii) If, moreover, $f, g \in \operatorname{Lip}(s, L^2(\Omega))$ and $K \in \operatorname{Lip}(s, L^2(\Omega^2))$, then

$$\sup_{k \in [N], t \in]t_{k-1}, t_k]} \|\mathbf{I}_n \mathbf{u}^{k-1} - u(\cdot, t)\|_{L^2(\Omega)} \leq C \exp(T/2) \left((1 + T^{1/2}) \delta_n^s + T^{1/2} \tau^{1/2} \right), \tag{47}$$

where C is positive constant that depends only on g, f, K and s .

Proof. C is any positive constant that may depend only on g, f, K and s , and that may be different at each line. We use the same notation as in the proof of Theorem 6.3.

- (i) Existence and uniqueness of u were proved in Theorem 5.1(i). Well-posedness of $\{\mathbf{u}^k\}_{k \in [N]}$ follows from Lemma 6.4 and Remark 6.1(iii).
- (ii) We have

$$\frac{\partial \check{\xi}_n(\mathbf{x}, t)}{\partial t} = \int_{\Omega} K_n(\mathbf{x}, \mathbf{y}) (\bar{w}_n(\mathbf{x}, \mathbf{y}, t) - w(\mathbf{x}, \mathbf{y}, t)) \, d\mathbf{y}$$

$$+ \int_{\Omega} (K_n(\mathbf{x}, \mathbf{y}) - K(\mathbf{x}, \mathbf{y}))w(\mathbf{x}, \mathbf{y}, t) \, d\mathbf{y} + (f_n(\mathbf{x}) - f(\mathbf{x})),$$

where w is the subgradient function associated to u (see Def. 4.2), and $\bar{w}_n(\mathbf{x}, \mathbf{y}, t) = \text{sign}(\bar{u}_n(\mathbf{y}, t) - \bar{u}_n(\mathbf{x}, t))$. Multiplying both sides by $\check{\xi}_n(\mathbf{x}, t)$, integrating and rearranging the terms, we get

$$\begin{aligned} \frac{1}{2} \frac{\partial}{\partial t} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2 &= \int_{\Omega^2} K_n(\mathbf{x}, \mathbf{y})(\bar{w}_n(\mathbf{x}, \mathbf{y}, t) - w(\mathbf{x}, \mathbf{y}, t))(\bar{u}_n(\mathbf{x}, t) - u(\mathbf{x}, t)) \, d\mathbf{x} \, d\mathbf{y} \\ &+ \int_{\Omega^2} (K_n(\mathbf{x}, \mathbf{y}) - K(\mathbf{x}, \mathbf{y}))w(\mathbf{x}, \mathbf{y}, t)\check{\xi}_n(\mathbf{x}, t) \, d\mathbf{x} \, d\mathbf{y} \\ &+ \int_{\Omega} K_n(\mathbf{x}, \mathbf{y})(\bar{w}_n(\mathbf{x}, \mathbf{y}, t) - w(\mathbf{x}, \mathbf{y}, t))(\check{u}_n(\mathbf{x}, t) - \bar{u}_n(\mathbf{x}, t)) \, d\mathbf{x} \, d\mathbf{y} \\ &+ \int_{\Omega} (f_n(\mathbf{x}) - f(\mathbf{x}))\check{\xi}_n(\mathbf{x}, t) \, d\mathbf{x}. \end{aligned} \tag{48}$$

As $u(\cdot, t) \in L^1$ and $\bar{u}_n(\cdot, t) \in L^2(\Omega) \subset L^1(\Omega)$ by Lemma 6.4, the monotonicity claim in Proposition 4.2(ii) yields that the first term in (48) is nonpositive. The second and third terms can be easily bounded as

$$\begin{aligned} \left| \int_{\Omega^2} (K_n(\mathbf{x}, \mathbf{y}) - K(\mathbf{x}, \mathbf{y}))w(\mathbf{x}, \mathbf{y}, t)\check{\xi}_n(\mathbf{x}, t) \, d\mathbf{x} \, d\mathbf{y} \right| &\leq \|K_n - K\|_{L^2(\Omega^2)} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)} \\ &\leq \frac{1}{4} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2 + \|K_n - K\|_{L^2(\Omega^2)}^2 \end{aligned}$$

and the third term using Lemma 6.5

$$\left| \int_{\Omega} K_n(\mathbf{x}, \mathbf{y})(\bar{w}_n(\mathbf{x}, \mathbf{y}, t) - w(\mathbf{x}, \mathbf{y}, t))(\check{u}_n(\mathbf{x}, t) - \bar{u}_n(\mathbf{x}, t)) \, d\mathbf{x} \, d\mathbf{y} \right| \leq 2\|K\|_{L^\infty, 2(\Omega^2)}^2 \tau.$$

Bounding the last term by Young inequality, we obtain

$$\frac{\partial}{\partial t} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2 \leq \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2 + 2\|f_n - f\|_{L^2(\Omega)}^2 + 2\|K_n - K\|_{L^2(\Omega^2)}^2 + C\tau.$$

Using the Gronwall’s lemma and (44), we get the claimed bound.

(iii) Insert (6) into (46).

□

6.2.2. Backward/Implicit Euler discretization

Forward Euler discretization was able to deal only with $p \in [1, 2]$. For backward Euler discretization, we will tackle $p \in]1, +\infty[$.

We consider the fully discrete problem with backward Euler time scheme

$$\begin{cases} \frac{\mathbf{u}^k - \mathbf{u}^{k-1}}{\tau_{k-1}} = -\widehat{\Delta}_p^{\mathbf{K}} \mathbf{u}^k + \mathbf{f}^k, & k \in [N], \\ \mathbf{u}^0 = \mathbf{g}, \end{cases} \tag{\mathcal{P}_p^{\text{TDB}}}$$

where $\mathbf{u}^k, \mathbf{f}^k \in \mathbb{R}^{n^d}$. This can also be written equivalently as

$$\mathbf{u}^k = J_{\tau_{k-1} \widehat{\Delta}_p^{\mathbf{K}}}(\mathbf{u}^{k-1} + \tau_{k-1} \mathbf{f}^k).$$

This is known as the proximal iteration, and is at the heart of so-called mild solutions as well as existence and uniqueness of solutions to (\mathcal{P}) through the theory of nonlinear semigroups [9, 10, 20, 36]. Denoting as before $u_n^k = \mathbf{I}_n \mathbf{u}^k$ and $f_n^k = \mathbf{I}_n \mathbf{f}^k$ the space continuum extensions of \mathbf{u}^k and \mathbf{f}^k , we also have

$$u_n^k = J_{\tau_{k-1} \Delta_p^{\mathbf{I}_n} \mathbf{K}}(u_n^{k-1} + \tau_{k-1} f_n^k).$$

We also let the time-space continuum extensions

$$\begin{aligned} \check{u}_n(\mathbf{x}, t) &= \frac{t_k - t}{\tau_{k-1}} u_n^{k-1}(\mathbf{x}) + \frac{t - t_{k-1}}{\tau_{k-1}} u_n^k(\mathbf{x}), & (\mathbf{x}, t) \in \Omega \times]t_{k-1}, t_k], k \in [N], \\ \bar{u}_n(\mathbf{x}, t) &= \sum_{k=1}^N u_n^k(\mathbf{x}) \chi_{]t_{k-1}, t_k]}(t) \quad \text{and} \quad \bar{f}_n(\mathbf{x}, t) = \sum_{k=1}^N f_n^k(\mathbf{x}) \chi_{]t_{k-1}, t_k]}(t), & (\mathbf{x}, t) \in \Omega \times]0, T]. \end{aligned}$$

Observe that the difference with the explicit Euler case lies in the definition of \bar{u}_n . From $(\mathcal{P}_p^{\text{TDB}})$ one clearly sees that \check{u}_n and \bar{u}_n then satisfy again (39) with $\bar{f}_n(\mathbf{x}, t)$ replacing $f_n(\mathbf{x})$.

The following estimates holds.

Lemma 6.6. *Consider problem $(\mathcal{P}_p^{\text{TDB}})$ with kernel \mathbf{K} and data (\mathbf{f}, \mathbf{g}) and step-sizes $\tau_k > 0$ for all k . Assume that $\mathbf{I}_n \mathbf{K}$ satisfies (H.1)–(H.3), that $\mathbf{I}_n \mathbf{g} \in L^{\max(p,q)}(\Omega)$, for some $q \in [1, +\infty]$, and $\sup_{n \in \mathbb{N}} \|\mathbf{I}_n \mathbf{g}\|_{L^q(\Omega)} < +\infty$, and that $\bar{f}_n \in L^1([0, T]; L^{\max(p,q)}(\Omega))$ and $\sup_{n \in \mathbb{N}} \|\bar{f}_n\|_{L^1([0, T]; L^q(\Omega))} < +\infty$. Then*

$$\bar{u}_n(\cdot, t) \in L^{\max(p,q)}(\Omega), \forall t \in [0, T], \quad \text{and} \quad \sup_{t \in [0, T], n \in \mathbb{N}} \|\bar{u}_n(\cdot, t)\|_{L^q(\Omega)} < +\infty.$$

Proof. Recall from Proposition 4.1(vii) that $J_{\lambda \Delta_p^{\mathbf{I}_n \mathbf{K}}}$, $\lambda > 0$, is single-valued on $L^p(\Omega)$ and non-expansive on $L^q(\Omega)$ for all $q \in [1, +\infty]$. Therefore, by induction, we have that for any $k \in [N]$,

$$\begin{aligned} \|u_n^k\|_{L^p(\Omega)} &\leq \|\mathbf{I}_n \mathbf{g}\|_{L^p(\Omega)} + \sum_{i=0}^k \tau_i \|f_n^i\|_{L^p(\Omega)} \leq \|\mathbf{I}_n \mathbf{g}\|_{L^p(\Omega)} + \sum_{i=0}^N \tau_i \|f_n^i\|_{L^p(\Omega)} \\ &= \|\mathbf{I}_n \mathbf{g}\|_{L^p(\Omega)} + \|\bar{f}_n\|_{L^1([0, T]; L^p(\Omega))}. \end{aligned}$$

Thus $u_n^k \in L^p(\Omega)$, for all $k \in [N]$. In turn, $J_{\tau_k \Delta_p^{\mathbf{I}_n \mathbf{K}}}(u_n^k)$ is single-valued for all k , and arguing as above, its non-expansiveness yields

$$\|u_n^k\|_{L^q(\Omega)} \leq \|\mathbf{I}_n \mathbf{g}\|_{L^q(\Omega)} + \|\bar{f}_n\|_{L^1([0, T]; L^q(\Omega))}.$$

Taking the supremum over k and n and using the definition of \bar{u}_n and the assumptions on \mathbf{g} and \mathbf{f} , we conclude. \square

Lemma 6.7. *Suppose that the assumptions of Lemma 6.6 are satisfied with $q = 2$ when $p \in]1, 2]$, $q = 2(p - 1)$ when $p \geq 2$. Assume in addition that $\sup_{n \in \mathbb{N}} \|\mathbf{I}_n \mathbf{K}\|_{L^{\infty,1}(\Omega^2)} < +\infty$ and $\sup_{n \in \mathbb{N}} \|\bar{f}_n\|_{\text{BV}([0, T]; L^2(\Omega))} < +\infty$. Then*

$$\sup_{t \in [0, T], n \in \mathbb{N}} \|\check{u}_n(\cdot, t) - \bar{u}_n(\cdot, t)\|_{L^2(\Omega)} \leq C\tau,$$

where C is a positive constant that does not depend on (n, N, T) .

Proof. For $t \in]t_{k-1}, t_k]$, $k \in \mathbb{N}$, we have

$$\begin{aligned} \|\check{u}_n(\cdot, t) - \bar{u}_n(\cdot, t)\|_{L^2(\Omega)} &= (t_k - t) \left\| \frac{u_n^{k-1} - u_n^k}{\tau_{k-1}} \right\|_{L^2(\Omega)} \\ &= (t_k - t) \|\Delta_p^{\mathbf{I}_n \mathbf{K}} u_n^k - f_n^k\|_{L^2(\Omega)} \\ &\leq \tau \|\Delta_p^{\mathbf{I}_n \mathbf{K}} u_n^k - f_n^k\|_{L^2(\Omega)} \\ &= \tau \|\Delta_p^{\mathbf{I}_n \mathbf{K}} \bar{u}_n(\cdot, t_k) - \bar{f}_n(\cdot, t_k)\|_{L^2(\Omega)} \end{aligned}$$

$$\begin{aligned}
&\leq \tau \left(\|\Delta_p^{\mathbf{I}_n \mathbf{K}} \bar{u}_n(\cdot, t_k)\|_{L^2(\Omega)} + \sum_{i=1}^k \|\bar{f}_n(\cdot, t_i) - \bar{f}_n(\cdot, t_{i-1})\|_{L^2(\Omega)} + \|\bar{f}_n(\cdot, 0)\|_{L^2(\Omega)} \right) \\
&\leq \tau \left(\|\Delta_p^{\mathbf{I}_n \mathbf{K}} \bar{u}_n(\cdot, t_k)\|_{L^2(\Omega)} + \text{Var}_q(\bar{f}_n) + \|\bar{f}_n(\cdot, 0)\|_{L^2(\Omega)} \right) \\
&= \tau \left(\|\Delta_p^{\mathbf{I}_n \mathbf{K}} \bar{u}_n(\cdot, t_k)\|_{L^2(\Omega)} + \|\bar{f}_n\|_{\text{BV}([0, T]; L^2(\Omega))} \right). \tag{49}
\end{aligned}$$

For $p \in]1, 2]$, we have from (18) that

$$\|\Delta_p^{\mathbf{I}_n \mathbf{K}} \bar{u}_n(\cdot, t_k)\|_{L^2(\Omega)} \leq 2^{p/2} C_2^{1/2} \|K\|_{L^\infty, 1(\Omega^2)} \|\bar{u}_n(\cdot, t)\|_{L^2(\Omega)}^{p-1}.$$

For $p \geq 2$, it is easy to show with simple arguments as before that

$$\|\Delta_p^{\mathbf{I}_n \mathbf{K}} \bar{u}_n(\cdot, t_k)\|_{L^2(\Omega)} \leq 2^{p-3/2} \|K\|_{L^\infty, 1(\Omega^2)} \|\bar{u}_n(\cdot, t)\|_{L^2(\Omega)}^{p-1}.$$

Inserting the last two estimates in (49), taking the supremum over t and n over both sides, and applying Lemma 6.6, we conclude. \square

Remark 6.4. As observed in the case of explicit time-discretization the uniform (over n) boundedness assumption made in the last two lemmas hold true if $\mathbf{g} = P_n g$, $\mathbf{K} = P_n K$ and $\mathbf{f}^k = \tau_k^{-1} \int_{t_{k-1}}^{t_k} P_n f(\cdot, t) dt$, where g , f and K verify simple assumptions. Indeed, in this case, we have thanks to Lemma 2.1 that for any $q \in [1, +\infty]$,

$$\begin{aligned}
\sup_{n \in \mathbb{N}} \|\mathbf{I}_n \mathbf{g}\|_{L^q(\Omega)} &\leq \|g\|_{L^q(\Omega)}, & \sup_{n \in \mathbb{N}} \|\mathbf{I}_n \mathbf{K}\|_{L^\infty, q(\Omega^2)} &\leq \|K\|_{L^\infty, q(\Omega^2)}, \\
\sup_{n \in \mathbb{N}} \|\bar{f}_n\|_{L^1([0, T]; L^q(\Omega))} &\leq \|f\|_{L^1([0, T]; L^q(\Omega))} & \text{and} & \sup_{n \in \mathbb{N}} \|\bar{f}_n\|_{\text{BV}([0, T]; L^q(\Omega))} &\leq \|f\|_{\text{BV}([0, T]; L^q(\Omega))}.
\end{aligned}$$

In fact, the condition $f \in \text{BV}([0, T]; L^q(\Omega))$ is sufficient to ensure that

$$\sup_{n \in \mathbb{N}} \|\bar{f}_n\|_{L^1([0, T]; L^q(\Omega))} < +\infty \quad \text{and} \quad \sup_{n \in \mathbb{N}} \|\bar{f}_n\|_{\text{BV}([0, T]; L^q(\Omega))} < +\infty.$$

Indeed, arguing as in Lemma A.1 of [15], this conditions implies $f \in L^\infty([0, T]; L^q(\Omega))$. In turn, using Lemma 2.1, we get

$$\begin{aligned}
\|\bar{f}_n\|_{L^1([0, T]; L^q(\Omega))} &\leq \|f\|_{L^1([0, T]; L^q(\Omega))} \leq \|f\|_{L^\infty([0, T]; L^q(\Omega))} \\
&\leq \|f(\cdot, 0)\|_{L^q(\Omega)} + \text{Var}_q(f) = \|f\|_{\text{BV}([0, T]; L^q(\Omega))}.
\end{aligned}$$

We are now in position to state the error bound for the fully discrete problem with backward/implicit Euler time discretization.

Theorem 6.5. *Suppose that $p \in]1, +\infty[$. Let u be a solution of (\mathcal{P}) with kernel K and data (f, g) , and $\{\mathbf{u}^k\}_{k \in [N]}$ is the sequence generated by $(\mathcal{P}_p^{\text{TDB}})$ with $\mathbf{K} = P_n K$, $\mathbf{g} = P_n g$, $\mathbf{f}^k = \tau_k^{-1} \int_{t_{k-1}}^{t_k} P_n f(\cdot, t) dt$. Assume that K satisfies (H.1), (H.2) and $K \in L^{\infty, 2}(\Omega^2)$, and that f, g satisfy either one of the conditions (a), (b) or (c) in Theorem 5.1, and that $f \in \text{BV}([0, T]; L^2(\Omega))$. Then, the following hold.*

- (i) u is the unique solution of (\mathcal{P}) , $\{\mathbf{u}^k\}_{k \in [N]}$ is uniquely defined and $\left\{ \|\mathbf{I}_n \mathbf{u}^k\|_{L^2(\Omega)} \right\}_{k \in [N]}$ is bounded uniformly in n .

(ii) We have the error estimate

$$\sup_{k \in [N], t \in]t_{k-1}, t_k]} \|\mathbf{I}_n \mathbf{u}^k - u(\cdot, t)\|_{L^2(\Omega)} \leq \exp(T/2) \left(\|\mathbf{I}_n \mathbf{P}_n g - g\|_{L^2(\Omega)} + \|\bar{f}_n - f\|_{L^1([0, T]; L^2(\Omega))} \right) + CT^{1/2} \begin{cases} \tau^{1/(3-p)} + \|\mathbf{I}_n \mathbf{P}_n K - K\|_{L^\infty, 2(\Omega^2)} & \text{under (a)} \\ \tau^{p/(2p-1)} + \|\mathbf{I}_n \mathbf{P}_n K - K\|_{L^\infty, 2(\Omega^2)} & \text{under (b)} \\ \tau^{1/(3-p)} + \|\mathbf{I}_n \mathbf{P}_n K - K\|_{L^2(\Omega^2)} & \text{under (c) when } p \in]1, 2] \\ \tau + \|\mathbf{I}_n \mathbf{P}_n K - K\|_{L^2(\Omega^2)} & \text{under (c) when } p \geq 2. \end{cases} \quad (50)$$

for τ sufficiently small, where C is positive constant that depends only on p, g, f and K .

(iii) If, moreover, $g \in L^\infty(\Omega) \cap \text{Lip}(s, L^2(\Omega))$, $K \in \text{Lip}(s, L^2(\Omega^2))$, and $f \in L^1([0, T]; L^\infty(\Omega)) \cap \text{Lip}(s, L^2(\Omega \times [0, T]))$ then

$$\sup_{k \in [N], t \in]t_{k-1}, t_k]} \|\mathbf{I}_n \mathbf{u}^k - u(\cdot, t)\|_{L^2(\Omega)} \leq C \exp(T/2) \left((1 + T^{1/2}) \delta_n^s + T^{1/2} \begin{cases} \tau^{\min(s, 1/(3-p))} & \text{when } p \in]1, 2] \\ \tau^s & \text{when } p \geq 2 \end{cases} \right). \quad (51)$$

for τ sufficiently small, where C is positive constant that depends only on p, g, f, K and s . The term τ^s in the dependence on τ disappears when f is time-independent.

Proof. In the proof, C is any positive constant that may depend solely on p, g, f, K and/or s , and that may be different at each line.

- (i) Existence and uniqueness of u were proved in Theorem 5.1(i). Well-posedness of the sequence $\{\mathbf{u}^k\}_{k \in [N]}$ is a consequence of Lemma 6.6 and Remark 6.4.
- (ii) For $p \in]1, 2]$, the proof of the error bound is exactly the same as that of (40) in Theorem 6.3 using the modified definition of \bar{u}_n and that now f is time-dependent, and thus we replace f_n there by \bar{f}_n . We also denote $g_n = \mathbf{I}_n \mathbf{P}_n g$ and $K_n = \mathbf{I}_n \mathbf{P}_n K$. For the $p \geq 2$, the argument is also similar, and the main change consists in bounding appropriately the third term in (42). We then invoke Lemma 6.7 to show that

$$\left| \int_{\Omega} (\Delta_p^{K_n} \bar{u}_n(\mathbf{x}, t) - \Delta_p^{K_n} u(\mathbf{x}, t)) (\bar{u}_n(\mathbf{x}, t) - u(\mathbf{x}, t)) \, d\mathbf{x} \right| \leq C \|\Delta_p^{K_n} \bar{u}_n(\cdot, t) - \Delta_p^{K_n} u(\cdot, t)\|_{L^2(\Omega)} \tau,$$

where C is indeed a finite constant owing to the assumption on f and Remark 6.4. We now use Lemma 4.1(ii) to get the bound

$$\begin{aligned} & \|\Delta_p^{K_n} \bar{u}_n(\cdot, t) - \Delta_p^{K_n} u(\cdot, t)\|_{L^2(\Omega)}^2 \\ &= \int_{\Omega} \left| \int_{\Omega} K_n(\mathbf{x}, \mathbf{y}) (\Psi(\bar{u}_n(\mathbf{y}, t) - \bar{u}_n(\mathbf{x}, t)) - \Psi(u(\mathbf{y}, t) - u(\mathbf{x}, t))) \, d\mathbf{y} \right|^2 \, d\mathbf{x} \\ &\leq \int_{\Omega} \left(\int_{\Omega} K_n(\mathbf{x}, \mathbf{y}) |\bar{\xi}_n(\mathbf{y}, t) - \bar{\xi}_n(\mathbf{x}, t)| (|\bar{u}_n(\mathbf{y}, t) - \bar{u}_n(\mathbf{x}, t)| + |u(\mathbf{y}, t) - u(\mathbf{x}, t)|)^{p-2} \, d\mathbf{y} \right)^2 \, d\mathbf{x}. \quad (52) \end{aligned}$$

For case (c), we infer from Lemma 6.6 (with $q = +\infty$) and Lemma 2.1 that

$$\begin{aligned}
 & \|\Delta_p^{K_n} \bar{u}_n(\cdot, t) - \Delta_p^{K_n} u(\cdot, t)\|_{L^2(\Omega)}^2 \\
 & \leq \left(4\left(\|g\|_{L^\infty(\Omega)} + \|f\|_{L^1([0,T];L^\infty(\Omega))}\right)\right)^{2(p-2)} \int_{\Omega} \left(\int_{\Omega} K_n(\mathbf{x}, \mathbf{y}) |\bar{\xi}_n(\mathbf{y}, t) - \bar{\xi}_n(\mathbf{x}, t)| d\mathbf{y}\right)^2 d\mathbf{x} \\
 & \leq \left(4\left(\|g\|_{L^\infty(\Omega)} \|f\|_{L^1([0,T];L^\infty(\Omega))}\right)\right)^{2(p-2)} \|K\|_{L^\infty,2(\Omega^2)} \int_{\Omega^2} K_n(\mathbf{x}, \mathbf{y}) |\bar{\xi}_n(\mathbf{y}, t) - \bar{\xi}_n(\mathbf{x}, t)|^2 d\mathbf{x} d\mathbf{y} \\
 & = 4\left(4\left(\|g\|_{L^\infty(\Omega)} \|f\|_{L^1([0,T];L^\infty(\Omega))}\right)\right)^{2(p-2)} \|K\|_{L^\infty,2(\Omega^2)} \int_{\Omega^2} K_n(\mathbf{x}, \mathbf{y}) |\bar{\xi}_n(\mathbf{x}, t)|^2 d\mathbf{x} d\mathbf{y} \\
 & \leq 4\left(4\left(\|g\|_{L^\infty(\Omega)} \|f\|_{L^1([0,T];L^\infty(\Omega))}\right)\right)^{2(p-2)} \|K\|_{L^\infty,2(\Omega^2)}^2 \|\bar{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2. \tag{53}
 \end{aligned}$$

It then follows by Cauchy–Schwartz inequality that

$$\begin{aligned}
 \left| \int_{\Omega} (\Delta_p^{K_n} \bar{u}_n(\mathbf{x}, t) - \Delta_p^{K_n} u(\mathbf{x}, t)) (\bar{u}_n(\mathbf{x}, t) - u(\mathbf{x}, t)) d\mathbf{x} \right| & \leq C \|\bar{\xi}_n(\cdot, t)\|_{L^2(\Omega)} \tau \\
 & \leq C \left(\|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)} \tau + \tau^2\right) \\
 & \leq \frac{1}{6} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2 + C\tau^2.
 \end{aligned}$$

Inserting this in (42), using again Young inequality for the last term, we have shown that when $p \geq 2$ and (c) holds,

$$\frac{\partial}{\partial t} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2 \leq \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2 + C\left(\tau^2 + \|\bar{f}_n(\cdot, t) - f(\cdot, t)\|_{L^2(\Omega)}^2 + \|K_n - K\|_{L^2(\Omega^2)}^2\right).$$

Using the Gronwall’s lemma, taking the square-root and using (44), we get the error bound in this case. It remains to consider the case (b), when $p \geq 2$. For this, we embark from (52), and use the continuity of Ψ in Lemma 4.1(i) (see (12)) with $\alpha = 1/p$. Combining this with Jensen and Hölder inequalities, we get

$$\begin{aligned}
 & \|\Delta_p^{K_n} \bar{u}_n(\cdot, t) - \Delta_p^{K_n} u(\cdot, t)\|_{L^2(\Omega)}^2 \\
 & \leq \|K\|_{L^\infty,1(\Omega^2)} \int_{\Omega^2} \left(K_n(\mathbf{x}, \mathbf{y}) |\bar{\xi}_n(\mathbf{y}, t) - \bar{\xi}_n(\mathbf{x}, t)|\right)^{2/p} \\
 & \quad \times (|\bar{u}_n(\mathbf{y}, t) - \bar{u}_n(\mathbf{x}, t)| + |u(\mathbf{y}, t) - u(\mathbf{x}, t)|)^{2(p-1)-2/p} d\mathbf{x} d\mathbf{y} \\
 & \leq \|K\|_{L^\infty,1(\Omega^2)} \int_{\Omega^2} \left(K_n(\mathbf{x}, \mathbf{y}) |\bar{\xi}_n(\mathbf{y}, t) - \bar{\xi}_n(\mathbf{x}, t)|\right)^{2/p} \\
 & \quad \times \left((K_n(\mathbf{x}, \mathbf{y}))^{(p-1)/p} (|\bar{u}_n(\mathbf{y}, t) - \bar{u}_n(\mathbf{x}, t)| + |u(\mathbf{y}, t) - u(\mathbf{x}, t)|)^{2(p-1)-2/p}\right) d\mathbf{x} d\mathbf{y} \\
 & \leq \|K\|_{L^\infty,1(\Omega^2)} \left(\int_{\Omega^2} K_n(\mathbf{x}, \mathbf{y}) |\bar{\xi}_n(\mathbf{y}, t) - \bar{\xi}_n(\mathbf{x}, t)|^2 d\mathbf{x} d\mathbf{y}\right)^{1/p} \\
 & \quad \times \left(\int_{\Omega^2} K_n(\mathbf{x}, \mathbf{y}) (|\bar{u}_n(\mathbf{y}, t) - \bar{u}_n(\mathbf{x}, t)| + |u(\mathbf{y}, t) - u(\mathbf{x}, t)|)^{2p-2/(p-1)} d\mathbf{x} d\mathbf{y}\right)^{(p-1)/p} \\
 & \leq \|K\|_{L^\infty,1(\Omega^2)} \left(4 \int_{\Omega^2} K_n(\mathbf{x}, \mathbf{y}) |\bar{\xi}_n(\mathbf{x}, t)|^2 d\mathbf{x} d\mathbf{y}\right)^{1/p} \\
 & \quad \times \left(2^{2p-2/(p-1)} \int_{\Omega^2} K_n(\mathbf{x}, \mathbf{y}) (|\bar{u}_n(\mathbf{x}, t)| + |u(\mathbf{x}, t)|)^{2p-2/(p-1)} d\mathbf{x} d\mathbf{y}\right)^{(p-1)/p}.
 \end{aligned}$$

$$\begin{aligned} &\leq 4\|K\|_{L^\infty,1(\Omega^2)}^2 \left(\int_{\Omega^2} |\bar{\xi}_n(\mathbf{x}, t)|^2 \, d\mathbf{x} \, d\mathbf{y} \right)^{1/p} \\ &\quad \times \left(\int_{\Omega^2} (|\bar{u}_n(\mathbf{x}, t)| + |u(\mathbf{x}, t)|)^{2p-2/(p-1)} \, d\mathbf{x} \, d\mathbf{y} \right)^{(p-1)/p}. \end{aligned}$$

Observe that $L^{2p-2/(p-1)}(\Omega) \subset L^{2(p-1)}(\Omega)$, hence by Hölder inequality and Lemma 6.6 with $q = 2(p - 1)$ and Lemma 2.1, the last term in the above display can be bounded as

$$\begin{aligned} \left(\int_{\Omega^2} (|\bar{u}_n(\mathbf{x}, t)| + |u(\mathbf{x}, t)|)^{2p-2/(p-1)} \, d\mathbf{x} \, d\mathbf{y} \right)^{(p-1)/p} &\leq \| |\bar{u}_n(\mathbf{x}, t)| + |u(\mathbf{x}, t)| \|_{L^{2(p-1)}(\Omega)}^{2(p-1)-2/p} \\ &\leq \left(\|g\|_{L^{2(p-1)}(\Omega)} + \|f\|_{L^1([0,T];L^{2(p-1)}(\Omega))} \right)^{2(p-1)-2/p}. \end{aligned}$$

We then arrive at

$$\| \Delta_p^{K_n} \bar{u}_n(\cdot, t) - \Delta_p^{K_n} u(\cdot, t) \|_{L^2(\Omega)}^2 \leq C \|K\|_{L^\infty,1(\Omega^2)}^2 \| \bar{\xi}_n \|_{L^2(\Omega)}^{2/p}.$$

Hence

$$\begin{aligned} \left| \int_{\Omega} (\Delta_p^{K_n} \bar{u}_n(\mathbf{x}, t) - \Delta_p^{K_n} u(\mathbf{x}, t)) (\check{u}_n(\mathbf{x}, t) - \bar{u}_n(\mathbf{x}, t)) \, d\mathbf{x} \right| &\leq C \| \bar{\xi}_n(\cdot, t) \|_{L^2(\Omega)}^{1/p} \tau \\ &\leq C \left(\| \check{\xi}_n(\cdot, t) \|_{L^2(\Omega)}^{1/p} \tau + \tau^{(p+1)/p} \right) \\ &\leq \frac{1}{6} \| \check{\xi}_n(\cdot, t) \|_{L^2(\Omega)}^2 + C(\tau^{2p/(2p-1)} + \tau^{(p+1)/p}). \end{aligned}$$

Inserting this into (42), using again Young inequality for the last term,

$$\begin{aligned} \frac{\partial}{\partial t} \| \check{\xi}_n(\cdot, t) \|_{L^2(\Omega)}^2 &\leq \| \check{\xi}_n(\cdot, t) \|_{L^2(\Omega)}^2 + C \left(\tau^{2p/(2p-1)} + \tau^{(p+1)/p} + \| \bar{f}_n(\cdot, t) - f(\cdot, t) \|_{L^2(\Omega)}^2 \right. \\ &\quad \left. + \| K_n - K \|_{L^2(\Omega^2)}^2 \right). \end{aligned}$$

Hence, using the Gronwall’s lemma, taking the square-root and using (44), we get the error bound in this case, after observing that the dependence on τ scales as $O(\tau^{p/(2p-1)})$ for τ sufficiently small (or N large enough) since $1/2 < p/(2p - 1) \leq (p + 1)/(2p)$ for $p \geq 2$.

(iii) Plug (6) into (50) after observing that

$$\| \bar{f}_n - f \|_{L^1([0,T];L^2(\Omega))} \leq T^{1/2} \| \bar{f}_n - f \|_{L^2([0,T];L^2(\Omega))} = T^{1/2} \| \bar{f}_n - f \|_{L^2(\Omega \times [0,T])} \leq CT^{1/2} \max(\tau^s, \delta_n^s).$$

For the scaling in τ , we use that $s \in]0, 1]$.

□

Another way to derive error bounds for $(\mathcal{P}_p^{\text{TDB}})$ is as follows. To lighten notation, denote $g_n = \mathbf{I}_n \mathbf{P}_n g$, $f_n(\cdot, t) = \mathbf{I}_n \mathbf{P}_n f(\cdot, t)$ for $t \in [0, T]$, and $K_n = \mathbf{I}_n \mathbf{P}_n \mathbf{K}$. Let u_n be a solution to (\mathcal{P}) with data (f_n, g_n) and kernel K_n . Under the assumptions of Theorem 6.5 on (f, g, K) , u_n is unique. Then one has

$$\| \check{u}_n - u \|_{C([0,T];L^2(\Omega))} \leq \| \check{u}_n - u_n \|_{C([0,T];L^2(\Omega))} + \| u_n - u \|_{C([0,T];L^2(\Omega))}.$$

Theorem 5.1 provides a bound on the last term of the right-hand side in the above display, which captures the space-discretization error. Bounds for the first term, which corresponds to the time-discretization error, were

derived in $C([0, T]; L^p(\Omega))$ by Crandall and Liggett in their seminal paper [20] for constant time step-size and $f = 0$, and then extended to non-uniform time partitions in [36], see also [41]. More precisely, using Theorem 1 of [41] and the fact that $\check{u}_n(\cdot, 0) = u_n(\cdot, 0) = g_n$, the following bound holds

$$\|\check{u}_n - u_n\|_{C([0, T]; L^p(\Omega))} \leq \|\bar{f}_n - f_n\|_{L^1([0, T]; L^p(\Omega))} + 2T^{1/2} \left(\|f_n^1 - \Delta_p^{K_n} g_n\|_{L^p(\Omega)} + \text{Var}_p(\bar{f}_n) \right) \tau^{1/2}.$$

The first term can be bounded as follows (for constant step-size to simplify)

$$\begin{aligned} \|\bar{f}_n - f_n\|_{L^1([0, T]; L^p(\Omega))} &= \sum_{k=1}^N \int_{t_{k-1}}^{t_k} \left\| \tau_k^{-1} \int_{t_{k-1}}^{t_k} f_n(\cdot, s) \, ds - f_n(\cdot, t) \right\|_{L^p(\Omega)} \, dt \\ &\leq \tau^{-1} \sum_{k=1}^N \int_{t_{k-1}}^{t_k} \int_{t_{k-1}}^{t_k} \|f(\cdot, s) - f(\cdot, t)\|_{L^p(\Omega)} \, ds \, dt \\ &\leq \tau^{-1} \int_{-\tau}^{\tau} \left(\int_0^T \|f(\cdot, t+s) - f(\cdot, t)\|_{L^p(\Omega)} \, dt \right) \, ds \\ &\leq \tau^{-1} \int_{-\tau}^{\tau} s \text{Var}_p(f) \, ds = \tau \text{Var}_p(f), \end{aligned}$$

where we used Lemma 2.1 in the first inequality and Lemma A.1 of [15] in the last one. Overall, this shows that the time discretization error $\|\check{u}_n - u_n\|_{C([0, T]; L^p(\Omega))}$ scales as $O\left((T\tau)^{1/2}\right)$ for τ sufficiently small. The rate $O(\tau^{1/2})$ is known to be optimal for general accretive operators in Banach spaces (see [41]). In turn, by standard comparisons of $L^q(\Omega)$ norms (assuming that (c) holds so that boundedness of \check{u}_n and u_n is in force), this strategy gives us a bound which scales as

$$\|\check{u}_n - u_n\|_{C([0, T]; L^p(\Omega))} = \begin{cases} O(\tau^{1/2}) & p \geq 2, \\ O(\tau^{p/4}) & p \in]1, 2]. \end{cases}$$

This is strictly worse than the rates in τ obtained from (50). There is however no contradiction in this and the reason is that the strategy outlined above is too general and does not exploit all properties of the operator Δ_p^K among which its continuity that was a key to derive better rates in τ . In this sense, our present results are optimal. We also remark that our rates are consistent with those in [30] for $p \geq 2$.

7. APPLICATION TO RANDOM GRAPH SEQUENCES

In this section, we study continuum limits of fully discrete problems on the random graph model of Definition 3.1 with backward/implicit Euler time discretization. Explicit discretization can also be treated following our results in Section 6.2.1, but we will not elaborate further on it for the sake of brevity.

Recall the notations in Section 3, in which case we now set $\Omega = [0, 1]$. Recall also the the construction of the random graph model in Definition 3.1 where each edge (i, j) is independently set to 1 with probability (7). This entails that the random matrix $\mathbf{\Lambda}$ is symmetric. However, it is worth emphasizing that the entries of $\mathbf{\Lambda}$ are not independent, but only the entries in each row are mutually independent³. This observation will be instrumental in deducing our error bound.

We consider the fully discrete on K -random graphs $\mathbf{G}(n, K, \rho_n)$ with backward Euler time scheme

$$\begin{cases} \frac{\mathbf{u}^k - \mathbf{u}^{k-1}}{\tau_{k-1}} = \frac{1}{\rho_n n} \sum_{j: (i, j) \in E(\mathbf{G}(n, K, \rho_n))} \Psi(\mathbf{u}_j^k - \mathbf{u}_i^k) + \mathbf{f}^k, & k \in [N], \\ \mathbf{u}^0 = \mathbf{g}, \end{cases} \quad (\mathcal{P}_p^{\text{TDB}, \mathbf{G}})$$

³This feature was already used in the proof of Lemma 3.1.

where $\mathbf{u}^k, \mathbf{f}^k \in \mathbb{R}^n$. It is important to keep in mind that, since $\mathbf{G}(n, K, \rho_n)$ is a random variable taking values in the set of simple graphs, the evolution problem $(\mathcal{P}_p^{\text{TDB}, \mathbf{G}})$ must be understood in this sense, *i.e.*, the \mathbf{u}^k 's are random variables. Observe that the normalization in $(\mathcal{P}_p^{\text{TDB}, \mathbf{G}})$ by $\rho_n n$ corresponds to the average degree (see Sect. 3.2 for details).

Problem $(\mathcal{P}_p^{\text{TDB}, \mathbf{G}})$ can be equivalently written as

$$\begin{cases} \frac{\mathbf{u}^k - \mathbf{u}^{k-1}}{\tau_{k-1}} = -\widehat{\Delta}_p^\Lambda \mathbf{u}^k + \mathbf{f}^k, & k \in [N], \\ \mathbf{u}^0 = \mathbf{g}. \end{cases}$$

We define the time-space continuum extensions \check{u}_n and \bar{u}_n and as in Section 6.2.2. One then sees that they satisfy

$$\begin{cases} \frac{\partial}{\partial t} \check{u}_n(x, t) = -\Delta_p^{\mathbf{I}_n \Lambda} \bar{u}_n(x, t) + \bar{f}_n(x, t), & (x, t) \in \Omega \times]0, T], \\ \check{u}_n(x, 0) = \mathbf{I}_n \mathbf{g}(x), & x \in \Omega. \end{cases} \tag{54}$$

Toward our goal of establishing error bounds, we define \mathbf{v} as the solution of the fully discrete problem $(\mathcal{P}_p^{\text{TDB}})$ with data (\mathbf{f}, \mathbf{g}) and discrete kernel $\widehat{\mathbf{K}}$. Its time-space continuum extensions, \check{v}_n and \bar{v}_n , defined similarly as above, fulfill

$$\begin{cases} \frac{\partial}{\partial t} \check{v}_n(x, t) = -\Delta_p^{\mathbf{I}_n \widehat{\mathbf{K}}} \bar{v}_n(x, t) + \bar{f}_n(x, t), & (x, t) \in \Omega \times]0, T], \\ \check{v}_n(x, 0) = \mathbf{I}_n \mathbf{g}(x), & x \in \Omega. \end{cases} \tag{55}$$

We have

$$\|\check{u}_n - u\|_{C([0, T]; L^2(\Omega))} \leq \|\check{u}_n - \check{v}_n\|_{C([0, T]; L^2(\Omega))} + \|\check{v}_n - u\|_{C([0, T]; L^2(\Omega))}. \tag{56}$$

This bound is composed of two terms: the first one captures the error of random sampling, and the second that of (space and time) discretization. We start by bounding the first term by comparing (54) and (55).

Lemma 7.1. *Assume that $(\mathbf{f}^k, \mathbf{g}, \mathbf{K}, f, g, K)$ verify the assumptions of Theorem 6.5. Assume also that $\rho_n \rightarrow 0$ and $n\rho_n = \omega((\log n)^\gamma)$ for some $\gamma > 1$. Then, for any $\beta \in]0, 1[$,*

$$\|\check{u}_n - \check{v}_n\|_{C([0, T]; L^2(\Omega))} \leq C \exp(T/2) T^{1/2} \left((\rho_n n)^{-\beta/2} + \begin{cases} \tau^{1/(3-p)} & p \in]1, 2], \\ \tau & p \geq 2, \end{cases} \right) \tag{57}$$

with probability at least $1 - (\rho_n n)^{-(1-\beta)}$. In particular,

$$\|\check{u}_n - \check{v}_n\|_{C([0, T]; L^2(\Omega))} \leq C \exp(T/2) T^{1/2} \left(o((\log n)^{-\gamma\beta/2}) + \begin{cases} \tau^{1/(3-p)} & p \in]1, 2], \\ \tau & p \geq 2, \end{cases} \right) \tag{58}$$

with probability at least $1 - o((\log n)^{-\gamma(1-\beta)})$.

Proof. Denote $\check{\xi}_n(x, t) = \check{v}_n(x, t) - \check{u}_n(x, t)$, $\bar{\xi}_n(x, t) = \bar{v}_n(x, t) - \bar{u}_n(x, t)$, $g_n = \mathbf{I}_n \mathbf{P}_n g$, $\widehat{K}_n = \mathbf{I}_n \widehat{\mathbf{K}}$ and $\Lambda_n = \mathbf{I}_n \Lambda$. We thus have from (54) and (55) that a.e.

$$\begin{aligned} \frac{\partial \check{\xi}_n(x, t)}{\partial t} &= -\left(\Delta_p^{\widehat{K}_n} (\bar{v}_n(x, t)) - \Delta_p^{\Lambda_n} (\bar{u}_n(x, t)) \right) \\ &= -\left(\Delta_p^{\widehat{K}_n} (\bar{v}_n(x, t)) - \Delta_p^{\Lambda_n} (\bar{v}_n(x, t)) \right) - \left(\Delta_p^{\Lambda_n} (\bar{v}_n(x, t)) - \Delta_p^{\Lambda_n} (\bar{u}_n(x, t)) \right). \end{aligned}$$

Multiplying both sides by $\check{\xi}_n(x, t)$, integrating and rearranging the terms, we get

$$\begin{aligned} \frac{1}{2} \frac{\partial}{\partial t} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2 &= - \int_{\Omega} (\Delta_p^{\Lambda_n} \bar{v}_n(x, t) - \Delta_p^{\Lambda_n} \bar{u}_n(x, t)) (\bar{v}_n(x, t) - \bar{u}_n(x, t)) \, dx \\ &\quad - \int_{\Omega} (\Delta_p^{\widehat{K}_n} \bar{v}_n(x, t) - \Delta_p^{\Lambda_n} \bar{v}_n(x, t)) \check{\xi}_n(x, t) \, dx \\ &\quad - \int_{\Omega} (\Delta_p^{\Lambda_n} \bar{v}_n(x, t) - \Delta_p^{\Lambda_n} \bar{u}_n(x, t)) ((\check{v}_n(x, t) - \bar{v}_n(x, t)) - (\check{u}_n(x, t) - \bar{u}_n(x, t))) \, dx. \end{aligned} \tag{59}$$

Under our condition on $n\rho_n$, Lemma 3.1 tells us that with probability 1,

$$\|\Lambda_n\|_{L^\infty,1(\Omega^2)} = \|\widehat{K}_n\|_{L^\infty,1(\Omega^2)} + o(1) \leq \|\mathbf{I}_n \mathbf{P}_n K\|_{L^\infty,1(\Omega^2)} + o(1) \leq \|K\|_{L^\infty,1(\Omega^2)} + o(1),$$

so in particular $\|\Lambda_n\|_{L^\infty,1(\Omega^2)}$ is uniformly bounded with probability 1. Λ_n is also positive and symmetric. Since $g \in L^q(\Omega)$ and $f \in L^1([0, T]; L^q(\Omega)) \cap \text{BV}([0, T]; L^2(\Omega))$, $q \in \{2, 2(p-1), +\infty\}$, the conclusions of Lemma 6.6 and Lemma 6.7 remain true which shows that with probability 1,

$$\sup_{t \in [0, T], n \in \mathbb{N}} \|\bar{u}_n(\cdot, t)\|_{L^q(\Omega)} < +\infty \quad \text{and} \quad \sup_{t \in [0, T], n \in \mathbb{N}} \|\check{u}_n(\cdot, t) - \bar{u}_n(\cdot, t)\|_{L^2(\Omega)} \leq C\tau.$$

The same claim holds for \check{v}_n and \bar{v}_n since $\|\widehat{K}_n\|_{L^\infty,1(\Omega^2)} \leq \|K\|_{L^\infty,1(\Omega^2)} < +\infty$ and \widehat{K}_n is positive and symmetric, *i.e.*, \widehat{K}_n obeys (H.1)–(H.3). Thus Proposition 4.1(iv) entails that the first term on the right-hand side of (59) is nonpositive with probability 1. Let us now bound the second term. Denote the random variables $\mathbf{Z}_i \stackrel{\text{def}}{=} \frac{1}{n} \sum_{j \in [n]} (\mathbf{\Lambda}_{ij} - \widehat{\mathbf{K}}_{ij}) \Psi(\mathbf{v}_j - \mathbf{v}_i)$. By Cauchy–Schwartz inequality, we have

$$\left| \int_{\Omega} (\Delta_p^{\Lambda_n} \bar{u}_n(x, t) - \Delta_p^{\widehat{K}_n} \bar{u}_n(x, t)) \check{\xi}_n(x, t) \, dx \right| \leq C \|\mathbf{I}_n \mathbf{Z}\|_{L^2(\Omega)} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}.$$

For the last term in (59), we argue as in the proof of Theorem 6.5 to show that, with probability 1,

$$\begin{aligned} &\left| \int_{\Omega} (\Delta_p^{\Lambda_n} \bar{u}_n(x, t) - \Delta_p^{\Lambda_n} \bar{v}_n(x, t)) ((\check{u}_n(x, t) - \bar{u}_n(x, t)) - (\check{v}_n(x, t) - \bar{v}_n(x, t))) \, dx \right| \\ &\leq C \begin{cases} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^{p-1} \tau + \tau^p & p \in]1, 2], \\ \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)} \tau + \tau^2 & p \geq 2. \end{cases} \end{aligned}$$

Collecting all these bounds, after using Young inequality, we have shown that (again with probability 1),

$$\frac{\partial}{\partial t} \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2 \leq \|\check{\xi}_n(\cdot, t)\|_{L^2(\Omega)}^2 + C \left(\|\mathbf{I}_n \mathbf{Z}\|_{L^2(\Omega)}^2 + \begin{cases} \tau^{2/(3-p)} + \tau^p & p \in]1, 2], \\ \tau^2 & p \geq 2. \end{cases} \right)$$

Using the Gronwall’s lemma and taking the square-root, we get for τ sufficiently small

$$\|\check{u}_n - \check{v}_n\|_{C([0, T]; L^2(\Omega))} \leq C \exp(T/2) T^{1/2} \left(\|\mathbf{I}_n \mathbf{Z}\|_{L^2(\Omega)} + \begin{cases} \tau^{1/(3-p)} & p \in]1, 2], \\ \tau & p \geq 2. \end{cases} \right) \tag{60}$$

It remains to bound the random variable $\|\mathbf{I}_n \mathbf{Z}\|_{L^2(\Omega)}$. For this purpose, we have by Markov inequality that for $\varepsilon > 0$

$$\mathbb{P}\left(\|\mathbf{I}_n \mathbf{Z}\|_{L^2(\Omega)} \geq \varepsilon\right) = \mathbb{P}\left(n^{-1} \sum_i \mathbf{Z}_i^2 \geq \varepsilon^2\right) \leq \varepsilon^{-2} n^{-1} \sum_i \mathbb{E}(\mathbf{Z}_i^2).$$

By independence of $(\mathbf{\Lambda}_{ij})_{j \in [n]}$, for each $i \in [n]$, we get

$$\begin{aligned} \mathbb{E}(\mathbf{Z}_i^2) &= (\rho_n n)^{-2} \sum_{j \in [n]} \mathbb{V}(\rho_n \mathbf{\Lambda}_{ij})(\Psi(\mathbf{v}_j - \mathbf{v}_i))^2 = (\rho_n n)^{-2} \sum_{j \in [n]} \rho_n \widehat{\mathbf{K}}_{ij} (1 - \rho_n \widehat{\mathbf{K}}_{ij})(\Psi(\mathbf{v}_j - \mathbf{v}_i))^2 \\ &\leq (\rho_n n^2)^{-1} \sum_{j \in [n]} \widehat{\mathbf{K}}_{ij} |\mathbf{v}_j - \mathbf{v}_i|^{2(p-1)}. \end{aligned}$$

In turn,

$$\begin{aligned} \mathbb{P}\left(\|\mathbf{I}_n \mathbf{Z}\|_{L^2(\Omega)} \geq \varepsilon\right) &\leq (\varepsilon^2 \rho_n n)^{-1} \frac{1}{n^2} \sum_{i,j \in [n]} \widehat{\mathbf{K}}_{ij} |\mathbf{v}_j - \mathbf{v}_i|^{2(p-1)} \\ &= (\varepsilon^2 \rho_n n)^{-1} \int_{\Omega^2} \widehat{K}_n(\mathbf{x}, \mathbf{y}) |\bar{v}_n(\mathbf{y}) - \bar{v}_n(\mathbf{x})|^{2(p-1)} d\mathbf{y} d\mathbf{x}. \end{aligned}$$

If the condition (a) holds, then by the symmetry of the kernel, Jensen inequality and Hölder inequality, one gets

$$\begin{aligned} \int_{\Omega^2} \widehat{K}_n(\mathbf{x}, \mathbf{y}) |\bar{v}_n(\mathbf{y}) - \bar{v}_n(\mathbf{x})|^{2(p-1)} d\mathbf{y} d\mathbf{x} &\leq 4 \int_{\Omega^2} \widehat{K}_n(\mathbf{x}, \mathbf{y}) |\bar{v}_n(\mathbf{x})|^{2(p-1)} d\mathbf{y} d\mathbf{x} \\ &\leq 4 \left\| \widehat{K}_n \right\|_{L^{\infty,1}(\Omega^2)} \int_{\Omega} |\bar{v}_n(\mathbf{x})|^{2(p-1)} d\mathbf{x} \\ &\leq 4 \left\| \widehat{K}_n \right\|_{L^{\infty,1}(\Omega^2)} \|\bar{v}_n\|_{L^2(\Omega)}^{2(p-1)}. \end{aligned}$$

Under the condition (b), by the symmetry of the kernel and Jensen inequality again, we have

$$\begin{aligned} \int_{\Omega^2} \widehat{K}_n(\mathbf{x}, \mathbf{y}) |\bar{v}_n(\mathbf{y}) - \bar{v}_n(\mathbf{x})|^{2(p-1)} d\mathbf{y} d\mathbf{x} &\leq 2^{2(p-1)} \int_{\Omega^2} \widehat{K}_n(\mathbf{x}, \mathbf{y}) |\bar{v}_n(\mathbf{x})|^{2(p-1)} d\mathbf{y} d\mathbf{x} \\ &\leq 2^{2(p-1)} \left\| \widehat{K}_n \right\|_{L^{\infty,1}(\Omega^2)} \|\bar{v}_n\|_{L^{2(p-1)}(\Omega)}^{2(p-1)}. \end{aligned}$$

Similarly, under condition (c), we have

$$\begin{aligned} \int_{\Omega^2} \widehat{K}_n(\mathbf{x}, \mathbf{y}) |\bar{v}_n(\mathbf{y}) - \bar{v}_n(\mathbf{x})|^{2(p-1)} d\mathbf{y} d\mathbf{x} &\leq 2^{2(p-1)} \|\bar{v}_n\|_{L^\infty(\Omega)}^{2(p-1)} \left\| \widehat{K}_n \right\|_{L^1(\Omega^2)} \\ &\leq 2^{2(p-1)} \left\| \widehat{K}_n \right\|_{L^{\infty,1}(\Omega^2)} \|\bar{v}_n\|_{L^\infty(\Omega)}^{2(p-1)}. \end{aligned}$$

Since $\left\| \widehat{K}_n \right\|_{L^{\infty,1}(\Omega^2)} \leq \|K\|_{L^{\infty,1}(\Omega^2)}$ (see (4) in Lem. 2.1), we have

$$\mathbb{P}\left(\|\mathbf{I}_n \mathbf{Z}\|_{L^2(\Omega)} \geq \varepsilon\right) \leq C (\varepsilon^2 \rho_n n)^{-1} \|K\|_{L^{\infty,1}(\Omega^2)},$$

where

$$C = \begin{cases} 4 \sup_n \|\bar{v}_n\|_{L^2(\Omega)}^{2(p-1)}, & \text{under (a),} \\ 2^{2(p-1)} \sup_n \|\bar{v}_n\|_{L^{2(p-1)}(\Omega)}^{2(p-1)}, & \text{under (b),} \\ 2^{2(p-1)} \sup_n \|\bar{v}_n\|_{L^\infty(\Omega)}^{2(p-1)}, & \text{under (c),} \end{cases}$$

and $C < +\infty$ thanks to Lemma 6.6. Taking $\varepsilon = \left(\frac{C \|K\|_{L^{\infty,1}(\Omega^2)}}{(\rho_n n)^\beta}\right)^{1/2}$, we get

$$\mathbb{P}\left(\|\mathbf{I}_n \mathbf{Z}\|_{L^2(\Omega)} \geq \varepsilon\right) \leq \frac{1}{(\rho_n n)^{1-\beta}}.$$

Plugging the latter into (60) completes the proof. □

Remark 7.1. Lemma 7.1 gives a deviation bound which holds with a controlled probability. One may ask whether almost sure convergence could be afforded. A naive and straightforward approach would be to invoke the Borel–Cantelli lemma as done in Remark 3.4(iv) from [31] for the case of graphons. But this argument does not apply to the more complex setting of L^q -graphons given that the probability of success in the statement Lemma 7.1 does not converge sufficiently fast. This is not even possible to make faster as ρ_n has to converge to 0. Thus, it is not clear at this stage whether this is even possible to achieve or not. We leave this to a future research.

Remark 7.2. One can also derive from Lemma 7.1 a bound in expectation. Let \mathcal{E}_n be the event on which (57) holds, and denote $\mathbf{1}_{\mathcal{E}_n} = 1$ if \mathcal{E}_n holds and 0 otherwise. We then have, for any $\beta \in]0, 1[$

$$\begin{aligned} \|\mathbb{E}(\check{u}_n) - \check{v}_n\|_{C([0,T];L^2(\Omega))} &\leq \mathbb{E}\left(\|\check{u}_n - \check{v}_n\|_{C([0,T];L^2(\Omega))}\right) \\ &= \mathbb{E}\left(\|\check{u}_n - \check{v}_n\|_{C([0,T];L^2(\Omega))}\mathbf{1}_{\mathcal{E}_n}\right) + \mathbb{E}\left(\|\check{u}_n - \check{v}_n\|_{C([0,T];L^2(\Omega))}\mathbf{1}_{\mathcal{E}_n^c}\right) \\ &\leq C \exp(T/2)T^{1/2} \left((\rho_n n)^{-\beta/2} + \begin{cases} \tau^{1/(3-p)} & p \in]1, 2], \\ \tau & p \geq 2. \end{cases} \right) + O((\rho_n n)^{-(1-\beta)}), \end{aligned}$$

where we have used the bound (57) and that \check{u}_n and \check{v}_n are bounded in $C([0, T]; L^2(\Omega))$ with probability one. We then conclude that $\|\mathbb{E}(\check{u}_n) - \check{v}_n\|_{C([0,T];L^2(\Omega))} \rightarrow 0$ as $n \rightarrow +\infty$.

We finally obtain the following error bound on fully discretized problems on sparse random graphs.

Theorem 7.1. *Suppose that $p \in]1, +\infty[$. Let u be a solution of (\mathcal{P}) with kernel K and data (f, g) , and $\{\mathbf{u}^k\}_{k \in [N]}$ is the sequence generated by $(\mathcal{P}_p^{\text{TDB,G}})$ with $\mathbf{K} = P_n K$, $\mathbf{g} = P_n g$, $\mathbf{f}^k = \tau_k^{-1} \int_{t_{k-1}}^{t_k} P_n f(\cdot, t) dt$. Assume that (f, g, K) satisfy the assumptions of Theorem 6.5, and that those of Lemma 7.1 also hold.*

(1) *For any $\beta \in]0, 1[$, with probability at least $1 - (\rho_n n)^{-(1-\beta)}$,*

$$\begin{aligned} \sup_{k \in [N], t \in]t_{k-1}, t_k]} \|\mathbf{I}_n \mathbf{u}^k - u(\cdot, t)\|_{L^2(\Omega)} &\leq \exp(T/2) \left(\|\mathbf{I}_n P_n g - g\|_{L^2(\Omega)} + \|\bar{f}_n - f\|_{L^1([0,T];L^2(\Omega))} + CT^{1/2}(\rho_n n)^{-\beta/2} \right. \\ &\quad \left. + CT^{1/2} \begin{cases} \tau^{1/(3-p)} + \|(K - \rho_n^{-1})_+\|_{L^{\infty,2}(\Omega^2)} + \|\mathbf{I}_n P_n K - K\|_{L^{\infty,2}(\Omega^2)} & \text{under (a)} \\ \tau^{p/(2p-1)} + \|(K - \rho_n^{-1})_+\|_{L^{\infty,2}(\Omega^2)} + \|\mathbf{I}_n P_n K - K\|_{L^{\infty,2}(\Omega^2)} & \text{under (b)} \\ \tau^{1/(3-p)} + \|(K - \rho_n^{-1})_+\|_{L^2(\Omega^2)} + \|\mathbf{I}_n P_n K - K\|_{L^2(\Omega^2)} & \text{under (c) when } p \in]1, 2] \\ \tau + \|(K - \rho_n^{-1})_+\|_{L^2(\Omega^2)} + \|\mathbf{I}_n P_n K - K\|_{L^2(\Omega^2)} & \text{under (c) when } p \geq 2. \end{cases} \right) \quad (61) \end{aligned}$$

for τ sufficiently small, where C is positive constant that depends only on p, g, f and K .

(2) *If, moreover, $g \in L^\infty(\Omega) \cap \text{Lip}(s, L^2(\Omega))$, $K \in \text{Lip}(s, L^2(\Omega^2))$, and $f \in L^1([0, T]; L^\infty(\Omega)) \cap \text{Lip}(s, L^2(\Omega \times [0, T]))$ then, for any $\delta \in]0, 1[$, with probability at least $1 - (\rho_n n)^{-(1-\beta)}$,*

$$\begin{aligned} \sup_{k \in [N], t \in]t_{k-1}, t_k]} \|\mathbf{I}_n \mathbf{u}^k - u(\cdot, t)\|_{L^2(\Omega)} &\leq C \exp(T/2) \left((1 + T^{1/2})n^{-s} + T^{1/2} \|(K - \rho_n^{-1})_+\|_{L^2(\Omega^2)} \right. \\ &\quad \left. + T^{1/2}(\rho_n n)^{-\beta/2} + T^{1/2} \begin{pmatrix} \tau^{\min(s, 1/(3-p))} & \text{when } p \in]1, 2] \\ \tau^s & \text{when } p \geq 2 \end{pmatrix} \right) \quad (62) \end{aligned}$$

for τ sufficiently small, where C is positive constant that depends only on p, g, f, K and s , and $\|(K - \rho_n^{-1})_+\|_{L^2(\Omega^2)} = o(1)$. The term τ^s in the dependence on τ disappears when f is time-independent.

Following the discussion in Remark 7.2, one can easily derive an error bound in expectation, *i.e.*, on $\sup_{k,t} \|\mathbb{I}_n \mathbb{E}(\mathbf{u}^k) - u(\cdot, t)\|_{L^2(\Omega)}$. We omit the details for the sake of brevity.

Proof. In view of (56), we shall use Theorem 6.5 to bound the second term, and a bound on the first term is provided by Lemma 7.1. Since $\mathbb{I}_n \widehat{\mathbf{K}}(x, y) \leq \mathbb{I}_n \mathbf{K}(x, y) = \mathbb{I}_n P_n K(x, y)$, the assumptions on \mathbf{K} transfer to $\widehat{\mathbf{K}}$, and the second term of (56) can then be bounded using (50), replacing $\mathbb{I}_n P_n K$ there by $\mathbb{I}_n \widehat{\mathbf{K}}$. Observing that

$$\begin{aligned} \|\mathbb{I}_n \widehat{\mathbf{K}} - K\|_{L^2(\Omega^2)} &= \|\min(\mathbb{I}_n P_n K, \rho_n^{-1}) - K\|_{L^2(\Omega^2)} \\ &\leq \|\min(\mathbb{I}_n P_n K, \rho_n^{-1}) - \mathbb{I}_n P_n K\|_{L^2(\Omega^2)} + \|\mathbb{I}_n P_n K - K\|_{L^2(\Omega^2)} \\ &= \|(\mathbb{I}_n P_n K - \rho_n^{-1})_+\|_{L^2(\Omega^2)} + \|\mathbb{I}_n P_n K - K\|_{L^2(\Omega^2)} \\ &\leq \|(K - \rho_n^{-1})_+\|_{L^2(\Omega^2)} + 2\|\mathbb{I}_n P_n K - K\|_{L^2(\Omega^2)}, \end{aligned}$$

and similarly for the L^∞ norm. The fact that $\|(K - \rho_n^{-1})_+\|_{L^2(\Omega^2)} = o(1)$ is because $\rho_n \rightarrow 0$ by the same argument as the end of the proof of Proposition 3.1. This completes the proof. \square

Acknowledgements. This work was supported by the Normandy Region grant MoNomads.

REFERENCES

- [1] F. Alvarez and J. Peypouquet, Asymptotic equivalence and Kobayashi-type estimates for nonautonomous monotone operators in Banach spaces. *Discrete Contin. Dyn. Syst.* **25** (2009) 1109–1128.
- [2] F. Andreu, J. Mazón, J. Rossi and J. Toledo, The Neumann problem for nonlocal nonlinear diffusion equations. *J. Evol. Equ.* **8** (2008) 189–215.
- [3] F. Andreu, J. Mazón, J. Rossi and J. Toledo, A nonlocal p -laplacian evolution equation with neumann boundary conditions. *J. Math. Pures Appl.* **90** (2008) 201–227.
- [4] F. Andreu-Vaillo, J.M. Mazón, J.D. Rossi and J.J. Toledo-Melero, Nonlocal Diffusion Problems. Vol. 165 of *Mathematical Surveys and Monographs*. American Mathematical Society (2010).
- [5] N. Ayi and N. Pouradier Duteil, Mean-field and graph limits for collective dynamics models with time-varying weights. *J. Differ. Equ.* **299** (2021) 65–110.
- [6] P.W. Bates and A. Chmaj, An integrodifferential model for phase transitions: stationary solutions in higher space dimensions. *J. Stat. Phys.* **95** (1999) 1119–1139.
- [7] P.W. Bates, P.C. Fife, X. Ren and X. Wang, Traveling waves in a convolution model for phase transitions. *Arch. Ration. Mech. Anal.* **138** (1997) 105–136.
- [8] H. Bauschke and P.L. Combettes, *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*. Springer (2011).
- [9] P. Bénilan, Solutions intégrales d'équations d'évolution dans un espace de banach. *C. R. Acad. Sci. Paris Ser. A-B* **274** (1972) A47–A50.
- [10] P. Bénilan and M. Crandall, Completely accretive operators, in *Semigroup Theory and Evolution Equations*. Vol. 135 of *Lecture Notes in Pure and Appl. Math.* Dekker, New York (1991) 41–75.
- [11] U. Biccari, D. Ko and E. Zuazua, Dynamics and control for multi-agent networked systems: a finite-difference approach. *Math. Models Methods Appl. Sci.* **29** (2019) 755–790.
- [12] B. Bollobás and O. Riordan, Metrics for sparse graphs, in *Surveys in Combinatorics 2009. London Mathematical Society Lecture Note Series*, edited by S. Huczynska, J.D. Mitchell and C.M.E. Roney-Dougal. Cambridge University Press, Cambridge (2009) 211–288.
- [13] C. Borgs, J.T. Chayes, H. Cohn and Y. Zhao, An L^p theory of sparse graph convergence II: LD convergence, quotients and right convergence. *Ann. Probab.* **46** (2018) 337–396.
- [14] C. Borgs, J.T. Chayes, H. Cohn and Y. Zhao, An L^p theory of sparse graph convergence I: limits, sparse random graph models, and power law distributions. *Trans. Amer. Math. Soc.* **372** (2019) 3019–3062.
- [15] H. Brézis, *Opérateurs Maximaux Monotones et semi-groupes de contractions dans les espaces de Hilbert*. North Holland, Amsterdam (1973).
- [16] T. Bühler and M. Hein, Spectral clustering based on the graph p -Laplacian, in *Proceedings of the 26th Annual International Conference on Machine Learning, ICML '09*. ACM, New York, NY, USA (2009) 81–88.
- [17] J. Byström, Sharp constants for some inequalities connected to the p -Laplace operator. *J. Inequalities Pure Appl. Math.* **6** (2005) 56.
- [18] J. Calder, The game theoretic p -laplacian and semi-supervised learning with few labels. *Nonlinearity* **32** (2018) 301.

- [19] C. Carrillo and P. Fife, Spatial effects in discrete generation population models. *J. Math. Biol.* **50** (2005) 161–188.
- [20] M.G. Crandall and T.M. Liggett, Generation of semigroups of nonlinear transformations on general banach spaces. *Amer. J. Math.* **93** (1971) 265–298.
- [21] M.G. Crandall and A. Pazy, Nonlinear evolution equations in banach spaces. *Isr. J. Math.* **11** (1972) 57–94.
- [22] R.A. DeVore and G.G. Lorentz, Constructive Approximation. Vol. 303 of *Grundlehren der mathematischen*. Springer-Verlag Berlin Heidelberg (1993).
- [23] A. El Alaoui, X. Cheng, A. Ramdas, M.J. Wainwright and M.I. Jordan, Asymptotic behavior of ℓ_p -based laplacian regularization in semi-supervised learning, in 29th Annual Conference on Learning Theory. PMLR (2016) 879–906.
- [24] A. Elmoataz, O. Lezoray, S. Bogleux and V.T. Ta, Unifying local and nonlocal processing with partial difference operators on weighted graphs, in International Workshop on Local and Non-Local Approximation in Image Processing. Switzerland (2008) 11–26.
- [25] A. Elmoataz, X. Desquesnes, Z. Lakhdari and O. L  zoray, Nonlocal infinity Laplacian equation on graphs with applications in image processing and machine learning. *Math. Comput. Simul.* **102** (2014) 153–163.
- [26] P. Erd  s and A. R  nyi, On the evolution of random graphs. *Publ. Math. Inst. Hung. Acad. Sci.* **5** (1960) 17–61.
- [27] P. Fife, Some nonclassical trends in parabolic and parabolic-like evolutions, in Trends in Nonlinear Analysis, edited by B. Fiedler. Springer-Verlag (2002).
- [28] P.C. Fife and X. Wang, A convolution model for interfacial motion: the generation and propagation of internal layers in higher space dimensions. *Adv. Differ. Equ.* **3** (1998) 85–110.
- [29] R. Glowinski and A. Marrocco, Sur l’approximation par   l  ments finis d’ordre un, et la r  solution, par p  nalisation-dualit  , d’une classe de probl  mes de Dirichlet non-lin  aires. *RAIRO: Anal. Num  r.* **2** (1975) 41–76.
- [30] Y. Hafiene, J. Fadili and A. Elmoataz, Nonlocal p -Laplacian evolution problems on graphs. *SIAM J. Numer. Anal.* **56** (2018) 1064–1090.
- [31] Y. Hafiene, M. Fadili, C. Chesneau and A.E. Moataz, Continuum limit of the nonlocal p -laplacian evolution problem on random in homogeneous graphs. *ESAIM: M2AN* **54** (2020) 565–589.
- [32] W. Hoeffding, The Strong Law of Large Numbers for u -Statistics. *Institute of Statistics Mimeograph Series 302*. North Carolina State University (1961).
- [33] D. Kaliuzhnyi-Verbovetskyi and G. Medvedev, The semilinear heat equation on sparse random graphs. *SIAM J. Math. Anal.* **49** (2017) 1333–1355.
- [34] T. Kato, Nonlinear semigroups and evolution equations. *J. Math. Soc. Jpn.* **19** (1967) 508–520.
- [35] B. Kawohl, Variations on the p -Laplacian. *Nonlinear Elliptic Part. Differ. Equ. Contemp. Math.* **540** (2011) 35–46.
- [36] Y. Kobayashi, Difference approximation of Cauchy problems for quasi-dissipative operators and generation of nonlinear semigroups. *J. Math. Soc. Jpn.* **27** (1975) 640–665.
- [37] K. Kobayashi, Y. Kobayashi and S. Oharu, Nonlinear evolution operators in Banach spaces. *Osaka J. Math.* **21** (1984) 281–310.
- [38] L. Lov  sz, Large Networks and Graph Limits. Vol. 60. American Mathematical Society (2012).
- [39] G.S. Medvedev, The nonlinear heat equation on dense graphs. *SIAM J. Math. Anal.* **46** (2014) 2743–2766.
- [40] G.S. Medvedev, The continuum limit of the kuramoto model on sparse random graphs. *Commun. Math. Sci.* **17** (2019) 883–898.
- [41] R.H. Nochetto and G. Savar  , Nonlinear evolution governed by accretive operators in banach spaces: error control and applications. *Math. Models Methods Appl. Sci.* **16** (2006) 439–477.
- [42] W. Rudin, Real and Complex Analysis, 3rd edition. McGraw-Hill (1987).
- [43] D. Slep  ev and M. Thorpe, Analysis of p -laplacian regularization in semisupervised learning. *SIAM J. Math. Anal.* **51** (2019) 2085–2120.
- [44] N.G. Trillos and D. Slep  ev, Continuum limit of total variation on point clouds. *Arch. Ration. Mech. Anal.* **220** (2016) 193–241.
- [45] X. Wang, Metastability and stability of patterns in a convolution model for phase transitions. *J. Differ. Equ.* **183** (2002) 434–461.

Please help to maintain this journal in open access!



This journal is currently published in open access under the Subscribe to Open model (S2O). We are thankful to our subscribers and supporters for making it possible to publish this journal in open access in the current year, free of charge for authors and readers.

Check with your library that it subscribes to the journal, or consider making a personal donation to the S2O programme by contacting subscribers@edpsciences.org.

More information, including a list of supporters and financial transparency reports, is available at <https://edpsciences.org/en/subscribe-to-open-s2o>.