

## A QUASI-INTERPOLATION OPERATOR YIELDING FULLY COMPUTABLE ERROR BOUNDS

THÉOPHILE CHAUMONT-FRELET<sup>1,\*</sup> AND MARTIN VOHRALÍK<sup>2,3</sup>

**Abstract.** We design a quasi-interpolation operator from the Sobolev space  $H_0^1(\Omega)$  to its finite-dimensional finite element subspace formed by piecewise polynomials on a simplicial mesh with a computable approximation constant. The operator (1) is defined on the entire  $H_0^1(\Omega)$ , no additional regularity is needed; (2) allows for an arbitrary polynomial degree; (3) works in any space dimension; (4) is defined locally, in vertex patches of mesh elements; (5) yields estimates optimal in the mesh size  $h$  for both the  $H^1$  seminorm and the  $L^2$  norm error; (6) yields estimates that bound the error in each computational mesh element by the best-approximation error in the element and in its vertex neighbors; (7) gives a computable constant for both the  $H^1$  seminorm and the  $L^2$  norm error; (8) leads to the equivalence of global-best and local-best errors; (9) possesses the projection property. Its construction follows the so-called potential reconstruction from *a posteriori* error analysis. Numerical experiments illustrate that our quasi-interpolation operator systematically gives the correct convergence rates in both the  $H^1$  seminorm and the  $L^2$  norm and its certified overestimation factor is rather sharp and stable in all tested situations.

**Mathematics Subject Classification.** 65N30, 65N15, 65D05.

Received July 16, 2025. Accepted October 17, 2025.

### 1. INTRODUCTION

Due to their ability to operate on general geometries, finite element methods based on unstructured meshes have become very popular to discretize boundary value problems over the past decades [10, 15]. The approximation properties of the finite element spaces are one of the key factors that eventually govern the accuracy of the resulting numerical schemes. Such approximation properties are often obtained thanks to (quasi-)interpolation operators, *i.e.*, mappings that associate to a given function  $u$  from the infinite-dimensional Sobolev space  $H_0^1(\Omega)$  an explicitly constructed approximation  $\mathcal{I}_h u$  in the finite-dimensional finite element space.

Given a polytopal domain  $\Omega \subset \mathbb{R}^d$ ,  $d \geq 1$ , triangulated by a simplicial mesh  $\mathcal{T}_h$ , this work focuses on conforming finite elements of polynomial degree  $p \geq 1$ ,  $\mathcal{P}_p(\mathcal{T}_h) \cap H_0^1(\Omega)$ , forming a finite-dimensional subspace of the Sobolev space  $H_0^1(\Omega)$ . We consider the Sobolev space  $H_0^1(\Omega)$  with the zero trace condition on the entire

---

*Keywords and phrases.* Finite element method, interpolation operator, stable projection, error estimate, guaranteed bound, minimal regularity.

<sup>1</sup> Inria, Univ. Lille, CNRS, UMR 8524 – Laboratoire Paul Painlevé, 40 Av. Halley, 59650 Villeneuve-d’Ascq, France.

<sup>2</sup> Inria, 48 rue Barrault, 75647 Paris, France.

<sup>3</sup> CERMICS, Ecole nationale des ponts et chaussées, IP Paris, 77455 Marne-la-Vallée, France.

\*Corresponding author: [theophile.chaumont@inria.fr](mailto:theophile.chaumont@inria.fr)

boundary  $\partial\Omega$  for the sake of simplicity of our exposition only. The construction and the results generalize easily to the case of zero trace condition only imposed on a part of  $\partial\Omega$  or not imposed at all. In this context, a natural way to associate to a function  $u \in H_0^1(\Omega)$  an interpolant  $\mathcal{I}_h u \in \mathcal{P}_p(\mathcal{T}_h) \cap H_0^1(\Omega)$  is by requiring that  $(\mathcal{I}_h u)(\mathbf{x}_\ell) = u(\mathbf{x}_\ell)$  for suitably chosen interpolation nodes  $\{\mathbf{x}_\ell\}_\ell$ . This process is known as Lagrange interpolation, and presents the advantage that it is fully local: For a mesh element  $K \in \mathcal{T}_h$ ,  $(\mathcal{I}_h u)|_K$  only depends on  $u|_K$  and consequently, the interpolation error  $(u - \mathcal{I}_h u)|_K$  will only depend on  $u|_K$ . Besides, fully-explicit error bounds expressed in terms of Sobolev semi-norms of  $u$  are available in the literature [1, 7, 21, 24–26, 28, 29].

The main drawback of Lagrange interpolation, however, is that it requires the nodal values of  $u$  to be well-defined, which is not the case (except in one space dimension,  $d = 1$ ) if  $u$  barely sits in  $H_0^1(\Omega)$ . When considering, *e.g.*, the Poisson problem in a convex domain with an  $L^2(\Omega)$  right-hand side, this is not an issue, since then the solution  $u$  belongs to  $H^2(\Omega)$ , enabling the use of Lagrange interpolation. However, the approach fails when considering more complex geometries and/or coefficients. Additionally, some applications, such as *a posteriori* error estimation [5, 31], preconditioning [37], or localized orthogonal decomposition methods [30], crucially hinge on interpolation operators defined for generic  $H_0^1(\Omega)$  target functions (*i.e.*, with the minimal variational regularity of the PDE under consideration).

There is therefore a need to define interpolation operators for non-smooth functions. The resulting operators are often called “quasi-interpolation” operators, since they do not, strictly-speaking, interpolate the target at any point. The seminal contribution in this direction is the work of Clément [11], where an operator defined over the whole  $L^1(\Omega)$  space is introduced with optimal  $h$ -approximation properties. The Clément operator does not preserve boundary conditions, and it is not a projection, *i.e.*,  $\mathcal{I}_h u_h \neq u_h$  for  $u_h$  already in the finite element space. These two properties have been incorporated later on in the key contribution of Scott and Zhang [38]. For these two operators, the degrees of freedom (dofs) of the interpolant  $\mathcal{I}_h u$  are fixed by suitably weighted mean values of the target function  $u$ . Another family of quasi-interpolation operators is due to Brenner [4] and Oswald [33]. There, a discontinuous approximation is first built by projecting element-wise  $u|_K$  onto a local polynomial space  $\mathcal{P}_p(K)$  of the mesh cell  $K \in \mathcal{T}_h$ . The dofs of this approximation that are shared by several elements are then suitably averaged to produce a conforming interpolant  $\mathcal{I}_h u \in \mathcal{P}_p(\mathcal{T}_h) \cap H_0^1(\Omega)$ . Since then, these constructions have been improved in several ways, and the design of quasi-interpolation operators is still an active field of research [14, 19, 20, 22, 23, 27, 36, 44].

Because the nodal evaluation of a generic target function  $u \in H_0^1(\Omega)$  is not possible, quasi-interpolation operators are not fully local. Indeed, the value of dofs needs to be obtained through some suitable averaging, meaning that the definition  $(\mathcal{I}_h u)|_K$  will not only depend on  $u|_K$ , but rather on  $u|_{\omega_K}$ , where  $\omega_K$  is the domain formed by other mesh elements surrounding  $K$ . Although this is not a huge drawback in practice, this fact makes the error analysis more complicated than for standard interpolation operators, since it is not possible to use arguments involving a single “reference element”. As a result, a shape-regularity requirement involving  $\omega_K$  rather than each element  $K$  individually often appears. In fact, the only fully-explicit approximation results we are aware of in this direction are [6, 40–42] but the scope of these results is limited, as they are specifically established with applications to *a posteriori* error estimation in mind.

Another important topic related to our work and connected to quasi-interpolation operators is the comparison between local-best and global-best approximations. Namely, we ask whether there exists a constant  $C > 0$  such that

$$\min_{u_h \in \mathcal{P}_p(\mathcal{T}_h) \cap H_0^1(\Omega)} \|\nabla(u - u_h)\|_\Omega \leq C \min_{v_h \in \mathcal{P}_p(\mathcal{T}_h)} \|\nabla_h(u - v_h)\|_\Omega. \quad (1.1)$$

In other words, we ask whether, up to a generic constant, the conforming finite element approximation is as good as the element-wise broken polynomial approximation of the target function. It is in fact clear that such a constant exists, because the left-hand side vanishes whenever the right-hand side does. However, the dependence of the constant  $C$  on key discretization parameters is a subtle issue. Following [2, 8, 39], recent results in this direction are given in [20, 27, 44]. Notice that once an inequality such as (1.1) is established, approximability estimates for the finite element space  $\mathcal{P}_p(\mathcal{T}_h) \cap H_0^1(\Omega)$  follow, since the right-hand side may be easily estimated in terms of (broken) Sobolev norms of  $u$  using element-wise Poincaré inequalities (see Sect. 4.5 below).

In this work, we propose a new quasi-interpolation operator  $\mathcal{J}_h^p : H_0^1(\Omega) \rightarrow \mathcal{P}_p(\mathcal{T}_h) \cap H_0^1(\Omega)$ . This operator can be computed locally by solving patch-wise finite element projection problems, works for any polynomial degree  $p \geq 1$ , preserves boundary conditions, is a projection, and, moreover, satisfies

$$\|\nabla(u - \mathcal{J}_h^p(u))\|_\Omega \leq C \min_{v_h \in \mathcal{P}_p(\mathcal{T}_h)} \|\nabla_h(u - v_h)\|_\Omega \quad \forall u \in H_0^1(\Omega) \quad (1.2)$$

for a constant  $C$  only depending on the shape-regularity parameter of the mesh  $\mathcal{T}_h$ , the space dimension  $d$ , and the polynomial degree  $p$ . In particular, our interpolation operator has the approximation power of discontinuous piecewise polynomials and leads to the local–global equivalence (1.1). We also establish localized versions of (1.2), as well as error estimates in the  $L^2(\Omega)$  norm (see Sect. 4 below). Our construction is inspired by the so-called potential reconstructions used in the context of a *a posteriori* error estimation for nonconforming and mixed finite element methods [16], as well as recent works of the authors on commuting quasi-interpolation operators [9, 18].

The main novelty of our work is that the constant  $C$  appearing in (1.2) is fully computable. Specifically, it is calculated through an algorithm that amounts to solving small, uncoupled, matrix eigenvalue problems that stem from patch-wise finite element spaces. We also establish localized versions of (1.2) and  $L^2(\Omega)$  error estimates with fully-computable constants. To the best of our knowledge, this work is the first to provide computable constants under minimal regularity, for arbitrary polynomial degree, and yielding the comparison of global-best and local-best errors. We emphasize that the proposed algorithm computes the generic constant  $C$  in (1.2) valid for all  $u \in H_0^1(\Omega)$  (without needing to know  $u$ ).

The remainder of this work is organized as follows. Section 2 gives the setting and recalls key preliminary results. In Section 3, we present elementwise Lagrange interpolation, the local-best approximation, and motivate our approach. The construction of our quasi-interpolation operator and statement of its key properties including (1.2) forms the content of Section 4. Finally, Section 5 collects the proofs and Section 6 reports on the actual behavior of our quasi-interpolation operator in several numerical experiments.

## 2. MESHES, SPACES, PATCHES, AND POINCARÉ’S INEQUALITY

### 2.1. Computational mesh

We consider an open, bounded, connected, Lipschitz polyhedral domain  $\Omega \subset \mathbb{R}^d$ ,  $d \geq 1$ , partitioned into a mesh  $\mathcal{T}_h$  that consists of (open) simplicial elements  $K$ . We assume that the mesh is matching in the sense that the intersection  $\overline{K_+} \cap \overline{K_-}$  of two elements  $K_\pm \in \mathcal{T}_h$  is either empty or a single common  $d'$ -dimensional subsimplex of  $\overline{K_+}$  and  $\overline{K_-}$ ,  $0 \leq d' \leq d - 1$  (a single vertex, edge, or face of  $K_-$  and  $K_+$  if  $d = 3$ ). This assumption is standard (see, *e.g.*, [10], Sect. 2.2 or [15], Def. 6.11); more general simplicial meshes with the so-called hanging nodes could be considered along the lines of the analysis carried out in [13, 17]. We denote by  $\mathcal{V}_h$  the set of vertices of  $\mathcal{T}_h$ .

For a simplex  $K \subset \mathbb{R}^d$ ,  $h_K$  denotes the diameter of  $K$  and  $\rho_K$  the diameter of the largest ball contained in  $\overline{K}$ . The shape-regularity parameter of  $K$  is then defined by

$$\kappa_K := \frac{h_K}{\rho_K}.$$

If  $\mathcal{T} \subset \mathcal{T}_h$  is a submesh, the shape-regularity parameter of  $\mathcal{T}$  is  $\kappa_{\mathcal{T}} := \max_{K \in \mathcal{T}} \kappa_K$ . Notice that  $\kappa_K \geq 1$  and that it is possible to design strongly graded meshes  $\mathcal{T}_h$  while maintaining  $\kappa_{\mathcal{T}_h}$  bounded, as long as the elements remain isotropic.

If  $K \subset \mathbb{R}^d$  is a simplex, we employ the notation  $\mathcal{F}_K$  for its  $(d - 1)$ -dimensional faces and  $\mathcal{V}_K$  for its vertices. Then, if  $\mathbf{a} \in \mathcal{V}_K$  is a vertex of  $K$ , the notation  $\tau_K^\mathbf{a}$  stands for the distance between  $\mathbf{a}$  and the (hyper)plane generated by the face of  $K$  opposite to  $\mathbf{a}$ . Notice that  $\rho_K \leq \tau_K^\mathbf{a} \leq h_K$ , so that in particular  $h_K/\tau_K^\mathbf{a} \leq \kappa_K$ .

### 2.2. Sobolev spaces

Throughout this manuscript, if  $\omega \subset \Omega$  is an open, bounded, connected, Lipschitz subset of  $\Omega$ , then  $L^2(\omega)$  is the Lebesgue space of scalar-valued square-integrable functions and  $\mathbf{L}^2(\omega) := [L^2(\omega)]^d$  is the space of vector-valued square-integrable functions. We denote by  $(\cdot, \cdot)_\omega$  and  $\|\cdot\|_\omega$  the inner products and norms of both spaces.

We employ the notation  $L^\infty(\omega)$  and  $\mathbf{L}^\infty(\omega)$  for essentially bounded scalar- and vector-valued functions, and denote by  $\|\cdot\|_{L^\infty(\omega)}$  and  $\|\cdot\|_{\mathbf{L}^\infty(\omega)}$  their usual norms.

The Sobolev spaces

$$H^1(\omega) := \{v \in L^2(\omega) \mid \nabla v \in \mathbf{L}^2(\omega)\}, \quad \mathbf{H}(\operatorname{div}, \omega) := \{\mathbf{w} \in \mathbf{L}^2(\omega) \mid \nabla \cdot \mathbf{w} \in L^2(\omega)\},$$

where  $\nabla$  and  $\nabla \cdot$  respectively denote the gradient and divergence operators defined in the sense of distributions, will be useful. We will also employ the following notation:

$$H_0^1(\omega) := \{v \in H^1(\omega) \mid v = 0 \text{ on } \partial\omega\}, \quad \mathbf{H}(\operatorname{div}^0, \omega) := \{\mathbf{v} \in \mathbf{H}(\operatorname{div}, \omega) \mid \nabla \cdot \mathbf{v} = 0\},$$

where the boundary value is understood in the trace sense.

For scalar-valued functions, we will also use higher-order Sobolev spaces. Namely for an integer  $s \in \mathbb{N}$ ,  $H^s(\omega)$  stands for the space of functions  $v \in L^2(\omega)$  such that

$$\partial_\alpha v \in L^2(\omega)$$

for all  $\alpha \in \mathbb{N}^d$  with  $|\alpha|_1 \leq s$ , whereby  $|\cdot|_1$  denotes the  $\ell_1$  norm on  $\mathbb{N}^d$ . We equip  $H^s(\omega)$  with the seminorm

$$|v|_{H^s(\omega)}^2 := \sum_{\substack{\alpha \in \mathbb{N}^d \\ |\alpha|_1 = s}} \|\partial_\alpha v\|_\omega^2 \quad \forall v \in H^s(\omega). \quad (2.1)$$

The broken Sobolev spaces

$$H^s(\mathcal{T}_h) = \{v \in L^2(\Omega) \mid v|_K \in H^s(K) \quad \forall K \in \mathcal{T}_h\}$$

will also be useful. The broken gradient is in particular defined element-wise as

$$(\nabla_h v)|_K := \nabla(v|_K) \quad \forall K \in \mathcal{T}_h$$

for all  $v \in H^1(\mathcal{T}_h)$ . It maps  $H^1(\mathcal{T}_h)$  into  $\mathbf{L}^2(\Omega)$ .

### 2.3. Finite element spaces

If  $K \subset \mathbb{R}^d$  is a simplex and  $q \geq 0$ , we denote by  $\mathcal{P}_q(K)$  the set of polynomial functions of total degree at most  $q$  on  $K$  and  $\mathcal{P}_q(K) := [\mathcal{P}_q(K)]^d$ . The set of Raviart–Thomas polynomials on  $K$  is then  $\mathcal{RT}_q(K) := \mathbf{x}\mathcal{P}_q(K) + \mathcal{P}_q(K)$ , see [32, 35].

If  $\mathcal{T}$  is a set of simplices with the corresponding domain  $\omega \subset \Omega$ , then  $\mathcal{P}_q(\mathcal{T})$  collects the functions  $v : \omega \rightarrow \mathbb{R}$  such that  $v|_K \in \mathcal{P}_q(K)$  for all  $K \in \mathcal{T}$ . The elementwise (broken) space  $\mathcal{RT}_q(\mathcal{T})$  is defined analogously.

Below, we fix a polynomial degree  $p \geq 1$ .

### 2.4. Hat functions

Throughout this work, the hat functions  $\{\psi_{\mathbf{a}}\}_{\mathbf{a} \in \mathcal{V}_h}$  will play an important role. For each mesh vertex  $\mathbf{a} \in \mathcal{V}_h$ ,  $\psi_{\mathbf{a}}$  is the only element of  $\mathcal{P}_1(\mathcal{T}_h) \cap H^1(\Omega)$  such that  $\psi_{\mathbf{a}}(\mathbf{b}) = \delta_{\mathbf{a}, \mathbf{b}}$  for all  $\mathbf{b} \in \mathcal{V}_h$ , where  $\delta$  is the Kronecker symbol. If  $K \in \mathcal{T}_h$  and  $\mathbf{a} \in \mathcal{V}_K$ , then

$$\|\psi_{\mathbf{a}}\|_{L^\infty(K)} = 1 \quad \|\nabla \psi_{\mathbf{a}}\|_{\mathbf{L}^\infty(K)} = \frac{1}{\tau_K^{\mathbf{a}}}. \quad (2.2)$$

We also denote by  $\omega_{\mathbf{a}}$  the open domain corresponding to the support of  $\psi_{\mathbf{a}}$ . Crucially, the hat functions form a partition of unity as

$$\sum_{\mathbf{a} \in \mathcal{V}_h} \psi_{\mathbf{a}} = 1. \quad (2.3)$$

## 2.5. Vertex patches

If  $\mathbf{a} \in \mathcal{V}_h$  we define the patch

$$\mathcal{T}_{\mathbf{a}} = \{K \in \mathcal{T}_h \mid \mathbf{a} \in \mathcal{V}_K\},$$

of elements having  $\mathbf{a}$  as a vertex. The elements  $K \in \mathcal{T}_{\mathbf{a}}$  correspond to the support  $\omega_{\mathbf{a}}$  of the hat function  $\psi_{\mathbf{a}}$ . For a vertex patch  $\mathcal{T}_{\mathbf{a}}$ , we define

$$h_{\mathbf{a}} := \max_{K \in \mathcal{T}_h \mid \mathbf{a} \in \mathcal{V}_K} h_K. \quad (2.4)$$

## 2.6. Poincaré's inequality

Since simplices are convex, for all  $K \in \mathcal{T}_h$  and for all  $v \in H^1(K)$  such that  $(v, 1)_K = 0$ , we have

$$\|v\|_K \leq \frac{h_K}{\pi} \|\nabla v\|_K, \quad (2.5)$$

see, *e.g.*, [3, 34].

## 3. ELEMENTWISE LAGRANGE INTERPOLATION, LOCAL-BEST APPROXIMATION, AND MAIN IDEAS

In this section, we present the elementwise Lagrange interpolation operator and discuss local-best approximation by  $H^1$  seminorm elementwise orthogonal projection. We then motivate the construction of our quasi-interpolation operator.

### 3.1. Elementwise Lagrange interpolation operator

We start by recalling the standard Lagrange interpolation operator on each mesh element  $K \in \mathcal{T}_h$ . We will only apply it to functions that are polynomials of degree  $p+1$  on  $K$ , so that we have  $\mathcal{I}_K^p : \mathcal{P}_{p+1}(K) \rightarrow \mathcal{P}_p(K)$ ,

$$(\mathcal{I}_K^p v_h)(\mathbf{x}_\ell) = v_h(\mathbf{x}_\ell) \quad (3.1a)$$

for all Lagrange interpolation nodes  $\{\mathbf{x}_\ell\}_\ell$  on  $K$  see, *e.g.*, Section 7.4 of [15]. Then, the elementwise Lagrange interpolation operator  $\mathcal{I}_h^p : \mathcal{P}_{p+1}(\mathcal{T}_h) \rightarrow \mathcal{P}_p(\mathcal{T}_h)$  is defined by

$$(\mathcal{I}_h^p v)|_K := \mathcal{I}_K^p(v|_K) \quad \forall K \in \mathcal{T}_h, \quad (3.1b)$$

for  $v \in \mathcal{P}_{p+1}(\mathcal{T}_h)$ . Let us point out that if  $v \in \mathcal{P}_{p+1}(\mathcal{T}_h) \cap H_0^1(\Omega)$ , then  $\mathcal{I}_h^p v \in \mathcal{P}_p(\mathcal{T}_h) \cap H_0^1(\Omega)$  and that  $\mathcal{I}_h^p$  is a projection.

**Remark 3.1** (Other interpolation operators). We focus on the Lagrange interpolation operator  $\mathcal{I}_K^p$  on each mesh element  $K$  for the sake of simplicity. In fact, any affine-equivalent interpolation operator  $\mathcal{I}_K^p : \mathcal{P}_{p+1}(K) \rightarrow \mathcal{P}_p(K)$  with the property that  $\mathcal{I}_h^p$  maps  $\mathcal{P}_{p+1}(\mathcal{T}_h) \cap H_0^1(\Omega)$  into  $\mathcal{P}_p(\mathcal{T}_h) \cap H_0^1(\Omega)$  can be used.

### 3.2. Local-best approximation

Another key ingredient of our quasi-interpolation operator is the so-called ‘‘local-best’’ approximation. For  $K \in \mathcal{T}_h$  and  $v \in H^1(K)$ , we denote by  $\pi_K^p v \in \mathcal{P}_p(K)$  the  $H^1$ -orthogonal projection of  $v$ . Specifically, it is defined as the only element of  $\mathcal{P}_p(K)$  such that  $(\pi_K^p v, 1)_K = (v, 1)_K$  and

$$(\nabla(\pi_K^p v), \nabla q_h)_K = (\nabla v, \nabla q_h)_K \quad \forall q_h \in \mathcal{P}_p(K). \quad (3.2a)$$

The ‘‘local-best’’ approximation of  $v \in H^1(\mathcal{T}_h)$ ,  $\pi_h^p v \in \mathcal{P}_p(\mathcal{T}_h)$ , is then defined by setting

$$(\pi_h^p v)|_K := \pi_K^p(v|_K) \quad \forall K \in \mathcal{T}_h. \quad (3.2b)$$

Notice that then

$$\|\nabla_h(v - \pi_h^p v)\|_K = \min_{v_h \in \mathcal{P}_p(K)} \|\nabla_h(v - v_h)\|_K$$

and

$$\|\nabla_h(v - \pi_h^p v)\|_\Omega = \min_{v_h \in \mathcal{P}_p(\mathcal{T}_h)} \|\nabla_h(v - v_h)\|_\Omega.$$

We stress that, in general,  $\pi_h^p v$  is discontinuous, *i.e.*,  $\pi_h^p v \notin H_0^1(\Omega)$  even if  $v \in H_0^1(\Omega)$ .

### 3.3. Main idea of the construction of the quasi-interpolation operator

Here, we explain the spirit of the definition of our operator. Given  $u \in H_0^1(\Omega)$ , we start by considering the local-best approximation  $\pi_h^p u \in \mathcal{P}_p(\mathcal{T}_h)$  of  $u$  given by (3.2). This function is locally defined and exhibits the best-possible approximation properties, but it is unfortunately nonconforming since  $\pi_h^p u \notin H_0^1(\Omega)$  in general.

The next stage is then to somehow “locally project”  $\pi_h^p u$  to the conforming Lagrange finite element space  $\mathcal{P}_p(\mathcal{T}_h) \cap H_0^1(\Omega)$ . The optimal projection

$$\arg \min_{v_h \in \mathcal{P}_p(\mathcal{T}_h) \cap H_0^1(\Omega)} \|\nabla_h(\pi_h^p u - v_h)\|_\Omega = \arg \min_{v_h \in \mathcal{P}_p(\mathcal{T}_h) \cap H_0^1(\Omega)} \|\nabla(u - v_h)\|_\Omega^2 \quad (3.3)$$

is not satisfactory for a quasi-interpolation operator since it is global: it requires a global system solve and does not lead to local approximation properties, since the minimizer from (3.3) restricted to  $K \in \mathcal{T}_h$  depends on the values of  $u$  in the whole domain  $\Omega$  in general. Note that the two minimizers from (3.3) indeed coincide, since the Euler–Lagrange optimality conditions use respectively

$$(\nabla_h \pi_h^p u, \nabla v_h) \quad \text{and} \quad (\nabla u, \nabla v_h), \quad v_h \in \mathcal{P}_p(\mathcal{T}_h) \cap H_0^1(\Omega),$$

which coincide in view of (3.2). Our key idea is to first set up  $\pi_h^p u$  and then project it locally in vertex patches.

We employ the partition of unity by the hat functions (2.3) and try to approximate  $\psi_\alpha \pi_h^p u$  into a local contribution  $s_h^\alpha \in \mathcal{P}_p(\mathcal{T}_\alpha) \cap H_0^1(\omega_\alpha)$ . This choice makes sense, since (a)  $\psi_\alpha \pi_h^p u$  is expected to be close to  $\psi_\alpha u$  which sits in  $H_0^1(\omega_\alpha)$  and (b) the boundary conditions ensure that the object resulting from the summation of the local contributions is globally conforming.

At this point, it is therefore tempting to define

$$s_h^\alpha := \arg \min_{v_h \in \mathcal{P}_p(\mathcal{T}_\alpha) \cap H_0^1(\omega_\alpha)} \|\nabla_h(\psi_\alpha \pi_h^p u - v_h)\|_{\omega_\alpha}, \quad \mathcal{J}_h^p(u) := \sum_{\alpha \in \mathcal{V}_h} s_h^\alpha. \quad (3.4)$$

The construction in (3.4) faithfully conveys the ideas behind the construction of our projector. However, as we shall see later, (3.4) would fail in the projection property and would only give  $p-1$  approximation order because we are trying to represent a polynomial of degree  $p+1$  with a polynomial of degree  $p$ . Indeed, the presence of the hat function  $\psi_\alpha$  increases the polynomial degree by 1; to bring it back, we will use the elementwise Lagrange interpolation operator (3.1).

### 3.4. Main idea for the computable approximation constant

The local problem (3.4) is an approximation of the piecewise polynomial but discontinuous (nonconforming) datum  $\psi_\alpha \pi_h^p u$  by the conforming finite element method. Our key idea to obtain a computable error bound is to employ duality, *i.e.*, define a local counterpart of problem (3.4) *via* Raviart–Thomas finite elements. This will allow to measure the nonconformity of the local-best approximation  $\pi_h^p u$  vertex patch by vertex patch *via* a local discrete eigenvalue problem and bring us to the precision of the  $H^1$ -orthogonal projection of (3.2). The projection (3.2) is defined separately on each mesh element and its approximation bound comes with a fully explicit, computable, constant, see (4.8) below. Combining these ingredients then gives estimates optimal in the mesh size  $h$  for both the  $H^1$  seminorm and the  $L^2$  norm with a fully computable constant (we do not perform  $p$ -explicit/ $p$ -optimal analysis).

## 4. MAIN RESULTS

In this section, we construct our quasi-interpolation operator and present its fully computable error bounds. The proofs are postponed until Section 5.

### 4.1. Local potential reconstruction

Let a mesh vertex  $\mathbf{a} \in \mathcal{V}_h$  be given. We develop further (3.4). As it will become apparent below, we need here to start from an arbitrary (discontinuous) piecewise polynomial  $u_h \in \mathcal{P}_p(\mathcal{T}_\mathbf{a})$ . We define  $s_h^\mathbf{a}(u_h)$  as the continuous piecewise polynomial with vanishing trace on the boundary  $\partial\omega_\mathbf{a}$  of the vertex patch subdomain  $\omega_\mathbf{a}$ ,  $s_h^\mathbf{a}(u_h) \in \mathcal{P}_p(\mathcal{T}_\mathbf{a}) \cap H_0^1(\omega_\mathbf{a})$ , by

$$s_h^\mathbf{a}(u_h) := \arg \min_{v_h \in \mathcal{P}_p(\mathcal{T}_\mathbf{a}) \cap H_0^1(\omega_\mathbf{a})} \|\nabla_h(\mathcal{I}_h^p(\psi_\mathbf{a}u_h) - v_h)\|_{\omega_\mathbf{a}}. \quad (4.1a)$$

Equivalently,  $s_h^\mathbf{a}$  is given by the Euler–Lagrange conditions of (4.1a), reading as: find  $s_h^\mathbf{a}(u_h) \in \mathcal{P}_p(\mathcal{T}_\mathbf{a}) \cap H_0^1(\omega_\mathbf{a})$  such that

$$(\nabla s_h^\mathbf{a}(u_h), \nabla v_h)_{\omega_\mathbf{a}} = (\nabla_h(\mathcal{I}_h^p(\psi_\mathbf{a}u_h)), \nabla v_h)_{\omega_\mathbf{a}} \quad \forall v_h \in \mathcal{P}_p(\mathcal{T}_\mathbf{a}) \cap H_0^1(\omega_\mathbf{a}). \quad (4.1b)$$

Since  $\|\nabla \cdot\|_{\omega_\mathbf{a}}$  is a norm on  $H_0^1(\omega_\mathbf{a})$ ,  $s_h^\mathbf{a}(u_h)$  is indeed uniquely defined. We observe that the Lagrange interpolation operator  $\mathcal{I}_h^p$  of (3.1) is employed element-wise in order to reduce the polynomial degree  $p+1$  of  $\psi_\mathbf{a}u_h$  back to  $p$ . This is legal, since  $\psi_\mathbf{a}u_h$  is a polynomial (it would have failed if immediately applied to  $\psi_\mathbf{a}u$ ). We implicitly extend  $s_h^\mathbf{a}(u_h)$  by zero outside of  $\omega_\mathbf{a}$ , so that  $s_h^\mathbf{a}$  is an operator from  $\mathcal{P}_p(\mathcal{T}_\mathbf{a})$  to  $\mathcal{P}_p(\mathcal{T}_h) \cap H_0^1(\Omega)$ . The operator  $s_h^\mathbf{a}$  is a crucial ingredient of our new quasi-interpolation operator in Section 4.5 below.

### 4.2. Local flux reconstruction

For each vertex  $\mathbf{a} \in \mathcal{V}_h$ , consider the local finite element space

$$\mathbf{W}_h^\mathbf{a} := \mathcal{RT}_p(\mathcal{T}_\mathbf{a}) \cap \mathbf{H}(\operatorname{div}^0, \omega_\mathbf{a}) \quad (4.2)$$

of divergence-free  $\mathbf{H}(\operatorname{div}, \omega_\mathbf{a})$ -conforming Raviart–Thomas piecewise polynomials defined over the vertex patch subdomain  $\omega_\mathbf{a}$ . For an arbitrary piecewise polynomial  $u_h \in \mathcal{P}_p(\mathcal{T}_\mathbf{a})$ , we define  $\mathbf{r}_h^\mathbf{a}(u_h)$  as the only element of  $\mathbf{W}_h^\mathbf{a}$  such that

$$\mathbf{r}_h^\mathbf{a}(u_h) := \arg \min_{\mathbf{w}_h \in \mathbf{W}_h^\mathbf{a}} \|\nabla_h(\psi_\mathbf{a}u_h) - \mathbf{w}_h\|_{\omega_\mathbf{a}}, \quad (4.3a)$$

or, equivalently,  $\mathbf{r}_h^\mathbf{a}(u_h) \in \mathbf{W}_h^\mathbf{a}$  such that

$$(\mathbf{r}_h^\mathbf{a}(u_h), \mathbf{w}_h)_{\omega_\mathbf{a}} = (\nabla_h(\psi_\mathbf{a}u_h), \mathbf{w}_h)_{\omega_\mathbf{a}} \quad \forall \mathbf{w}_h \in \mathbf{W}_h^\mathbf{a}. \quad (4.3b)$$

Notice that we then have

$$\|\mathbf{r}_h^\mathbf{a}(u_h)\|_{\omega_\mathbf{a}} = \max_{\substack{\mathbf{w}_h \in \mathbf{W}_h^\mathbf{a} \\ \|\mathbf{w}_h\|_{\omega_\mathbf{a}}=1}} (\nabla_h(\psi_\mathbf{a}u_h), \mathbf{w}_h)_{\omega_\mathbf{a}}. \quad (4.4)$$

From a practical standpoint, given  $u_h \in \mathcal{P}_p(\mathcal{T}_\mathbf{a})$ ,  $\mathbf{r}_h^\mathbf{a}(u_h)$  can be computed as  $\mathbf{r}_h^\mathbf{a}(u_h) = \mathbf{r}_h$ , where  $(\mathbf{r}_h, q_h) \in \mathbf{X}_h^\mathbf{a} \times Y_h^\mathbf{a}$  is the unique pair such that

$$\begin{cases} (\mathbf{r}_h, \mathbf{v}_h)_{\omega_\mathbf{a}} + (q_h, \nabla \cdot \mathbf{v}_h)_{\omega_\mathbf{a}} = (\nabla_h(\psi_\mathbf{a}u_h), \mathbf{v}_h)_{\omega_\mathbf{a}}, \\ (\nabla \cdot \mathbf{r}_h, p_h)_{\omega_\mathbf{a}} = 0, \end{cases} \quad (4.5)$$

for all  $(\mathbf{v}_h, p_h) \in \mathbf{X}_h^\mathbf{a} \times Y_h^\mathbf{a}$ , with

$$\mathbf{X}_h^\mathbf{a} := \mathcal{RT}_p(\mathcal{T}_\mathbf{a}) \cap \mathbf{H}(\operatorname{div}, \omega_\mathbf{a}), \quad Y_h^\mathbf{a} := \mathcal{P}_p(\mathcal{T}_\mathbf{a}).$$

Indeed, although (4.3) and (4.5) are equivalent, in contrast to  $\mathbf{W}_h^\mathbf{a}$ , both  $\mathbf{X}_h^\mathbf{a}$  and  $Y_h^\mathbf{a}$  can be easily equipped with an explicit basis. The operator  $\mathbf{r}_h^\mathbf{a}$  is employed below to obtain the computable factors  $\lambda_\mathbf{a}$ .

### 4.3. A local primal–dual quotient and its computation by a local matrix eigenvalue problem

Consider a vertex  $\mathbf{a} \in \mathcal{V}_h$ . For  $u_h \in \mathcal{P}_p(\mathcal{T}_{\mathbf{a}})$ , let  $s_h^{\mathbf{a}}(u_h)$  be defined by (4.1) and  $\mathbf{r}_h^{\mathbf{a}}(u_h)$  by (4.3). We define the primal–dual quotient

$$\lambda_{\mathbf{a}} := \max_{u_h \in \mathcal{P}_p(\mathcal{T}_{\mathbf{a}})} \frac{\|\nabla_h(\mathcal{I}_h^p(\psi_{\mathbf{a}}u_h) - s_h^{\mathbf{a}}(u_h))\|_{\omega_{\mathbf{a}}}}{\|\mathbf{r}^{\mathbf{a}}(u_h)\|_{\omega_{\mathbf{a}}}}. \quad (4.6)$$

Theorem 5.1 below warrants that  $\lambda_{\mathbf{a}}$  is finite and only depends on the polynomial degree  $p$ , the space dimension  $d$ , and the shape-regularity parameter  $\kappa_{\mathcal{T}_{\mathbf{a}}}$  of the patch.

The number  $\lambda_{\mathbf{a}}$  can be computed through a small-size matrix eigenvalue problem. Specifically, given a basis of  $\mathcal{P}_p(\mathcal{T}_{\mathbf{a}})$ , since the operator

$$u_h \in \mathcal{P}_p(\mathcal{T}_{\mathbf{a}}) \rightarrow \mathcal{P}_p(\mathcal{T}_{\mathbf{a}}) \ni \mathcal{I}_h^p(\psi_{\mathbf{a}}u_h) - s_h^{\mathbf{a}}(u_h)$$

is linear, its action may be encoded as a matrix  $D \in \mathbb{R}^{|\mathcal{P}_p(\mathcal{T}_{\mathbf{a}})| \times |\mathcal{P}_p(\mathcal{T}_{\mathbf{a}})|}$ . Similarly, the action of the operator  $\mathbf{r}_h^{\mathbf{a}} : \mathcal{P}_p(\mathcal{T}_{\mathbf{a}}) \rightarrow \mathbf{W}_h^{\mathbf{a}} \subset \mathbf{X}_h^{\mathbf{a}}$  can be easily encoded as a simple matrix  $R \in \mathbb{R}^{|\mathcal{P}_p(\mathcal{T}_{\mathbf{a}})| \times |\mathbf{X}_h^{\mathbf{a}}|}$  after inverting the matrix representation of the left-hand side of (4.5). Then, if  $M$  and  $K$  respectively denote the mass and stiffness matrices of  $\mathbf{X}_h^{\mathbf{a}} = \mathcal{RT}_p(\mathcal{T}_{\mathbf{a}}) \cap \mathbf{H}(\text{div}, \omega_{\mathbf{a}})$  and  $\mathcal{P}_p(\mathcal{T}_{\mathbf{a}})$ ,  $\lambda_{\mathbf{a}}$  can be obtained as the largest eigenvalue of the problem: Find  $\mu \in \mathbb{R}$  and  $\mathbf{v} \in \mathbb{R}^{|\mathcal{P}_p(\mathcal{T}_{\mathbf{a}})|}$  such that

$$D^T K D \mathbf{v} = \mu R^T M R \mathbf{v}.$$

### 4.4. A local mesh shape-regularity characterization

Recall the notation of Section 2.1. For a vertex  $\mathbf{a} \in \mathcal{V}_h$ , define the constant

$$\rho_{\mathbf{a}} := 1 + \frac{1}{\pi} \max_{K \in \mathcal{T}_{\mathbf{a}}} \frac{h_K}{\tau_K^{\mathbf{a}}}. \quad (4.7)$$

It is fully computable and only depends on the shape-regularity parameter  $\kappa_{\mathcal{T}_{\mathbf{a}}}$  of  $\mathcal{T}_{\mathbf{a}}$ .

### 4.5. Local-best approximation error estimate

For the local-best approximation of (3.2), we observe that, applying the Poincaré inequality stated in (2.5) repeatedly, see Corollary 12.13 of [15] or Theorem 1.8.1 of [43] for details,

$$\|\nabla_h(v - \pi_h^p v)\|_K^2 \leq (s+1)! \left(\frac{h_K}{\pi}\right)^{2s} |v|_{H^{1+s}(K)}^2 \quad \forall v \in H^{1+s}(K), \forall K \in \mathcal{T}_h \quad (4.8)$$

for all integer  $0 \leq s \leq p$ . Crucially, the constant appearing in (4.8) is fully explicit (computable). We remark, though, that (4.8) is not explicit/optimal in the polynomial degree  $p$ ; estimates optimal with respect to  $p$  are available in the literature with, however, an unknown (not explicit) constant.

**Remark 4.1** (Uniform-in- $s$  constant for a redefined Sobolev seminorm). If one redefines the seminorm  $|v|_{H^s(\omega)}$  from (2.1) to repetitively contain the indices, starting with  $|v|_{H^2(\omega)} = \sum_{i=1}^d \sum_{j=1}^d \|\partial_{\mathbf{x}_j} \partial_{\mathbf{x}_i} v\|_K^2$  in place of  $|v|_{H^2(\omega)} = \sum_{i=1}^d \sum_{j=i}^d \|\partial_{\mathbf{x}_j} \partial_{\mathbf{x}_i} v\|_K^2$ , then the factor  $(s+1)!$  from (4.8) can be removed.

### 4.6. The quasi-interpolation operator $\mathcal{J}_h^p$

Let  $u \in H_0^1(\Omega)$ . To define our quasi-interpolation operator, we employ (4.1a) with the argument  $u_h$  being the local-best approximation  $\pi_h^p u$  of the target function  $u$  given by (3.2), restricted to the patch subdomain  $\omega_{\mathbf{a}}$ :

$$\mathcal{J}_h^p u := \sum_{\mathbf{a} \in \mathcal{V}_h} s_h^{\mathbf{a}}((\pi_h^p u)|_{\omega_{\mathbf{a}}}). \quad (4.9)$$

Since the computations of  $\pi_h^p u$  and  $s_h^\alpha((\pi_h^p u)|_{\omega_\alpha})$  amount to element-wise/patch-wise finite element solves,  $\mathcal{J}_h^p u$  is computable. This operator is also local in the sense that for any element  $K \in \mathcal{T}_h$ ,  $(\mathcal{J}_h^p u)|_K$  only depends on the values of  $u$  in the vertex patch subdomains  $\omega_\alpha$  for the vertices  $\mathbf{a} \in \mathcal{V}_K$ .

#### 4.7. $H^1$ seminorm error estimate for $\mathcal{J}_h^p$ with computable constants

We now state our main result on the approximation error of  $\mathcal{J}_h^p u$  in the  $H^1$  seminorm (energy norm). We present it in a Pythagoras form that we are lead to by the orthogonal projection property (3.2).

**Theorem 4.2** ( $H^1$  seminorm error estimate with computable constants). *Let  $u \in H_0^1(\Omega)$  and let the quasi-interpolation operator  $\mathcal{J}_h^p$  be given by (3.2), (4.1), (4.9). Then*

$$\begin{aligned} \|\nabla(u - \mathcal{J}_h^p(u))\|_K &\leq \left\{ \|\nabla_h(u - \pi_h^p u)\|_K^2 + \left( \sum_{\mathbf{a} \in \mathcal{V}_K} \rho_\alpha \lambda_\alpha \|\nabla_h(u - \pi_h^p u)\|_{\omega_\alpha} \right)^2 \right\}^{1/2} \\ &\leq \left( \frac{1}{(d+1)} + (d+1)c_K^2 \right)^{1/2} \left\{ \sum_{\mathbf{a} \in \mathcal{V}_K} \|\nabla_h(u - \pi_h^p u)\|_{\omega_\alpha}^2 \right\}^{1/2} \quad \forall K \in \mathcal{T}_h, \end{aligned} \quad (4.10)$$

with the computable constant

$$c_K := \max_{\mathbf{a} \in \mathcal{V}_K} (\rho_\alpha \lambda_\alpha).$$

There also holds

$$\|\nabla(u - \mathcal{J}_h^p u)\|_\Omega \leq (1 + (d+1)^2 c_\Omega^2)^{1/2} \|\nabla_h(u - \pi_h^p u)\|_\Omega \quad (4.11)$$

with the computable constant

$$c_\Omega := \max_{\mathbf{a} \in \mathcal{V}_h} (\rho_\alpha \lambda_\alpha).$$

Computable quasi-interpolation estimates follow from combination with (4.8).

#### 4.8. $L^2$ error estimate for $\mathcal{J}_h^p$ with computable constants

We state here our main result on the approximation error of  $\mathcal{J}_h^p u$  in the  $L^2$  norm. We present it in a triangle-inequality form, since there is no orthogonal projection here. Recall the definitions of  $c_K$  and  $c_\Omega$  above.

**Theorem 4.3** ( $L^2$  error estimate with computable constants). *Let  $u \in H_0^1(\Omega)$  and let the quasi-interpolation operator  $\mathcal{J}_h^p$  be given by (3.2), (4.1), (4.9). Then*

$$\begin{aligned} \|u - \mathcal{J}_h^p u\|_K &\leq \frac{h_K}{\pi} \|\nabla_h(u - \pi_h^p u)\|_K + \frac{2}{d} \sum_{\mathbf{a} \in \mathcal{V}_K} \rho_\alpha \lambda_\alpha h_\alpha \|\nabla_h(u - \pi_h^p u)\|_{\omega_\alpha} \\ &\leq \left( \frac{1}{\pi} + \frac{2c_K}{d} \right) \sum_{\mathbf{a} \in \mathcal{V}_K} h_\alpha \|\nabla_h(u - \pi_h^p u)\|_{\omega_\alpha} \quad \forall K \in \mathcal{T}_h. \end{aligned} \quad (4.12)$$

There also holds

$$\|u - \mathcal{J}_h^p u\|_\Omega \leq \left( \frac{1}{\pi \sqrt{d+1}} + \frac{2}{d} \sqrt{d+1} c_\Omega \right) \left\{ \sum_{\mathbf{a} \in \mathcal{V}_h} h_\alpha^2 \|\nabla_h(u - \pi_h^p u)\|_{\omega_\alpha}^2 \right\}^{1/2}. \quad (4.13)$$

Computable quasi-interpolation estimates follow from combination with (4.8).

#### 4.9. Projection property

From (4.11) or (4.13), when  $u \in \mathcal{P}_p(\mathcal{T}_h)$ , then  $u = \pi_h^p u$ . Hence, we immediately have:

**Theorem 4.4** (Projection). *The quasi-interpolation operator  $\mathcal{J}_h^p$  from (3.2), (4.1), (4.9) is a projector, i.e.,*

$$\mathcal{J}_h^p(u) = u \quad \forall u \in \mathcal{P}_p(\mathcal{T}_h) \cap H_0^1(\Omega).$$

### 5. PROOFS OF THE MAIN RESULTS

#### 5.1. Preliminary results and notation

For each vertex  $\mathbf{a} \in \mathcal{V}_h$ , there exists a finite set  $\widehat{R}(\kappa_{\mathcal{T}_a})$  of reference patches only depending on the shape-regularity parameter  $\kappa_{\mathcal{T}_a}$  and the space dimension  $d$  and satisfying the following properties. Each  $\widehat{\mathcal{T}}_0 \in \widehat{R}(\kappa_{\mathcal{T}_a})$  is a conforming mesh of simplices sharing the vertex  $\mathbf{0}$ , and  $\widehat{\psi}_0$  denotes the piecewise affine ‘‘hat function’’ taking value 1 at  $\mathbf{0}$  and 0 at all the other vertices of  $\widehat{\mathcal{T}}_0$ ;  $\widehat{\omega}_0$  is the open domain corresponding to  $\widehat{\mathcal{T}}_0$ . All the elements  $\widehat{K} \in \widehat{\mathcal{T}}_0$  satisfy

$$c(\kappa_{\mathcal{T}_a}, d) \leq \rho_{\widehat{K}}, \quad h_{\widehat{K}} \leq C(\kappa_{\mathcal{T}_a}, d),$$

for two generic constants only depending on  $\kappa_{\mathcal{T}_a}$  and  $d$ . In addition, there exists a piecewise affine bilipchitz mapping  $\phi : \widehat{\omega}_0 \rightarrow \omega_a$  that transforms the elements of  $\widehat{\mathcal{T}}_0$  into those of  $\mathcal{T}_a$ . We denote by  $\phi^g$  and  $\phi^d$  the associated gradient- and divergence-preserving Piola mappings,  $\phi^g : L^2(\widehat{\omega}_0) \rightarrow L^2(\omega_a)$  and  $\phi^d : \mathbf{L}^2(\widehat{\omega}_0) \rightarrow \mathbf{L}^2(\omega_a)$  respectively defined by

$$\phi^g(\widehat{v}) := \widehat{v} \circ \phi^{-1} \quad \text{and} \quad \phi^d(\widehat{\mathbf{w}}) := \left( \frac{\mathbb{J}}{|\mathbb{J}|} \widehat{\mathbf{w}} \right) \circ \phi^{-1} \quad (5.1)$$

for all  $\widehat{v} \in L^2(\widehat{\omega}_0)$  and  $\widehat{\mathbf{w}} \in \mathbf{L}^2(\widehat{\omega}_0)$ , where  $\mathbb{J}$  is the Jacobian matrix of  $\phi$  and  $|\mathbb{J}|$  is its determinant (see, e.g., [15], Sect. 7.2). The following properties of Piola mappings will be useful. First, if  $\widehat{v} \in H^1(\widehat{\mathcal{T}}_0)$  and  $\widehat{\mathbf{w}} \in \mathbf{L}^2(\widehat{\omega}_0)$ , we have

$$(\nabla_h(\phi^g \widehat{v}), \phi^d \widehat{\mathbf{w}})_{\omega_a} = (\nabla_h \widehat{v}, \widehat{\mathbf{w}})_{\widehat{\omega}_0}. \quad (5.2)$$

The Piola mappings are invertible, and  $\widehat{\psi}_0 := (\phi^g)^{-1} \psi_a$ . We have  $\phi^g(\widehat{\psi}_0 \widehat{v}) = \psi_a \phi^g(\widehat{v})$  and

$$\|\nabla_h(\phi^g \widehat{v})\|_{\omega_a} \leq C(\kappa_{\mathcal{T}_a}, d) h_a^{d/2-1} \|\nabla_h \widehat{v}\|_{\widehat{\omega}_0} \quad (5.3)$$

for all  $\widehat{v} \in H^1(\widehat{\mathcal{T}}_0)$ , with  $h_a$  introduced in (2.4). Besides,  $\phi^d$  is an isomorphism between  $\mathcal{RT}_p(\widehat{\mathcal{T}}_0) \cap \mathbf{H}(\text{div}^0, \widehat{\omega}_0)$  and  $\mathcal{RT}_p(\mathcal{T}_a) \cap \mathbf{H}(\text{div}^0, \omega_a)$ , and we have

$$\|\phi^d \widehat{\mathbf{w}}\|_{\omega_a} \leq C(\kappa_{\mathcal{T}_a}, d) h_a^{1-d/2} \|\widehat{\mathbf{w}}\|_{\widehat{\omega}_0} \quad \forall \widehat{\mathbf{w}} \in \mathbf{L}^2(\widehat{\omega}_0). \quad (5.4)$$

We denote by  $\widehat{\mathcal{F}}_0$  the faces of a reference patch  $\widehat{\mathcal{T}}_0$ . The subset

$$\widehat{\mathcal{F}}_0^i := \left\{ \widehat{F} \in \widehat{\mathcal{F}}_0 \mid \widehat{\psi}_0|_{\widehat{F}} \neq 0 \right\},$$

coinciding with interior faces for an interior vertex and in general the faces sharing  $\mathbf{0}$ , will be useful. If  $\widehat{F} \in \widehat{\mathcal{F}}_0$  is a face,  $\mathcal{P}_q(\widehat{F})$  is the set of functions  $v : \widehat{F} \rightarrow \mathbb{R}$  that are polynomial of degree less than or equal to  $q$  in the  $(d-1)$  directions tangential to  $\widehat{F}$ . For a collection of faces  $\widehat{\mathcal{F}} \subset \widehat{\mathcal{F}}_0$ , we write  $\mathcal{P}_q(\widehat{\mathcal{F}}) := \cup_{\widehat{F} \in \widehat{\mathcal{F}}} \mathcal{P}_q(\widehat{F})$ . We also associate with each face  $\widehat{F} \in \widehat{\mathcal{F}}_0$  a unit normal vector  $\mathbf{n}_{\widehat{F}}$ . If  $\widehat{F} \subset \partial \widehat{\omega}_0$ , we assume that  $\mathbf{n}_{\widehat{F}}$  points outward  $\widehat{\omega}_0$ . For interior faces, the orientation of  $\mathbf{n}_{\widehat{F}}$  is arbitrary, but fixed. If  $\widehat{v} \in \mathcal{P}_{p+1}(\widehat{\mathcal{T}}_0)$ , the jump of  $\widehat{v}$  of through an interior face  $\widehat{F} = \partial \widehat{K}_- \cap \partial \widehat{K}_+ \in \widehat{\mathcal{F}}_0$  is defined by

$$[[\widehat{v}]]_{\widehat{F}} := \widehat{v}_-|_{\widehat{F}}(\mathbf{n}_- \cdot \mathbf{n}_{\widehat{F}}) + \widehat{v}_+|_{\widehat{F}}(\mathbf{n}_+ \cdot \mathbf{n}_{\widehat{F}}),$$

where  $\widehat{v}_\pm$  is the restriction of  $\widehat{v}$  to  $\widehat{K}_\pm$  and  $\mathbf{n}_\pm$  is the unit normal vector to  $\partial\widehat{K}_\pm$  pointing outward  $\widehat{K}_\pm$ ; note that  $\mathbf{n}_\pm \cdot \mathbf{n}_{\widehat{F}} = \pm 1$  only determines the sign. For exterior faces  $\widehat{F} \subset \partial\widehat{\omega}_0$ , we simply set

$$[[\widehat{v}]]_{\widehat{F}} = \widehat{v}|_{\widehat{F}}.$$

Thanks to these definitions, the integration by parts formula

$$(\nabla_h \widehat{v}, \widehat{\mathbf{w}})_{\widehat{\omega}_0} = \sum_{\widehat{F} \in \widehat{\mathcal{F}}_0} ([[ \widehat{v} ]]_{\widehat{F}}, \widehat{\mathbf{w}} \cdot \mathbf{n}_{\widehat{F}})_{\widehat{F}} - (\widehat{v}, \nabla \cdot \widehat{\mathbf{w}})_{\widehat{\omega}_0} \quad (5.5)$$

holds true for all  $\widehat{\mathbf{w}} \in \mathcal{RT}_p(\widehat{\mathcal{T}}_0) \cap \mathbf{H}(\operatorname{div}, \widehat{\omega}_0)$  and  $\widehat{v} \in \mathcal{P}_{p+1}(\widehat{\mathcal{T}}_0)$ .

## 5.2. Proof of the properties of $\lambda_a$ from (4.6)

Here, we establish the following important result:

**Theorem 5.1** (Upper bound on  $\lambda_a$ ). *The constant  $\lambda_a$  from (4.6) is finite and only depends on the shape-regularity parameter of  $\mathcal{T}_a$ , the space dimension  $d$ , and the polynomial degree  $p$ , i.e.,*

$$\lambda_a \leq C(\kappa_{\mathcal{T}_a}, d, p). \quad (5.6)$$

In addition, we have

$$\|\nabla_h(\mathcal{I}_h^p(\psi_a u_h) - s_h^\alpha(u_h))\|_{\omega_a} \leq \lambda_a \min_{v \in H_0^1(\omega_a)} \|\nabla_h(\psi_a u_h - v)\|_{\omega_a} \quad (5.7)$$

for all  $u_h \in \mathcal{P}_p(\mathcal{T}_a)$ .

Before proving Theorem 5.1, we first establish a few intermediate results. If  $\widehat{\mathcal{T}}_0$  is a reference patch, we use the notation  $\widehat{\mathbf{W}}_h^0 := \mathcal{RT}_p(\widehat{\mathcal{T}}_0) \cap \mathbf{H}(\operatorname{div}^0, \widehat{\omega}_0)$  for the counterpart to  $\mathbf{W}_h^\alpha$  of (4.2). We first show that even though functions in  $\widehat{\mathbf{W}}_h^0$  satisfy a divergence constraint, we are still free to assign their normal traces on relevant faces (faces from  $\widehat{\mathcal{F}}_0^i$ , namely interior faces for interior vertices, and in general the faces sharing  $\mathbf{0}$ ).

**Lemma 5.2** (Traces of Raviart–Thomas elements). *Let  $\widehat{\mathcal{T}}_0$  be a reference patch. For each  $\widehat{F} \in \widehat{\mathcal{F}}_0^i$ , consider  $\widehat{q}_{\widehat{F}} \in \mathcal{P}_p(\widehat{F})$ . Then, there exists  $\widehat{\mathbf{w}}_h \in \widehat{\mathbf{W}}_h^0$  such that  $\widehat{\mathbf{w}}_h \cdot \mathbf{n}_{\widehat{F}} = \widehat{q}_{\widehat{F}}$  for all  $\widehat{F} \in \widehat{\mathcal{F}}_0^i$ .*

*Proof.* We will construct a suitable function  $\widehat{\mathbf{w}}_h \in \widehat{\mathbf{W}}_h^0$  element by element. Let  $\widehat{K} \in \widehat{\mathcal{T}}_0$  be fixed. Note that given  $\widehat{r}_{\widehat{F}} \in \mathcal{P}_p(\widehat{\mathcal{F}}_{\widehat{K}})$  and  $\widehat{r}_{\widehat{K}} \in \mathcal{P}_p(\widehat{K})$  (which we actually take zero), there exists a function  $\widehat{\mathbf{w}}_{\widehat{K}} \in \mathcal{RT}_p(\widehat{K})$  such that  $\widehat{\mathbf{w}}_{\widehat{K}} \cdot \mathbf{n}_{\widehat{K}} = \widehat{r}_{\widehat{F}}$  on  $\partial\widehat{K}$  and  $\nabla \cdot \widehat{\mathbf{w}}_{\widehat{K}} = \widehat{r}_{\widehat{K}}$  in  $\widehat{K}$  if and only if the Neumann compatibility condition  $(\widehat{r}_{\widehat{K}}, 1)_{\widehat{K}} = (\widehat{r}_{\widehat{F}}, 1)_{\partial\widehat{K}}$  is satisfied. The key observation is that there is always a face  $\widehat{F}^* \in \widehat{\mathcal{F}}_{\widehat{K}}$  such that  $\widehat{F}^* \notin \widehat{\mathcal{F}}_0^i$ , i.e.,  $\widehat{\psi}_0|_{\widehat{F}^*} = 0$ : this is the face opposite to the vertex  $\mathbf{0}$ . Then, we select a boundary datum  $\widehat{r}_{\widehat{F}} \in \mathcal{P}_p(\widehat{\mathcal{F}}_{\widehat{K}})$  such that  $\widehat{r}_{\widehat{F}}|_{\widehat{F}} = \widehat{q}_{\widehat{F}}$  for all faces  $\widehat{F}$  of  $\widehat{K}$  such that  $\widehat{F} \in \widehat{\mathcal{F}}_0^i$  and, by fixing a suitable Neumann value on  $\widehat{F}^*$ , such that  $(\widehat{r}_{\widehat{F}}, 1)_{\widehat{\mathcal{F}}_{\widehat{K}}} = 0$ . Finally, we are eligible to impose  $\widehat{\mathbf{w}}_h|_{\widehat{K}} \cdot \mathbf{n}_{\widehat{K}} = \widehat{r}_{\widehat{F}}$  on  $\partial\widehat{K}$  and  $\nabla \cdot (\widehat{\mathbf{w}}_h|_{\widehat{K}}) = 0$ .  $\square$

By carefully selecting their normal traces, we can then show that functions in  $\widehat{\mathbf{W}}_h^0$  (and, after mapping, in  $\mathbf{W}_h^\alpha$ ) can be successfully employed to measure the level of nonconformity of a piecewise polynomial function.

**Corollary 5.3** (Control of the nonconformity). *For all  $\widehat{u}_h \in \mathcal{P}_p(\widehat{\mathcal{T}}_0)$ , if*

$$\left( \nabla_h(\widehat{\psi}_0 \widehat{u}_h), \widehat{\mathbf{v}}_h \right)_{\widehat{\omega}_0} = 0 \quad \forall \widehat{\mathbf{v}}_h \in \widehat{\mathbf{W}}_h^0,$$

then  $\widehat{\psi}_0 \widehat{u}_h \in H_0^1(\widehat{\omega}_0)$ . In particular,  $\widehat{u}_h \in H^1(\widehat{\omega}_0)$ .

*Proof.* Let  $\hat{u}_h \in \mathcal{P}_p(\hat{\mathcal{T}}_0)$ . For all  $\hat{v}_h \in \widehat{\mathcal{W}}_h^0$ , since  $\nabla \cdot \hat{v}_h = 0$ , Stokes' formula (5.5) shows that

$$\left( \nabla_h \left( \hat{\psi}_0 \hat{u}_h \right), \hat{v}_h \right)_{\hat{\omega}_0} = \sum_{\hat{F} \in \hat{\mathcal{F}}_0} \left( \left[ \hat{\psi}_0 \hat{u}_h \right]_{\hat{F}}, \hat{v}_h \cdot \mathbf{n}_{\hat{F}} \right)_{\hat{F}} = \sum_{\hat{F} \in \hat{\mathcal{F}}_0^i} \left( \hat{\psi}_0 \llbracket \hat{u}_h \rrbracket_{\hat{F}}, \hat{v}_h \cdot \mathbf{n}_{\hat{F}} \right)_{\hat{F}},$$

since  $\hat{\psi}_0$  is continuous and  $\hat{\psi}_0|_{\hat{F}} = 0$  for all  $\hat{F} \in \hat{\mathcal{F}}_0 \setminus \hat{\mathcal{F}}_0^i$ . Following Lemma 5.2, we can now pick  $\hat{v}_h^* \in \widehat{\mathcal{W}}_h^0$  such that  $\hat{v}_h^* \cdot \mathbf{n}_{\hat{F}} = \llbracket \hat{u}_h \rrbracket_{\hat{F}}$  for all  $\hat{F} \in \hat{\mathcal{F}}_0^i$ . This gives

$$\sum_{\hat{F} \in \hat{\mathcal{F}}_0^i} \left\| \hat{\psi}_0^{1/2} \llbracket \hat{u}_h \rrbracket_{\hat{F}} \right\|_{\hat{F}}^2 = \left( \nabla_h \left( \hat{\psi}_0 \hat{u}_h \right), \hat{v}_h^* \right)_{\hat{\omega}_0} = 0.$$

Since  $\hat{\psi}_0 > 0$  a.e. on each  $\hat{F} \in \hat{\mathcal{F}}_0^i$ , this shows that  $\llbracket \hat{u}_h \rrbracket_{\hat{F}} = 0$  for all  $\hat{F} \in \hat{\mathcal{F}}_0^i$ . Hence  $\hat{u}_h \in H^1(\hat{\omega}_0)$ . There also holds  $\hat{u}_h = 0$  on  $\hat{F} \subset \partial \hat{\omega}_0$ ,  $\hat{F} \in \hat{\mathcal{F}}_0^i$  (such faces only exist when the vertex  $\mathbf{0}$  lies on the boundary of  $\hat{\omega}_0$ ). Thus, it immediately follows that  $\hat{\psi}_0 \hat{u}_h \in H_0^1(\hat{\omega}_0)$ .  $\square$

We are now ready to establish our main result concerning  $\lambda_{\mathbf{a}}$ . We apply below the argument of equivalence of norms in finite dimension, which is not explicit/optimal in the polynomial degree  $p$ . We actually do not search to be explicit and optimal in  $p$ , we are merely justifying that the estimates of Theorems 4.2 and 4.3 indeed come with all constants bounded (and fully computable).

*Proof of Theorem 5.1.* Fix a vertex  $\mathbf{a} \in \mathcal{V}_h$ , and consider the associated reference patch  $\hat{\mathcal{T}}_0 \in \hat{R}(\kappa_{\mathcal{T}_{\mathbf{a}}})$ .

If  $\hat{u}_h \in \mathcal{P}_p(\hat{\mathcal{T}}_0)$ , we define  $\hat{s}_h^0(\hat{u}_h) \in \mathcal{P}_p(\hat{\mathcal{T}}_0) \cap H_0^1(\hat{\omega}_0)$  by requiring that

$$\left( \nabla_h \hat{s}_h^0(\hat{u}_h), \nabla_h \hat{v}_h \right)_{\hat{\omega}_0} = \left( \nabla_h \left( \mathcal{I}_h^p \left( \hat{\psi}_0 \hat{u}_h \right) \right), \nabla_h \hat{v}_h \right)_{\hat{\omega}_0} \quad \forall \hat{v}_h \in \mathcal{P}_p(\hat{\mathcal{T}}_0) \cap H_0^1(\hat{\omega}_0), \quad (5.8)$$

and similarly we let  $\hat{\mathbf{r}}_h^0(\hat{u}_h)$  be the only element of  $\widehat{\mathcal{W}}_h^0$  such that

$$\left( \hat{\mathbf{r}}_h^0(\hat{u}_h), \hat{\mathbf{w}}_h \right)_{\hat{\omega}_0} = \left( \nabla_h \left( \hat{\psi}_0 \hat{u}_h \right), \hat{\mathbf{w}}_h \right)_{\hat{\omega}_0} \quad \forall \hat{\mathbf{w}}_h \in \widehat{\mathcal{W}}_h^0. \quad (5.9)$$

These are the natural counterparts to  $s_h^{\mathbf{a}}(u_h)$  from (4.1) and  $\mathbf{r}_h^{\mathbf{a}}(u_h)$  from (4.3) on the reference patch  $\hat{\mathcal{T}}_0$ .

Now consider an arbitrary  $u_h \in \mathcal{P}_p(\mathcal{T}_{\mathbf{a}})$ . Since  $s_h^{\mathbf{a}}(u_h)$  is the minimizer, see (4.1a), we have

$$\begin{aligned} \left\| \nabla_h \left( \mathcal{I}_h^p(\psi_{\mathbf{a}} u_h) - s_h^{\mathbf{a}}(u_h) \right) \right\|_{\omega_{\mathbf{a}}} &\leq \left\| \nabla_h \left( \mathcal{I}_h^p(\psi_{\mathbf{a}} u_h) - \phi^{\mathcal{S}}(\hat{v}_h) \right) \right\|_{\omega_{\mathbf{a}}} \\ &\leq C(\kappa_{\mathcal{T}_{\mathbf{a}}}, d) h_{\mathbf{a}}^{d/2-1} \left\| \nabla_h \left( \hat{\mathcal{I}}_h^p \left( \hat{\psi}_0 \hat{u}_h \right) - \hat{v}_h \right) \right\|_{\hat{\omega}_0} \end{aligned}$$

for any  $\hat{v}_h \in \mathcal{P}_p(\hat{\mathcal{T}}_0) \cap H_0^1(\hat{\omega}_0)$ , with  $\hat{u}_h := (\phi^{\mathcal{S}})^{-1}(u_h) \in \mathcal{P}_p(\hat{\mathcal{T}}_0)$ . Here, we employed the fact that  $\phi^{\mathcal{S}}$  preserves piecewise polynomial functions as well as point values in the Lagrange interpolation nodes (affine-equivalent degrees of freedom) and used (5.3) in the last inequality. We can minimize the last norm in the right-hand side, leading to

$$\left\| \nabla_h \left( \mathcal{I}_h^p(\psi_{\mathbf{a}} u_h) - s_h^{\mathbf{a}}(u_h) \right) \right\|_{\omega_{\mathbf{a}}} \leq C(\kappa_{\mathcal{T}_{\mathbf{a}}}, d) h_{\mathbf{a}}^{d/2-1} \left\| \nabla_h \left( \hat{\mathcal{I}}_h^p \left( \hat{\psi}_0 \hat{u}_h \right) - \hat{s}_h^0(\hat{u}_h) \right) \right\|_{\hat{\omega}_0}, \quad (5.10)$$

by definition (5.8) of  $\hat{s}_h^0(\hat{u}_h)$ .

On the other hand, still for the chosen  $u_h \in \mathcal{P}_p(\mathcal{T}_{\mathbf{a}})$ , because of (5.2), for any  $\hat{\mathbf{w}}_h \in \widehat{\mathcal{W}}_h^0$ , we also have

$$\left( \nabla_h(\psi_{\mathbf{a}} u_h), \phi^{\mathcal{d}}(\hat{\mathbf{w}}_h) \right)_{\omega_{\mathbf{a}}} = \left( \nabla_h \left( \hat{\psi}_0 \hat{u}_h \right), \hat{\mathbf{w}}_h \right)_{\hat{\omega}_0}. \quad (5.11)$$

Moreover, as in (4.4), from (5.9), there holds

$$\|\widehat{\mathbf{r}}_h^{\mathbf{0}}(\widehat{u}_h)\|_{\widehat{\omega}_0} = \max_{\substack{\widehat{\mathbf{w}}_h \in \widehat{\mathbf{W}}_h^{\mathbf{0}} \\ \|\widehat{\mathbf{w}}_h\|_{\widehat{\omega}_0}=1}} \left( \nabla_h(\widehat{\psi}_0 \widehat{u}_h), \widehat{\mathbf{w}}_h \right)_{\widehat{\omega}_0}. \quad (5.12)$$

Thus, using the maximizer from (5.12),  $\widehat{\mathbf{w}}_h^* \in \widehat{\mathbf{W}}_h^{\mathbf{0}}$  with  $\|\widehat{\mathbf{w}}_h^*\|_{\widehat{\omega}_0} = 1$ , and employing  $\phi^d(\widehat{\mathbf{w}}_h^*) / \|\phi^d(\widehat{\mathbf{w}}_h^*)\|_{\omega_\alpha}$  in (4.4), we have

$$\begin{aligned} \|\mathbf{r}_h^\alpha(u_h)\|_{\omega_\alpha} &\stackrel{(4.4)}{\geq} \frac{1}{\|\phi^d(\widehat{\mathbf{w}}_h^*)\|_{\omega_\alpha}} \left( \nabla_h(\psi_\alpha u_h), \phi^d(\widehat{\mathbf{w}}_h^*) \right)_{\omega_\alpha} \\ &\stackrel{(5.11)}{=} \frac{1}{\|\phi^d(\widehat{\mathbf{w}}_h^*)\|_{\omega_\alpha}} \left( \nabla_h(\widehat{\psi}_0 \widehat{u}_h), \widehat{\mathbf{w}}_h^* \right)_{\widehat{\omega}_0} \stackrel{(5.12)}{=} \frac{1}{\|\phi^d(\widehat{\mathbf{w}}_h^*)\|_{\omega_\alpha}} \|\widehat{\mathbf{r}}_h^{\mathbf{0}}(\widehat{u}_h)\|_{\widehat{\omega}_0}. \end{aligned}$$

Finally, we use (5.4) to establish that

$$\|\phi^d(\widehat{\mathbf{w}}_h^*)\|_{\omega_\alpha} \leq C(\kappa_{\mathcal{T}_\alpha}, d) h_\alpha^{1-d/2} \|\widehat{\mathbf{w}}_h^*\|_{\widehat{\omega}_0} = C(\kappa_{\mathcal{T}_\alpha}, d) h_\alpha^{1-d/2},$$

which leads to

$$\frac{1}{\|\mathbf{r}_h^\alpha(u_h)\|_{\omega_\alpha}} \leq C(\kappa_{\mathcal{T}_\alpha}, d) \frac{h_\alpha^{1-d/2}}{\|\widehat{\mathbf{r}}_h^{\mathbf{0}}(\widehat{u}_h)\|_{\widehat{\omega}_0}}. \quad (5.13)$$

Combining (5.10) and (5.13) gives

$$\frac{\|\nabla_h(\mathcal{I}_h^p(\psi_\alpha u_h) - s_h^\alpha(u_h))\|_{\omega_\alpha}}{\|\mathbf{r}_h^\alpha(u_h)\|_{\omega_\alpha}} \leq C(\kappa_{\mathcal{T}_\alpha}, d) \frac{\left\| \nabla_h \left( \mathcal{I}_h^p(\widehat{\psi}_0 \widehat{u}_h) - \widehat{s}_h^{\mathbf{0}}(\widehat{u}_h) \right) \right\|_{\widehat{\omega}_0}}{\|\widehat{\mathbf{r}}_h^{\mathbf{0}}(\widehat{u}_h)\|_{\widehat{\omega}_0}} \leq C(\kappa_{\mathcal{T}_\alpha}, d) \widehat{\lambda}_0,$$

where

$$\widehat{\lambda}_0 := \max_{\widehat{u}_h \in \mathcal{P}_p(\widehat{\mathcal{T}}_0)} \frac{\left\| \nabla_h \left( \mathcal{I}_h^p(\widehat{\psi}_0 \widehat{u}_h) - \widehat{s}_h^{\mathbf{0}}(\widehat{u}_h) \right) \right\|_{\widehat{\omega}_0}}{\|\widehat{\mathbf{r}}_h^{\mathbf{0}}(\widehat{u}_h)\|_{\widehat{\omega}_0}}.$$

In other words, since the last inequality is valid for all  $u_h \in \mathcal{P}_p(\mathcal{T}_\alpha)$ , we have actually shown that  $\lambda_\alpha \leq C(\kappa_{\mathcal{T}_\alpha}, d) \widehat{\lambda}_0$ .

We next show that  $\widehat{\lambda}_0$  is finite, only depending on the shape regularity  $\kappa_{\mathcal{T}_\alpha}$ , the space dimension  $d$ , and the polynomial degree  $p$ . Since  $\mathcal{P}_p(\widehat{\mathcal{T}}_0)$  is a finite-dimensional vector space (with dimension only depending on  $\kappa_{\mathcal{T}_\alpha}$ ,  $d$ , and  $p$ ), it suffices to show that  $\|\widehat{\mathbf{r}}_h^{\mathbf{0}}(\widehat{u}_h)\|_{\widehat{\omega}_0} = 0$  implies that  $\|\nabla_h(\mathcal{I}_h^p(\widehat{\psi}_0 \widehat{u}_h) - \widehat{s}_h^{\mathbf{0}}(\widehat{u}_h))\|_{\widehat{\omega}_0} = 0$  for all  $\widehat{u}_h \in \mathcal{P}_p(\widehat{\mathcal{T}}_0)$ . Thus, we consider  $\widehat{u}_h \in \mathcal{P}_p(\widehat{\mathcal{T}}_0)$  such that  $\|\widehat{\mathbf{r}}_h^{\mathbf{0}}(\widehat{u}_h)\|_{\widehat{\omega}_0} = 0$ . Due to (5.12) and Corollary 5.3,  $\widehat{\psi}_0 \widehat{u}_h \in \mathcal{P}_{p+1}(\widehat{\mathcal{T}}_0) \cap H_0^1(\widehat{\omega}_0)$ . Then, using the minimization property of  $\widehat{s}_h^{\mathbf{0}}(\widehat{u}_h)$  that follows from (5.8), cf. (4.1a), we have

$$\left\| \nabla_h \left( \mathcal{I}_h^p(\widehat{\psi}_0 \widehat{u}_h) - \widehat{s}_h^{\mathbf{0}}(\widehat{u}_h) \right) \right\|_{\widehat{\omega}_0} \leq \left\| \nabla_h \left( \mathcal{I}_h^p(\widehat{\psi}_0 \widehat{u}_h) - \mathcal{I}_h^p(\widehat{\psi}_0 \widehat{u}_h) \right) \right\|_{\widehat{\omega}_0} = 0,$$

since  $\mathcal{I}_h^p(\widehat{\psi}_0 \widehat{u}_h) \in \mathcal{P}_p(\widehat{\mathcal{T}}_0) \cap H_0^1(\widehat{\omega}_0)$ . Thus, we have established (5.6).

It remains to establish the bound (5.7). We first note that

$$\left\| \nabla_h(\mathcal{I}_h^p(\psi_\alpha u_h) - s_h^\alpha(u_h)) \right\|_{\omega_\alpha} \leq \lambda_\alpha \|\mathbf{r}_h^\alpha(u_h)\|_{\omega_\alpha} \quad \forall u_h \in \mathcal{P}_p(\mathcal{T}_\alpha)$$

by the definition of  $\lambda_\alpha$ . On the other hand, using (4.4), we have

$$\|\mathbf{r}_h^\alpha(u_h)\|_{\omega_\alpha} = \max_{\substack{\mathbf{w}_h \in \mathbf{W}_h^\alpha \\ \|\mathbf{w}_h\|_{\omega_\alpha}=1}} \left( \nabla_h(\psi_\alpha u_h), \mathbf{w}_h \right)_{\omega_\alpha} = \max_{\substack{\mathbf{w}_h \in \mathbf{W}_h^\alpha \\ \|\mathbf{w}_h\|_{\omega_\alpha}=1}} \left( \nabla_h(\psi_\alpha u_h - v), \mathbf{w}_h \right)_{\omega_\alpha}$$

$$\leq \|\nabla_h(\psi_{\mathbf{a}}u_h - v)\|_{\omega_{\mathbf{a}}}$$

for all  $v \in H_0^1(\omega_{\mathbf{a}})$ , since we always have

$$(\nabla_h v, \mathbf{w}_h)_{\omega_{\mathbf{a}}} = (\nabla v, \mathbf{w}_h)_{\omega_{\mathbf{a}}} = -(v, \nabla \cdot \mathbf{w}_h)_{\omega_{\mathbf{a}}} = 0$$

for all  $\mathbf{w}_h \in \mathbf{W}_h^{\mathbf{a}}$ . □

### 5.3. Proof of $H^1$ seminorm error estimates of Theorem 4.2

In this section, we now derive the local and global error estimates in the  $H^1$  norm stated in Theorem 4.2. We start with a fundamental result concerning the local contributions  $s_h^{\mathbf{a}}$ .

**Lemma 5.4** (Patch-wise estimates). *The estimate*

$$\|\nabla_h(\mathcal{I}_h^p(\psi_{\mathbf{a}}\pi_h^p u) - s_h^{\mathbf{a}}((\pi_h^p u)|_{\omega_{\mathbf{a}}}))\|_{\omega_{\mathbf{a}}} \leq \rho_{\mathbf{a}}\lambda_{\mathbf{a}}\|\nabla_h(u - \pi_h^p u)\|_{\omega_{\mathbf{a}}} \quad (5.14)$$

holds true for each vertex  $\mathbf{a} \in \mathcal{V}_h$ .

*Proof.* We start with (5.7). Next, since  $\psi_{\mathbf{a}}u \in H_0^1(\omega_{\mathbf{a}})$ , we have

$$\min_{v \in H_0^1(\omega_{\mathbf{a}})} \|\nabla_h(\psi_{\mathbf{a}}\pi_h^p u - v)\|_{\omega_{\mathbf{a}}} \leq \|\nabla_h(\psi_{\mathbf{a}}(\pi_h^p u - u))\|_{\omega_{\mathbf{a}}}.$$

Recalling that  $(\pi_h^p u, 1)_K = (u, 1)_K$  for all  $K \in \mathcal{T}_{\mathbf{a}}$  from (3.2), we then employ the product rule element-wise to show that

$$\begin{aligned} \|\nabla(\psi_{\mathbf{a}}(u - \pi_h^p u))\|_K &\leq \|\nabla\psi_{\mathbf{a}}\|_{L^\infty(K)}\|u - \pi_h^p u\|_K + \|\psi_{\mathbf{a}}\|_{L^\infty(K)}\|\nabla(u - \pi_h^p u)\|_K \\ &\leq \left(1 + \frac{h_K}{\pi\tau_K^{\mathbf{a}}}\right)\|\nabla(u - \pi_h^p u)\|_K, \end{aligned}$$

where we employed (2.2) and (2.5). The desired result (5.14) follows by the definition of  $\rho_{\mathbf{a}}$  in (4.7). □

*Proof of Theorem 4.2.* From (3.2a), the Pythagoras equality yields

$$\|\nabla(u - \mathcal{J}_h^p(u))\|_K^2 = \|\nabla(u - \pi_h^p u)\|_K^2 + \|\nabla(\pi_h^p u - \mathcal{J}_h^p(u))\|_K^2.$$

The partition of unity (2.3) and the projection and linearity properties of  $\mathcal{I}_h^p$  give

$$\sum_{\mathbf{a} \in \mathcal{V}_h} \mathcal{I}_h^p(\psi_{\mathbf{a}}\pi_h^p u) = \mathcal{I}_h^p\left(\sum_{\mathbf{a} \in \mathcal{V}_h} (\psi_{\mathbf{a}}\pi_h^p u)\right) = \mathcal{I}_h^p(\pi_h^p u) = \pi_h^p u. \quad (5.15)$$

Thus, together with definition (4.9) and Lemma 5.4, we estimate the second term as

$$\begin{aligned} \|\nabla(\pi_h^p u - \mathcal{J}_h^p(u))\|_K &= \left\| \sum_{\mathbf{a} \in \mathcal{V}_K} \nabla(\mathcal{I}_h^p(\psi_{\mathbf{a}}\pi_h^p u) - s_h^{\mathbf{a}}((\pi_h^p u)|_{\omega_{\mathbf{a}}})) \right\|_K \\ &\leq \sum_{\mathbf{a} \in \mathcal{V}_K} \|\nabla(\mathcal{I}_h^p(\psi_{\mathbf{a}}\pi_h^p u) - s_h^{\mathbf{a}}((\pi_h^p u)|_{\omega_{\mathbf{a}}}))\|_K \\ &\leq \sum_{\mathbf{a} \in \mathcal{V}_K} \|\nabla_h(\mathcal{I}_h^p(\psi_{\mathbf{a}}\pi_h^p u) - s_h^{\mathbf{a}}((\pi_h^p u)|_{\omega_{\mathbf{a}}}))\|_{\omega_{\mathbf{a}}} \\ &\leq \sum_{\mathbf{a} \in \mathcal{V}_K} \rho_{\mathbf{a}}\lambda_{\mathbf{a}}\|\nabla_h(u - \pi_h^p u)\|_{\omega_{\mathbf{a}}}, \end{aligned}$$

and the first inequality in (4.10) immediately follows.

We also have

$$\begin{aligned} \left( \sum_{\mathbf{a} \in \mathcal{V}_K} \rho_{\mathbf{a}} \lambda_{\mathbf{a}} \|\nabla_h(u - \pi_h^p u)\|_{\omega_{\mathbf{a}}} \right)^2 &\leq (d+1) \sum_{\mathbf{a} \in \mathcal{V}_K} (\rho_{\mathbf{a}} \lambda_{\mathbf{a}})^2 \|\nabla_h(u - \pi_h^p u)\|_{\omega_{\mathbf{a}}}^2 \\ &\leq (d+1) \max_{\mathbf{a} \in \mathcal{V}_K} (\rho_{\mathbf{a}} \lambda_{\mathbf{a}})^2 \sum_{\mathbf{a} \in \mathcal{V}_K} \|\nabla_h(u - \pi_h^p u)\|_{\omega_{\mathbf{a}}}^2, \end{aligned}$$

which proves the second inequality in (4.10).

Finally, to establish (4.11), we treat sharply

$$\begin{aligned} \|\nabla_h(\pi_h^p u - \mathcal{I}_h^p(u))\|_{\Omega}^2 &= \sum_{K \in \mathcal{T}_h} \|\nabla(\pi_h^p u - \mathcal{I}_h^p(u))\|_K^2 \\ &\leq \sum_{K \in \mathcal{T}_h} (d+1) \sum_{\mathbf{a} \in \mathcal{V}_K} \|\nabla(\mathcal{I}_h^p(\psi_{\mathbf{a}} \pi_h^p u) - s_h^{\mathbf{a}}((\pi_h^p u)|_{\omega_{\mathbf{a}}}))\|_K^2 \\ &= (d+1) \sum_{\mathbf{a} \in \mathcal{V}_h} \|\nabla_h(\mathcal{I}_h^p(\psi_{\mathbf{a}} \pi_h^p u) - s_h^{\mathbf{a}}((\pi_h^p u)|_{\omega_{\mathbf{a}}}))\|_{\omega_{\mathbf{a}}}^2 \\ &\leq (d+1) \sum_{\mathbf{a} \in \mathcal{V}_h} (\rho_{\mathbf{a}} \lambda_{\mathbf{a}})^2 \|\nabla_h(u - \pi_h^p u)\|_{\omega_{\mathbf{a}}}^2 \\ &\leq \max_{\mathbf{a} \in \mathcal{V}_h} (\rho_{\mathbf{a}} \lambda_{\mathbf{a}})^2 (d+1)^2 \|\nabla_h(u - \pi_h^p u)\|_{\Omega}^2. \end{aligned} \tag{5.16}$$

□

#### 5.4. Proof of $L^2$ error estimates of Theorem 4.3

We finally prove the local and global error estimates presented in Theorem 4.3. We start by establishing a Poincaré inequality.

**Lemma 5.5** (Element-wise Poincaré). *Consider an element  $K \in \mathcal{T}_h$ , one of its vertices  $\mathbf{a} \in \mathcal{V}_K$  and the face  $F \in \mathcal{F}_K$  opposite  $\mathbf{a}$ . Then, if  $v \in H^1(K)$  satisfies  $v = 0$  on  $F$ , we have*

$$\|v\|_K \leq \frac{2h_K}{d} \|\nabla v\|_K. \tag{5.17}$$

*Proof.* Let  $\mathbf{y} : \mathbf{x} \rightarrow \mathbf{x} - \mathbf{a}$ . Since  $\nabla \cdot \mathbf{y} = d$ , we have

$$d \int_K |v|^2 = \int_K \nabla \cdot \mathbf{y} |v|^2 = \int_{\partial K} \mathbf{y} \cdot \mathbf{n} |v|^2 - 2 \int_K \mathbf{y} \cdot \nabla v v.$$

We then observe that the boundary term vanishes as  $\mathbf{y} \cdot \mathbf{n} = 0$  on all the faces  $F' \in \mathcal{F}_K$  sharing the vertex  $\mathbf{a}$ , and  $v = 0$  on the remaining face  $F$ . We then use that  $|\mathbf{y}| \leq h_K$ , and therefore

$$d \|v\|_K^2 \leq 2h_K \|\nabla v\|_K \|v\|_K$$

using the Cauchy–Schwarz inequality. The result follows. □

**Corollary 5.6** (Closeness to the local-best approximation in  $L^2$ ). *For all  $\mathbf{a} \in \mathcal{V}_h$ , we have*

$$\|\mathcal{I}_h^p(\psi_{\mathbf{a}} \pi_h^p u) - s_h^{\mathbf{a}}((\pi_h^p u)|_{\omega_{\mathbf{a}}})\|_{\omega_{\mathbf{a}}} \leq \frac{2}{d} h_{\mathbf{a}} \|\nabla_h(\mathcal{I}_h^p(\psi_{\mathbf{a}} \pi_h^p u) - s_h^{\mathbf{a}}((\pi_h^p u)|_{\omega_{\mathbf{a}}}))\|_{\omega_{\mathbf{a}}}. \tag{5.18}$$

In addition, the estimates

$$\|\pi_h^p u - \mathcal{J}_h^p u\|_K \leq \frac{2}{d} \sum_{\mathbf{a} \in \mathcal{V}_K} \rho_{\mathbf{a}} \lambda_{\mathbf{a}} h_{\mathbf{a}} \|\nabla_h(u - \pi_h^p u)\|_{\omega_{\mathbf{a}}} \quad (5.19)$$

and

$$\|\pi_h^p u - \mathcal{J}_h^p u\|_{\Omega} \leq \frac{2}{d} \sqrt{d+1} c_{\Omega} \left\{ \sum_{\mathbf{a} \in \mathcal{V}_h} h_{\mathbf{a}}^2 \|\nabla_h(u - \pi_h^p u)\|_{\omega_{\mathbf{a}}}^2 \right\}^{1/2} \quad (5.20)$$

hold true.

*Proof.* The estimate in (5.18) simply follows by applying Lemma 5.5 in each element of the patch, noting that, for each  $K \in \mathcal{T}_{\mathbf{a}}$ , both  $\mathcal{I}_h^p(\psi_{\mathbf{a}} \pi_h^p u)$  and  $s_h^{\mathbf{a}}(\pi_h^p u)|_{\omega_{\mathbf{a}}}$  vanish on the face of  $K$  opposite to  $\mathbf{a}$ , see (3.1) and (4.1).

For the second estimate (5.19), we observe from (5.15) and (4.9) that

$$(\pi_h^p u - \mathcal{J}_h^p u)|_K = \sum_{\mathbf{a} \in \mathcal{V}_K} \{\mathcal{I}_h^p(\psi_{\mathbf{a}} \pi_h^p u) - s_h^{\mathbf{a}}(\pi_h^p u)|_{\omega_{\mathbf{a}}}\}|_K$$

and therefore, using the triangle inequality together with (2.4) and (5.14),

$$\begin{aligned} \|\pi_h^p u - \mathcal{J}_h^p u\|_K &\leq \sum_{\mathbf{a} \in \mathcal{V}_K} \|\mathcal{I}_h^p(\psi_{\mathbf{a}} \pi_h^p u) - s_h^{\mathbf{a}}(\pi_h^p u)|_{\omega_{\mathbf{a}}}\|_K \\ &\leq \frac{2}{d} \sum_{\mathbf{a} \in \mathcal{V}_K} h_K \|\nabla_h(\mathcal{I}_h^p(\psi_{\mathbf{a}} \pi_h^p u) - s_h^{\mathbf{a}}(\pi_h^p u)|_{\omega_{\mathbf{a}}})\|_K, \\ &\leq \frac{2}{d} \sum_{\mathbf{a} \in \mathcal{V}_K} h_{\mathbf{a}} \|\nabla_h(\mathcal{I}_h^p(\psi_{\mathbf{a}} \pi_h^p u) - s_h^{\mathbf{a}}(\pi_h^p u)|_{\omega_{\mathbf{a}}})\|_{\omega_{\mathbf{a}}} \\ &\leq \frac{2}{d} \sum_{\mathbf{a} \in \mathcal{V}_K} h_{\mathbf{a}} \rho_{\mathbf{a}} \lambda_{\mathbf{a}} \|\nabla_h(u - \pi_h^p u)\|_{\omega_{\mathbf{a}}}. \end{aligned}$$

To establish (5.20), we estimate sharply, as in (5.16),

$$\begin{aligned} \|\pi_h^p u - \mathcal{J}_h^p u\|_{\Omega}^2 &\leq (d+1) \sum_{\mathbf{a} \in \mathcal{V}_h} \|\mathcal{I}_h^p(\psi_{\mathbf{a}} \pi_h^p u) - s_h^{\mathbf{a}}(\pi_h^p u)|_{\omega_{\mathbf{a}}}\|_{\omega_{\mathbf{a}}}^2 \\ &\stackrel{(5.18)}{\leq} \frac{4}{d^2} (d+1) \sum_{\mathbf{a} \in \mathcal{V}_h} h_{\mathbf{a}}^2 \|\nabla_h(\mathcal{I}_h^p(\psi_{\mathbf{a}} \pi_h^p u) - s_h^{\mathbf{a}}(\pi_h^p u)|_{\omega_{\mathbf{a}}})\|_{\omega_{\mathbf{a}}}^2 \\ &\stackrel{(5.14)}{\leq} \frac{4}{d^2} (d+1) \max_{\mathbf{a} \in \mathcal{V}_h} (\rho_{\mathbf{a}} \lambda_{\mathbf{a}})^2 \sum_{\mathbf{a} \in \mathcal{V}_h} h_{\mathbf{a}}^2 \|\nabla_h(u - \pi_h^p u)\|_{\omega_{\mathbf{a}}}^2. \end{aligned}$$

□

*Proof of Theorem 4.3.* The estimates follow from the ones in Corollary 5.6, triangle inequalities, and the fact that

$$\|u - \pi_h^p u\|_K \leq \frac{h_K}{\pi} \|\nabla(u - \pi_h^p u)\|_K$$

for all  $K \in \mathcal{T}_h$ . For (4.13), we namely use  $\|u - \mathcal{J}_h^p u\|_{\Omega} \leq \|u - \pi_h^p u\|_{\Omega} + \|\pi_h^p u - \mathcal{J}_h^p u\|_{\Omega}$  together with (5.20) and

$$\begin{aligned} \|u - \pi_h^p u\|_{\Omega}^2 &\leq \sum_{K \in \mathcal{T}_h} \frac{h_K^2}{\pi^2} \|\nabla(u - \pi_h^p u)\|_K^2 = \frac{1}{d+1} \sum_{\mathbf{a} \in \mathcal{V}_h} \sum_{K \in \mathcal{T}_h | \mathbf{a} \in \mathcal{V}_K} \frac{h_K^2}{\pi^2} \|\nabla(u - \pi_h^p u)\|_K^2 \\ &\leq \frac{1}{\pi^2 (d+1)} \sum_{\mathbf{a} \in \mathcal{V}_h} h_{\mathbf{a}}^2 \|\nabla_h(u - \pi_h^p u)\|_{\omega_{\mathbf{a}}}^2. \end{aligned}$$

□

## 6. NUMERICAL EXAMPLES

We present here the results of several numerical experiments illustrating the actual practical behavior of our quasi-interpolate.

### 6.1. Setting

For different two-dimensional domains  $\Omega \subset \mathbb{R}^2$ , functions  $u \in H_0^1(\Omega) \cap C^0(\overline{\Omega})$  with various additional regularity, and triangular meshes  $\mathcal{T}_h$ , we compute the (broken) local-best approximation  $\pi_h^1 u \in \mathcal{P}_1(\mathcal{T}_h)$ ,

$$\pi_h^1 u := \arg \min_{v_h \in \mathcal{P}_1(\mathcal{T}_h)} \|\nabla_h(u - v_h)\|_{\Omega},$$

the global-best approximation

$$u_h^1 := \arg \min_{v_h \in \mathcal{P}_1(\mathcal{T}_h) \cap H_0^1(\Omega)} \|\nabla(u - u_h^1)\|_{\Omega},$$

the Lagrange interpolant  $\mathcal{I}_h^1 u$  of Section 3.1, and our projection  $\mathcal{J}_h^1 u$  of Section 4.5, both for the polynomial degree  $p = 1$ . Except from the local-best approximation, all these approximations of  $u$  are conforming and sit in the finite element space  $\mathcal{P}_1(\mathcal{T}_h) \cap H_0^1(\Omega)$ .

We employ both structured and unstructured meshes  $\mathcal{T}_h$ . Our structured meshes are simply constructed by first building a Cartesian grid of squares, and then breaking each square into four triangles by connecting its vertices to its barycenter. The unstructured meshes are generated by the `mmg` software package [12]. We use both uniform and adapted unstructured meshes. In the uniform case, the meshes are simply generated using the `-hmax` flag of `mmg`, whereas the local mesh size is specified through a “metic file” *via* the `-sol` flag for adapted meshes.

For a given mesh  $\mathcal{T}_h$ ,  $N := \dim \mathcal{P}_1(\mathcal{T}_h) \cap H_0^1(\Omega)$  denotes the number of degrees of freedom of the associated finite element space. For a uniform mesh with maximal size  $h$ , we have  $N \sim (\text{diam}(\Omega)/h)^2$ .

For each configuration of domain and function, our results are summarized in a figure with 4 panels. Panels (a) and (b) aim at comparing the quality of approximation of different projectors. Specifically, the projection errors measured in the  $H^1$  seminorm and in the  $L^2$  norm are respectively represented on panels (a) and (b). On the other hand, in panels (c) and (d), we measure the quality of our guaranteed upper bounds, using that the local-best approximation error  $\|\nabla_h(u - \pi_h^1 u)\|_{\Omega}$  is known. On panel (c), we plot  $\|\nabla(u - \mathcal{J}_h^1 u)\|_{\Omega}$  and  $\|u - \mathcal{J}_h^1 u\|_{\Omega}$  together with the right-hand sides in the guaranteed estimations in (4.11) and (4.13), written as

$$\|\nabla(u - \mathcal{J}_h^1 u)\|_{\Omega} \leq (1 + (d+1)^2 c_{\Omega}^2)^{1/2} \|\nabla_h(u - \pi_h^1 u)\|_{\Omega} =: \eta_{H^1}(\pi_h^1 u)$$

and

$$\|u - \mathcal{J}_h^1 u\|_{\Omega} \leq \left( \frac{1}{\pi\sqrt{d+1}} + \frac{2}{d}\sqrt{d+1}c_{\Omega} \right) \left\{ \sum_{\alpha \in \mathcal{V}_h} h_{\alpha}^2 \|\nabla_h(u - \pi_h^1 u)\|_{\omega_{\alpha}}^2 \right\}^{1/2} =: \eta_{L^2}(\pi_h^1 u).$$

Finally, we plot the ratios on panel (d), thereby measuring the overestimation factor in our guaranteed bounds.

We obtain similar results in all the test cases below, so that we give a discussion in Section 6.5.

### 6.2. Smooth function

We start with the square  $\Omega := (-1, 1)^2$  and the smooth function

$$u(\mathbf{x}) = \sin(\pi \mathbf{x}_1) \sin(\pi \mathbf{x}_2).$$

We employ uniform unstructured meshes generated by `mmg`. The result are presented in Figure 1.




FIGURE 1. Smooth solution example.

### 6.3. Circular irregularity

We keep the domain  $\Omega := (-1, 1)^2$  and consider the function

$$u(\mathbf{x}) = (1 - |\mathbf{x}|)\mathbf{1}_{|\mathbf{x}| < 1}.$$

This function is  $C^0$ , but not  $C^1$ . In fact, the gradient has a line of discontinuity across the unit circle, so that  $h^{1/2}$  and  $h^{3/2}$  (i.e.,  $N^{-1/4}$  and  $N^{-3/4}$ ) rates are expected for the  $H^1$  and  $L^2$  errors. We employ uniform structured meshes. Figure 2 summarizes the results.




FIGURE 2. Circular interface solution example.

#### 6.4. Corner singularity

We now consider the  $L$ -shape domain  $\Omega := (-1, 1)^2 \setminus [0, 1]^2$ , and the standard corner singularity function

$$u(\mathbf{x}) = \chi(\mathbf{x})|\mathbf{x}|^\alpha \sin(\alpha\theta(\mathbf{x}))$$

where  $\alpha := 2/3$  and  $\theta(\mathbf{x})$  is an angular variable defined such that  $\theta = 0$  for  $\mathbf{x} \in (0, 1) \times \{0\}$  and  $3\pi/2$  when  $\mathbf{x} \in \{0\} \times (0, 1)$  and

$$\chi(\mathbf{x}) := (1 - |\mathbf{x}|^2)\mathbf{1}_{|\mathbf{x}| < 1}$$

is a cutoff function that enforces boundary conditions. We first use uniform meshes generated by `mmg` and present the corresponding results in Figure 3. In this case, we expect and observe convergence rates of order  $N^{-1/3}$  and  $N^{-5/6}$  in the  $H^1$  and  $L^2$  norms. We then employed adapted meshes (still generated by `mmg`) with corresponding results presented in Figure 4. Optimal  $h$ -convergence rates are expected and observed in this




FIGURE 3. Singular function example.

case. The adapted meshes are generated by requiring that  $h_K \lesssim \max(h_{\max}^{3/2}, |\mathbf{x}_K|^{1/3} h_{\max})$  for all  $K \in \mathcal{T}_h$ , where  $\mathbf{x}_K$  is the barycenter of  $K$ .

## 6.5. Discussion

In all the benchmarks, we obtain the expected rates of convergence for our quasi-interpolation operator  $\mathcal{J}_h^1$ . In the  $H^1$  seminorm, the errors of the global-best approximation  $u_h^1$ , the Lagrange interpolant  $\mathcal{I}_h^1 u$ , and our quasi-interpolant  $\mathcal{J}_h^1 u$  are always similar. In the  $L^2$  norm, our quasi-interpolant  $\mathcal{J}_h^1 u$  is always more accurate than the Lagrange interpolant  $\mathcal{I}_h^1 u$  and the global-best approximation  $u_h^1$ , and sometimes significantly so. In both the  $H^1$  seminorm and the  $L^2$  norm, the local-best approximation provides better errors than the conforming approximation, but the improvement is typically marginal. Our guaranteed upper bounds never underestimate the errors as predicted by the theory of Theorems 4.2 and 4.3, and behave with the correct rates. They allow for




FIGURE 4. Singular function example with adapted meshes.

error certification. The overestimation factor is remarkably stable in all tested situations, with values around 10 in the  $H^1$  seminorm and 100 in the  $L^2$  norm.

## DATA AVAILABILITY STATEMENT

No new data/codes were created or analyzed in this study.

## REFERENCES

- [1] R. Arcangeli and J.L. Gout, Sur l'évaluation de l'erreur d'interpolation de Lagrange dans un ouvert de  $\mathbb{R}^2$ . *RAIRO Analyse numérique* **10** (1976) 5–27.
- [2] M. Aurada, M. Feischl, J. Kemetmüller, M. Page and D. Praetorius, Each  $H^{1/2}$ -stable projection yields convergence and quasi-optimality of adaptive FEM with inhomogeneous Dirichlet data in  $\mathbb{R}^d$ . *ESAIM Math. Model. Numer. Anal.* **47** (2013) 1207–1235.

- [3] M. Bebendorf, A note on the Poincaré inequality for convex domains. *Z. Anal. Anwendungen* **22** (2003) 751–756.
- [4] S.C. Brenner, An optimal-order multigrid method for P1 nonconforming finite elements. *Math. Comp.* **52** (1989) 1–15.
- [5] C. Carstensen, Quasi-interpolation and a posteriori error analysis in finite element methods. *ESAIM Math. Model. Numer. Anal.* **33** (1999) 1187–1202.
- [6] C. Carstensen and S.A. Funken, Constants in Clément-interpolation error and residual based a posteriori error estimates in finite element methods. *East-West J. Numer. Math.* **8** (2000) 153–175.
- [7] C. Carstensen, J. Gedicke and D. Rim, Explicit error estimates for Courant, Crouzeix-Raviart and Raviart-Thomas finite element methods. *J. Comput. Math.* **30** (2012) 337–353.
- [8] C. Carstensen, D. Peterseim and M. Schedensack, Comparison results of finite element methods for the Poisson model problem. *SIAM J. Numer. Anal.* **50** (2012) 2803–2823.
- [9] T. Chaumont-Frelet and M. Vohralík, A stable local commuting projector and optimal  $hp$  approximation estimates in  $\mathbf{H}(\text{curl})$ . *Numer. Math.* **156** (2024) 2293–2342.
- [10] P.G. Ciarlet, The Finite Element Method For Elliptic Problems. SIAM (2002).
- [11] P. Clément, Approximation by finite element functions using local regularization. *ESAIM: Math. Model. Numer. Anal.* **9** (1975) 77–84.
- [12] C. Dobrzynski, MMG3D: User guide. Tech. Report 422, Inria (2012).
- [13] V. Dolejší, A. Ern and M. Vohralík,  $hp$ -adaptation driven by polynomial-degree-robust a posteriori error estimates for elliptic problems. *SIAM J. Sci. Comput.* **38** (2016) A3220–A3246.
- [14] A. Ern and J.-L. Guermond, Finite element quasi-interpolation and best approximation. *ESAIM: Math. Model. Numer. Anal.* **51** (2017) 1367–1385.
- [15] A. Ern and J.-L. Guermond, Finite Elements I. Approximation and Interpolation. *Texts in Applied Mathematics*. Vol. 72. Springer International Publishing, Springer Nature Switzerland AG (2021).
- [16] A. Ern and M. Vohralík, Polynomial-degree-robust a posteriori estimates in a unified setting for conforming, non-conforming, discontinuous Galerkin, and mixed discretizations. *SIAM J. Numer. Anal.* **53** (2015) 1058–1081.
- [17] A. Ern, I. Smears and M. Vohralík, Discrete  $p$ -robust  $\mathbf{H}(\text{div})$ -liftings and a posteriori estimates for elliptic problems with  $H^{-1}$  source terms. *Calcolo* **54** (2017) 1009–1025.
- [18] A. Ern, T. Gudi, I. Smears and M. Vohralík, Equivalence of local- and global-best approximations, a simple stable local commuting projector, and optimal  $hp$  approximation estimates in  $\mathbf{H}(\text{div})$ . *IMA J. Numer. Anal.* **42** (2022) 1023–1049.
- [19] R.S. Falk and R. Winther, Local bounded cochain projections. *Math. Comp.* **83** (2014) 2631–2656.
- [20] E. Gawlik, M.J. Holst and M.W. Licht, Local finite element approximation of Sobolev differential forms. *ESAIM: Math. Model. Numer. Anal.* **55** (2021) 2075–2099.
- [21] T. Hao, X. Guan, S. Mao and S. Chen, Computable interpolation error constants for the geometric simplex finite elements. *J. Sci. Comput.* **87** (2021) 21.
- [22] R. Hiptmair and C. Pechstein, Discrete regular decompositions of tetrahedral discrete 1-forms, in collection Maxwell’s Equations, De Gruyter (2019) 199–258.
- [23] M. Karkulik and J.M. Melenk, Local high-order regularization and applications to  $hp$ -methods. *Comput. Math. Appl.* **70** (2015) 1606–1639.
- [24] F. Kikuchi and X. Liu, Estimation of interpolation error constants for the  $P_0$  and  $P_1$  triangular finite elements. *Comput. Methods Appl. Mech. Eng.* **196** (2007) 3750–3758.
- [25] K. Kobayashi, On the interpolation constants over triangular elements. *App. Math.* **2015** (2015) 110–124.
- [26] K. Kobayashi and T. Tsuchiya, A Babuška-Aziz proof of the circumradius condition. *Jpn. J. Indust. Appl. Math.* **31** (2014) 193–210.
- [27] M.W. Licht, Averaging-based local projections in finite element exterior calculus. Preprint [arXiv:2301.03007](https://arxiv.org/abs/2301.03007) (2023).
- [28] X. Liu and F. Kikuchi, Analysis and estimation of error constants for  $P_0$  and  $P_1$  interpolations over triangular finite elements. *J. Math. Sci. Univ. Tokyo* **17** (2010) 27–78.
- [29] X. Liu and C. You, Explicit bound for quadratic Lagrange interpolation constant on triangular finite elements. *Appl. Math. Comput.* **319** (2018) 693–701.
- [30] A. Målqvist and D. Peterseim, Localization of elliptic multiscale problems. *Math. Comp.* **83** (2014) 2583–2603.
- [31] J.M. Melenk,  $hp$ -interpolation of nonsmooth functions and an application to  $hp$ -a posteriori error estimation. *SIAM J. Numer. Anal.* **43** (2005) 127–155.
- [32] J.C. Nédélec, Mixed finite elements in  $\mathbb{R}^3$ . *Numer. Math.* **35** (1980) 315–341.

- [33] P. Oswald, On a BPX-preconditioner for P1 elements. *Computing* **51** (1993) 125–133.
- [34] L.E. Payne and H.F. Weinberger, An optimal Poincaré inequality for convex domains. *Arch. Ration. Mech. Anal.* **5** (1960) 1005–1036.
- [35] P.A. Raviart and J.M. Thomas, A mixed finite element method for 2nd order elliptic problems, in *Mathematical Aspects of Finite Element Methods*. Springer-Verlag (1977).
- [36] J. Schöberl, Commuting quasi-interpolation operators for mixed finite elements. Tech. Report ISC-01-10-MATH, Texas A&M University (2001).
- [37] J. Schöberl, J.M. Melenk, C. Pechstein and S. Zaglmayr, Additive Schwarz preconditioning for  $p$ -version triangular and tetrahedral finite elements. *IMA J. Numer. Anal.* **28** (2008) 1–24.
- [38] L.R. Scott and S. Zhang, Finite element interpolation of nonsmooth functions satisfying boundary conditions. *Math. Comp.* **54** (1990) 483–493.
- [39] A. Veerer, Approximating gradients with continuous piecewise polynomial functions. *Found. Comput. Math.* **16** (2016) 723–750.
- [40] A. Veerer and R. Verfürth, Explicit upper bounds for dual norms of residuals. *SIAM J. Numer. Anal.* **47** (2009) 2387–2405.
- [41] A. Veerer and R. Verfürth, Poincaré constants for finite element stars. *IMA J. Numer. Anal.* **32** (2012) 30–47.
- [42] R. Verfürth, Error estimates for some quasi-interpolation operators. *ESAIM: Math. Model. Numer. Anal.* **33** (1999) 695–713.
- [43] M. Vohralík, Analysis and approximation of partial differential equations by finite elements. Lecture notes, ENSTA, Institut Polytechnique de Paris (2025).
- [44] M. Vohralík,  $p$ -robust equivalence of global continuous and local discontinuous approximation, a  $p$ -stable local projector, and optimal elementwise  $hp$  approximation estimates in  $H^1$ . HAL Preprint 04436063, submitted for publication (2025).



**Please help to maintain this journal in open access!**

This journal is currently published in open access under the Subscribe to Open model (S2O). We are thankful to our subscribers and supporters for making it possible to publish this journal in open access in the current year, free of charge for authors and readers.

Check with your library that it subscribes to the journal, or consider making a personal donation to the S2O programme by contacting [subscribers@edpsciences.org](mailto:subscribers@edpsciences.org).

More information, including a list of supporters and financial transparency reports, is available at <https://edpsciences.org/en/subscribe-to-open-s2o>.